

Question 1 (12 marks) Start a NEW booklet.**Marks**

- a) Find the remainder when the polynomial $P(x) = 2x^3 - 4x + 3$ is divided by $(x+1)$. 1
- b) Using the points $A(-4,1)$ and $B(2,-2)$ find the coordinates of the point $P(x,y)$ that divides the interval AB **externally** in the ratio $5:2$. 2
- c) Solve $\frac{2}{x-1} \leq 3$. 3
- d) i) Show that $f(x) = \frac{x^3}{1+x^2}$ is an odd function. 1
- ii) Hence or otherwise evaluate $\int_{-1}^1 \frac{x^3}{1+x^2} dx$. 1
- e) Neatly sketch $y = 3\cos^{-1} 2x$ clearly showing the domain and range. 2
- f) Show that $\frac{d}{dx}(\sec x) = \sec x \tan x$. 2

Question 2 (12 marks) Start a NEW booklet.**Marks**

- a) Find $\int \frac{1}{\sqrt{3-x^2}} dx$. 1
- b) The parametric equations of a curve are given by $x = 6r$, $y = \frac{3}{r}$.
Find the cartesian equation of the curve. 1
- c) Given the function $f : y = \frac{x-1}{x+2}$, find its inverse function $f^{-1}(x)$ in terms of x and **state** the range of the inverse function. 3
- d) Solve $\sin 2\theta = \sin \theta$ for $0 \leq \theta \leq 2\pi$. 3
- e) If α, β and γ are the roots of $2x^3 + 12x^2 - 6x + 1 = 0$, find
- i) $\alpha + \beta + \gamma$ 1
- ii) $\alpha\beta\gamma$ 1
- iii) $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$ 2

Question 3 (12 marks) Start a NEW booklet.**Marks**


- a) Find $\lim_{\theta \rightarrow 0} \frac{\sin 2\theta}{\tan 3\theta}$. 2
- b) Use the substitution $u = 5 - x$ to evaluate $\int_1^5 x\sqrt{5-x} \, dx$ 3
- c) i) Express $\cos\theta - \sqrt{3}\sin\theta$ in the form $R\cos(\theta + \alpha)$, 2
 where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$.
- ii) Hence solve $\cos\theta - \sqrt{3}\sin\theta = 2$ for $0 \leq \theta \leq 2\pi$. 2
- d) A particle is moving in a straight line x centimetres from the origin O.
 After t minutes its displacement is given by $x = 3 - 5\cos 2t$.
- i) Show that its acceleration is given by $\ddot{x} = -4(x - 3)$. 2
- ii) Assuming it is moving in simple harmonic motion find its centre of motion. 1

Question 4 (12 marks) Start a NEW booklet.**Marks**

- a) The acceleration of a body P is given by $\frac{d^2x}{dt^2} = 18x^3 + 18x$, where x is the displacement of P from O at time t . The velocity is v .

Given that $t = 0, x = 0, v = 3$ and that $v > 0$ throughout the motion:

- i) Find v in terms of x . 2
 - ii) Show that $x = \tan 3t$. 2
- b) $P(2ap, ap^2)$ is a point on the parabola $x^2 = 4ay$. SN is perpendicular to the normal at P , where S is the focus of the parabola and N the foot of the perpendicular from S to the normal.




- i) Show that the equation of the normal at P is $x + py = 2ap + ap^3$. 2
- ii) Find the equation of SN . 2
- iii) Show that the coordinates of the point N are $(ap, ap^2 + a)$. 2
- iv) Find the locus of N as P moves on the parabola. 2

Question 5 (12 marks) Start a NEW booklet.**Marks**

- a) i) Find the values of a and b such that $x^2 - 4x + 7 = (x - a)^2 + b$. 1
- ii) Hence, state the largest positive domain for which $y = x^2 - 4x + 7$ has an inverse function. 1

- b) Evaluate $\sin\left(\tan^{-1}\frac{1}{2}\right)$ in exact form. 2

- c) The altitudes PM and QN of an acute angled triangle PQR meet at H . PM produced cuts the circle PQR at A . [A **larger diagram** is included at the end of the exam paper, **remove** it, use it and **submit** it with your **solutions**]



- i) Explain why $PQMN$ is a cyclic quadrilateral. 1
- ii) With the aid of congruent triangles prove that $HM = MA$. 3
- d) The rate of growth of the number of bacteria in a colony is proportional to the excess of the colony's population over 5000 and is given by $\frac{dN}{dt} = k(N - 5000)$.
- i) Show that $N = 5000 + Ae^{kt}$ is a solution to the differential equation. 1
- ii) If the initial population is 15 000 and reaches 20 000 after 2 days find the values of A and k . 2
- iii) Hence, calculate the expected population after 7 days. 1

Question 6 (12 marks) Start a NEW booklet.**Marks**

a) Using $t = \tan \frac{\theta}{2}$, prove that $\frac{1 + \cos \theta}{1 - \cos \theta} = \cot^2 \frac{\theta}{2}$. 2

b) Air is being pumped into a spherical balloon at the rate of $20 \text{ cm}^3 / \text{s}$. 4

Find the rate of increase of the balloon's surface area when the radius is 5cm.

$$\left[V = \frac{4}{3} \pi r^3, SA = 4\pi r^2 \right]$$

c) For the function $y = \frac{x^2 - 4}{x^2 - 1}$

i) Write down the equations of any horizontal and vertical asymptotes. 2

ii) Find any stationary points and determine their nature. 2

iii) Neatly sketch the graph showing the above features. 2

Question 7 (12 marks) Start a NEW booklet.**Marks**


- a) Prove by mathematical induction that for any positive integer $n \geq 1$.

4

$$\frac{1}{1 \times 5} + \frac{1}{5 \times 9} + \frac{1}{9 \times 13} + \dots + \frac{1}{(4n-3)(4n+1)} = \frac{n}{4n+1}$$

- b) From a balloon 500 metres above a road junction X , the angle of depression to a point P , which lies due south of the junction X is 42° .

To a point Q , which lies at a bearing of 280° from the road junction, the angle of depression to the balloon is 32° .




- i) Clearly explain why $\angle QXP = 100^\circ$.
- ii) Calculate the distance PQ (correct to the nearest metre).
- c) A particle is moving in a straight line with simple harmonic motion. The velocity of the particle is respectively $\sqrt{20} \text{ ms}^{-1}$ and 4 ms^{-1} at distances of 1 metre and 2 metres from the centre of motion. Find the period and amplitude of the motion.


1

3

4

END OF EXAM

This image shows a single page of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page, leaving small margins at the top and bottom. There are no vertical margin lines, text, or other markings on the page.

CHS YEAR 12 EXTENSION 1 2011	TRIAL HSC SOLUTIONS
1a) $P -1 = -2 + 4 + 3 = 5$ ✓ b) $x = \frac{5 \times 2 + -2 \times -4}{5 - 2}$ $y = \frac{5 \times -2 + -2 \times 1}{5 - 2}$ ✓ $= 6$ $= -4$ ✓ $\therefore P_{x,y} = 6, -4$ c) CV1: $2 = 3x - 1 \rightarrow x = \frac{5}{3}$ ✓ CV2: $x = 1$ ✓ On testing $x < 1$ and $x \geq \frac{5}{3}$, since $x \neq 1$ ✓ d) i) An odd function exists if $f(-x) = -f(x)$ $f(-x) = \frac{-x^3}{1 + (-x)^2}$ $= \frac{-x^3}{1 + x^2} = -f(x)$ $\therefore$ an odd function. ✓ ii) $\therefore \int_{-1}^1 \frac{x^3}{1 + x^2} dx = 0$ ✓ an odd function with symmetrical limits. e)  ✓✓	f) $\frac{d}{dx} \sec x = \frac{d}{dx} \cos x^{-1}$ $= -\cos x^{-2} \times -\sin x$ $= \frac{\sin x}{\cos^2 x}$ $= \frac{\sin x}{\cos x} \times \frac{1}{\cos x} = \tan x \sec x$ ✓ 2a) $\int \frac{1}{\sqrt{3-x^2}} dx = \sin^{-1}\left(\frac{x}{\sqrt{3}}\right) + c$ ✓ b) $x = 6r \rightarrow r = \frac{x}{6}$ $\therefore y = \frac{3}{r} \rightarrow y = \frac{3}{x/6}$ Hence $xy = 18$ c) $f^{-1}(x) : x = \frac{y-1}{y+2}$ ✓ $xy + 2x = y - 1$ $2x + 1 = y - xy$ $2x + 1 = y(1 - x)$ $\therefore y = \frac{2x + 1}{1 - x}$ ✓ Range is all real y except $y = -2$ ✓ d) $\sin 2\theta = \sin \theta$ for $0 \leq \theta \leq 2\pi$ $2\sin \theta \cos \theta - \sin \theta = 0$ $2\sin \theta (2\cos \theta - 1) = 0$ $\therefore \sin \theta = 0 \text{ or } \cos \theta = \frac{1}{2}$ ✓ $\therefore \theta = 0, \pi, 2\pi \text{ or } \theta = \frac{\pi}{3}, \frac{5\pi}{3}$ ✓

$$\text{e) i) } \alpha + \beta + \gamma = \frac{-b}{a} = -6$$

✓

$$\text{ii) } \alpha\beta\gamma = \frac{d}{a} = \frac{-1}{2}$$

✓

$$\text{iii) } \frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\beta\gamma + \alpha\gamma + \alpha\beta}{\alpha\beta\gamma}$$

✓

$$= \frac{-6/2}{-1/2} = 6$$

✓

$$\text{3a) } \lim_{\theta \rightarrow 0} \frac{\sin 2\theta}{\tan 3\theta} = \lim_{\theta \rightarrow 0} \frac{2}{3} \left[\frac{\sin 2\theta}{2\theta} \times \frac{3\theta}{\tan 3\theta} \right]$$

✓

$$= \frac{2}{3}$$

✓

$$\text{b) } \text{When } u = 5 - x \rightarrow x = 5 - u$$

$$\text{When } x = 1, u = 4; \quad x = 5, u = 0.$$

$$du = -dx$$

$$I = - \int_4^0 5 - u \sqrt{u} \, du$$

✓

$$= \int_0^4 5u^{\frac{1}{2}} - u^{\frac{3}{2}} \, du$$

$$= \left[\frac{10u^{\frac{3}{2}}}{3} - \frac{2u^{\frac{5}{2}}}{5} \right]_0^4$$

✓

$$= \left[\frac{10 \cdot 4^{\frac{3}{2}}}{3} - \frac{2 \cdot 4^{\frac{5}{2}}}{5} \right] - 0$$

$$= \frac{208}{15}$$

✓

$$\text{c) i) } \cos\theta - \sqrt{3}\sin\theta = R\cos\theta + \alpha$$

$$= R\cos\alpha\cos\theta - R\sin\alpha\sin\theta$$

$$\therefore R\cos\alpha = 1 \text{ and } R\sin\alpha = \sqrt{3}$$

✓

$$\therefore R^2\cos^2\alpha + R^2\sin^2\alpha = 1 + 3$$

$$\therefore R^2 = 4 \rightarrow R = 2$$

$$\text{and hence } \alpha = \frac{\pi}{3}$$

✓

$$\therefore \cos\theta - \sqrt{3}\sin\theta = 2\cos\left(\theta + \frac{\pi}{3}\right)$$

$$\text{ii) } \therefore 2\cos\left(\theta + \frac{\pi}{3}\right) = 2$$

$$\therefore \cos\left(\theta + \frac{\pi}{3}\right) = 1$$

✓

$$\therefore \theta + \frac{\pi}{3} = 0, 2\pi \rightarrow \theta = \frac{5\pi}{3} \text{ only.}$$

✓

$$\text{d) i) } x = 3 - 5\cos 2t$$

$$\therefore \dot{x} = 10\sin 2t$$

$$\ddot{x} = 20\cos 2t$$

✓

$$\ddot{x} = 20\left(\frac{3-x}{5}\right) = 4(3-x)$$

✓

$$\therefore \ddot{x} = -4(x-3)$$

$$\text{ii) } \text{Centre of motion is at } x = 3.$$

✓

$$\text{4a) i) } \frac{d}{dx}\left(\frac{1}{2}v^2\right) = 18x^3 + 18x$$

$$\therefore \frac{1}{2}v^2 = \frac{9x^4}{2} + 9x^2 + c$$

$$v^2 = 9x^4 + 18x^2 + d$$

$$\text{when } x = 0, v = 3$$

$$\therefore 9 = 0 + 0 + d \rightarrow d = 9$$

✓

$$v^2 = 9x^4 + 18x^2 + 9$$

$$= 9x^4 + 2x^2 + 1$$


$$= 9x^2 + 1^2$$

$$\therefore v = 3x^2 + 1, \text{ since } v > 0.$$

✓

$$\text{ii) } \frac{dx}{dt} = 3x^2 + 1$$

$$\therefore \frac{dt}{dx} = \frac{1}{3x^2 + 1}$$

$\therefore t = \frac{1}{3}\tan^{-1}x + c$	✓	5a) i) $x^2 - 4x + 7 = x - a^2 + b$ $x^2 - 4x + 4 + 3 = x - a^2 + b$ $x - 2^2 + 3 = x - a^2 + b$ $\therefore a = 2, b = 3$	✓
$\therefore 0 = \frac{1}{3}\tan^{-1}0 + c \rightarrow c = 0$		ii) all $x \geq 2$	✓
$\therefore t = \frac{1}{3}\tan^{-1}x$		b) Let $u = \tan^{-1}\frac{1}{2}$ and evaluate $\sin u$	
$\therefore 3t = \tan^{-1}x \rightarrow x = \tan 3t$	✓	$u = \tan^{-1}\frac{1}{2} \rightarrow \tan u = \frac{1}{2}$	✓
b) i) $x^2 = 4ay \rightarrow y = \frac{x^2}{4a}$		 $\sin u = \frac{1}{\sqrt{5}}$	✓
$\therefore \frac{dy}{dx} = \frac{x}{2a} \rightarrow m_1 = \frac{2ap}{2a} = p$		c) i) $\angle QMP = \angle PNQ$ { both 90° } $\therefore PQMN$ is a cyclic quad - $\angle$'s in same segment standing on same arc or chord PQ .	✓
$\therefore \text{gradient of the normal at } P \text{ is } m_2 = \frac{-1}{p}$	✓	ii) Join QA : $\angle AQR = \angle APR$ { $= \theta$ say } { $\angle$'s in same segment on arc AR } Also $\angle MQN = \angle MPN$ { $= \theta$ } { $\angle$'s in same segment on arc MN }	✓
$\therefore \text{Equation of the normal:}$		$\therefore$ in Δ 's QHM and QAM	
$y - ap^2 = \frac{-1}{p} x - 2ap$	✓	$\angle QMH = \angle QMA$ { both 90° } $\angle HQM = \angle AQM$ { proven above ($= \theta$) } $QM = QM$ { common }	✓
$py - ap^3 = -x + 2ap$		$\therefore \Delta QHM \equiv \Delta QAM$ (AAS)	
$\therefore x + py = 2ap + ap^3$		Hence $HM = MA$ { corresponding sides in $\equiv \Delta$'s }	✓
ii) gradient of SN is p and S has coordinates $(0, a)$	✓	d) i) $N = 5000 + Ae^{kt} \rightarrow Ae^{kt} = N - 5000$	
$\therefore \text{eq}^n \text{ of } SN \text{ is } y - a = p(x - 0) \rightarrow y = px + a$	✓	$LHS = \frac{d}{dt} 5000 + Ae^{kt}$	
iii) $x + py = 2ap + ap^3$		$= k Ae^{kt}$	✓
$y = px + a$		$= k N - 5000 = RHS$	
$\therefore [2] \text{ in } [1] \rightarrow x + p \quad px + a = 2ap + ap^3$		ii) $t = 0 \rightarrow N = 15\,000$	
$\therefore x + p^2x + ap = 2ap + ap^3$		$15\,000 = 5000 + Ae^0 \rightarrow A = 10\,000$	✓
$\therefore x(1 + p^2) = ap(1 + p^2)$	✓	$t = 2 \rightarrow N = 20\,000$	
$\therefore x = ap$	✓	$20\,000 = 5000 + 10\,000e^{2k}$	
and $y = ap^2 + a$	✓		
$\therefore N \quad ap, ap^2 + a$			
iv) $x = ap \rightarrow p = \frac{x}{a}$	✓		
$\therefore y = a\left(\frac{x}{a}\right)^2 + a \rightarrow y = \frac{x^2}{a} + a$	✓		
$x^2 = a(y - a)$			

$$\therefore e^{2k} = \frac{3}{2} \rightarrow 2k = \ln \frac{3}{2}$$

$$k = \frac{1}{2} \ln \frac{3}{2} \approx 0.202733$$

iii) When $t = 7$, $N = 5000 + 10\,000e^{7k}$

$$\approx 46\,335$$

6a) Using $t = \tan \frac{\theta}{2}$:

$$LHS = \frac{1 + \frac{1-t^2}{1+t^2}}{1 - \frac{1-t^2}{1+t^2}}$$

$$= \frac{1+t^2 + 1-t^2}{1+t^2 - 1-t^2}$$

$$= \frac{2}{2t^2} = \frac{1}{t^2}$$

$$= \frac{1}{\tan^2 \frac{\theta}{2}} = \cot^2 \frac{\theta}{2}$$

b) $\frac{dV}{dt} = 20$, $V = \frac{4}{3}\pi r^3$, $\frac{dV}{dr} = \pi r^2$, $A = 4\pi r^2$, $\frac{dA}{dr} = 8\pi r$

$$\frac{dV}{dt} = \frac{dV}{dr} \times \frac{dr}{dt} \rightarrow 20 = 4\pi r^2 \times \frac{dr}{dt}$$

$$\therefore \frac{dr}{dt} = \frac{5}{\pi r^2}$$

Also $\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$

$$= 8\pi r \times \frac{5}{\pi r^2} = \frac{40}{r}$$

and when $r = 5$, $\frac{dA}{dt} = 8\text{ cm}^2 / \text{s}$.

c) i) Vertical asymptotes: $x = \pm 1$

Horizontal asymptote: $y = 1$

ii) $\frac{dy}{dx} = \frac{x^2 - 1 \cdot 2x - x^2 - 4 \cdot 2x}{x^2 - 1^2}$

$$= \frac{6x}{x^2 - 1^2}$$

Stationary points when $\frac{dy}{dx} = 0$


$\therefore$ when $6x = 0 \rightarrow x = 0$

at $x = 0$, $y = 4$

x	$-\frac{1}{2}$	0	$\frac{1}{2}$
y'	-ve	0	+ve

$\therefore$ local minimum turning point at $(0, 4)$

iii)



$$7a) \frac{1}{1 \times 5} + \frac{1}{5 \times 9} + \frac{1}{9 \times 13} + \dots + \frac{1}{4n-3} \frac{1}{4n+1} = \frac{n}{4n+1}$$

$$\text{When } n = 1, LHS = \frac{1}{1 \times 5} = \frac{1}{5}; RHS = \frac{1}{4 \times 1 + 1} = \frac{1}{5}$$

$\therefore$ true for $n = 1$. ☒

Assume true for $n = k$

$$\text{i.e. } \frac{1}{1 \times 5} + \frac{1}{5 \times 9} + \frac{1}{9 \times 13} + \dots + \frac{1}{4k-3} \frac{1}{4k+1} = \frac{k}{4k+1}$$

Prove true for $n = k + 1$

$$\text{i.e. } S_k + T_{k+1} = S_{k+1}$$

$$S_k = \frac{k}{4k+1}, T_{k+1} = \frac{1}{4k+1} \frac{1}{4k+5}, S_{k+1} = \frac{k+1}{4k+5} \quad \checkmark$$

$$LHS = S_k + T_{k+1}$$

$$LHS = \frac{k}{4k+1} + \frac{1}{4k+1} \frac{1}{4k+5}$$

$$= \frac{k}{4k+1} \frac{4k+5}{4k+5} + \frac{1}{4k+1} \frac{1}{4k+5} \quad \checkmark$$

$$= \frac{4k^2 + 5k + 1}{4k+1} \frac{1}{4k+5}$$

$$= \frac{4k+1}{4k+1} \frac{k+1}{4k+5}$$

$$= \frac{k+1}{4k+5} = S_{k+1} = RHS \quad \checkmark$$


$\therefore$ If true for $n = k$, then true for $n = k + 1$.

Hence by the principle of mathematical induction,
the result is true for all $n \geq 1$.

b) i) P is due south of X , i.e. at a bearing of 180° .
 Q is at a bearing of 280° .


$\therefore$ the angle between P and Q on the ground is
 $280^\circ - 180^\circ = 100^\circ$. ☒

ii)



$$\tan 48^\circ = \frac{PX}{500}$$

$$\therefore PX = 500 \tan 48^\circ \quad \checkmark$$



$$\tan 58^\circ = \frac{QX}{500}$$

$$\therefore QX = 500 \tan 58^\circ$$

$\therefore$ using ΔQXP :

$$PQ^2 = PX^2 + QX^2 - 2 \cdot PX \cdot QX \cdot \cos 100^\circ \quad \checkmark$$

$$= 500^2 \tan^2 48^\circ + 500^2 \tan^2 58^\circ - 2 \times 500 \tan 42^\circ \times 500 \tan 58^\circ \times \cos 100^\circ$$

$$\therefore PQ^2 = 1\,102\,949.62 \rightarrow PQ \approx 1050\,m \quad \checkmark$$

c) $v^2 = n^2 a^2 - x^2$; $v = \sqrt{20}$, $x = 1$; $v = 4$, $x = 2$.

$$20 = n^2 a^2 - 1 \quad \text{--- [1]} \quad \checkmark$$

$$16 = n^2 a^2 - 4 \quad \text{--- [2]} \quad \checkmark$$

$$\text{[1]} \div \text{[2]} \rightarrow \frac{5}{4} = \frac{a^2 - 1}{a^2 - 4} \quad \checkmark$$

$$\therefore 5a^2 - 20 = 4a^2 - 4$$

$$\therefore a^2 = 16 \rightarrow a = 4$$

$$\therefore \frac{16}{12} = \frac{4}{3} = n^2$$

$$\therefore n = \frac{2}{\sqrt{3}}$$

$$\text{hence the period} = \frac{2\pi}{n} = \frac{2\pi}{\frac{2}{\sqrt{3}}} = \pi\sqrt{3} \text{ seconds} \quad \checkmark$$

$$\text{amplitude} = 4 \text{ metres.} \quad \checkmark$$