



Caringbah High School

2013

Trial HSC Examination

Mathematics Extension I

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen (Black pen is preferred)
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11–14, show relevant mathematical reasoning and/or calculations

Total marks – 70

Section I Pages 2 – 4

10 marks

- Attempt Questions 1–10
- Allow about 15 minutes for this section

Section II Pages 5 – 8

60 marks

- Attempt Questions 11–14
- Allow about 1 hour and 45 minutes for this section

Section I


10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

For Questions 1–10, use the multiple-choice answer sheet on page 9. Please detach this from the exam paper and submit with your answer booklets.

1 This is the graph of $y = f(x)$, $0 \leq x \leq 8$.



The value of $\int_0^8 f(x) dx$ is

- A) $\frac{\pi}{4} - 2$ B) $\pi - 2$
C) $\frac{\pi}{4} + 10$ D) $10 + \pi$
- 2 Let $f(x) = 3\cos^{-1}\left(\frac{x}{2}\right)$.
The domain of the function $f(x)$ is given by
- A) $-\frac{1}{3} \leq x \leq \frac{1}{3}$ B) $-\frac{1}{2} \leq x \leq \frac{1}{2}$
C) $-2 \leq x \leq 2$ D) $-3 \leq x \leq 3$
- 3 The point $(3, -4)$ divides the interval AB externally in the ratio 3:2. If the coordinates of A are $(6, 5)$, then the coordinates of B are
- A) $(5, 2)$ B) $(8, 11)$
C) $(4, -1)$ D) $(1, -10)$

4

$$\frac{d(\tan^{-1} 3x)}{dx} =$$

A) $\frac{3}{9 + x^2}$

B) $\frac{1}{9 + x^2}$

C) $\frac{1}{1 + 9x^2}$

D) $\frac{3}{1 + 9x^2}$

5

The variable point P ($5t, t^2$) lies on a parabola. The Cartesian equation for this parabola is

A) $y = \frac{x^2}{4}$

B) $x^2 = 10y$

C) $y = 25x^2$

D) $x^2 = 25y$

6

α, β and γ are roots of the equation $x^3 - 3x^2 + 1 = 0$.

The value of $\alpha\beta + \alpha\gamma + \beta\gamma$ is

A) -1

B) 0

C) 1

D) 3

7

A particle undergoes SHM about the origin. Its displacement in cm is given by

$$x = 3 \cos\left(2t + \frac{\pi}{3}\right).$$

The particle is at rest when

A) $x = -3$

B) $x = 0$

C) $t = \frac{\pi}{6}$

D) $t = 0$

8

$$\int \frac{-dx}{\sqrt{9 - x^2}}$$

Which of the following may be a solution?

A) $\cos^{-1} \frac{x}{3}$

B) $\sin^{-1} \frac{x}{3}$

C) $\cos^{-1} 3x$

D) $\sin^{-1} 3x$

- 4

Section II

60 marks

Attempt Questions 11–14

Allow about 1 hour and 45 minutes for this section

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11–14, your responses should include relevant mathematical reasoning and/or calculations.

Question 11 (15 marks) Use a SEPARATE writing booklet.

- a) Evaluate $\lim_{x \rightarrow \infty} \frac{3x + 2}{5 - x}$ 1
- b) Differentiate $\sin^{-1}(x^2)$, with respect to x . 2
- c) Sketch the graph of $y = 3 \sin\left(\frac{x}{2}\right)$ in the domain $0 \leq x \leq 2\pi$. 2
- d) Using the substitution $u = 4 - x^3$, 3
Evaluate $\int_{-1}^1 x^2 \sqrt{4 - x^3} dx$
- e) The line $y = 2x - 3$ intersects with the curve $y = 2x^3 - 15$ at the point 3
(2,1). Find the size of the angle between the line and the curve at the
point of intersection. (Answer to nearest degree)
- f) Find all values for x that satisfy $\frac{5}{x - 4} \leq x$ 2
- g) The function $f(x) = x^2 - e^x$ has a root near $x = 3$. Use one application 2
of Newton's method to find a better approximation.

Question 12 (15 marks) Use a SEPARATE writing booklet.

a) Evaluate $\lim_{x \rightarrow 0} \frac{3x}{\sin 2x}$ 1

b) Find the general solution to 2

$$\sqrt{3} \tan x - 1 = 0$$

Answer as an exact value in radians.

c) A pendulum swings freely due to gravity and is friction free. When viewed from above, the end of the pendulum executes simple harmonic motion, with a period of π seconds and an amplitude of 1.2 m.

i) Explain why the acceleration, $\ddot{x}$, of the pendulum is given by $\ddot{x} = -4x$, where x is the position at any time, t . 1

ii) Using part i), show that the maximum velocity of the end of the pendulum is $2 \cdot 4 \text{ ms}^{-1}$. 2

d) i) $x^2 + 8x + 20$ can be expressed in the form $(x + a)^2 + b^2$. 2
Find values for a and b .

ii) Hence or otherwise find 2

$$\int \frac{1}{x^2 + 8x + 20} dx$$

e) A spherical beach ball is being inflated at a rate of 12 mm^3 per second. 2
Calculate the rate that the radius is increasing when the surface area is $5\,000 \text{ mm}^2$. (NB. $V = \frac{4}{3}\pi r^3$ and $S = 4\pi r^2$)

f) The half-life of a substance is the time taken for half of the substance to decay. The carbon isotope ^{14}C decays at a rate proportional to its mass. It has been shown that ^{14}C has a half-life of 5580 years. 3

A fossil that was tested contained 40% of the ^{14}C it would have originally contained.

Estimate the age of the fossil.

Question 13 (15 marks) Use a SEPARATE writing booklet.

a) Solve, $\cos 2x = \sqrt{3} \cos x - 1$, for $0 \leq \theta \leq \frac{\pi}{2}$. 2

b) i) Differentiate $x \sin^{-1}(x) + \sqrt{1-x^2}$ 2

ii) Hence evaluate $\int_0^1 \sin^{-1} x \, dx$ 2

c) The 2 points P ($2ap, ap^2$) and Q ($2aq, aq^2$) lie on the parabola $x^2 = 4ay$, with $a > 0$.

The chord PQ passes through the focus of the parabola.


i) Show that $pq = -1$. 1

ii) The tangent at P has the equation $y = px - ap^2$ 2

The tangents from P and Q intersect at T. Show that T lies on the directrix, $y = -a$.

d) Find all values of θ , $0 \leq \theta \leq \pi$ such that $\sqrt{2} \sin \theta + \cos \theta = 1$ 3

e) Two circles with centres P and Q touch externally at C and have a common tangent that touches at A and B, as shown. The common tangent at C meets AB in T.



i) Show that T is the midpoint of AB. 1

ii) Show that C, T, A and P are concyclic. 2

Question 14 (15 marks) Use a SEPARATE writing booklet.

a) Find $\int \sin^2 x \, dx$ 2

b) Let $f(x) = \frac{1}{1+x^3}$ for all x . 1

Find an expression for the inverse function $f^{-1}(x)$, in terms of x .

c) i) Sketch the curve $y = \cos^{-1} x$ 1

ii) The area between the curve $y = \cos^{-1} x$, the line $x = -1$ and the x -axis is rotated about the x -axis. Use Simpson's rule with 5 function values to approximate the volume of the solid formed. 3

d) Prove by mathematical induction that 3

$$7^{2n} - 3^{3n} \text{ is divisible by } 11, \text{ for all integers } n \geq 1.$$

e) If an archer fires an arrow with a velocity of 50 ms^{-1} at an angle of θ to the horizontal, it can be shown that the equations of motion are given by


$$x = 50t \cos \theta \quad \text{and} \quad y = \frac{-gt^2}{2} + 50t \sin \theta \quad (\text{do not prove this}).$$

i) Show the Cartesian equation for the flight of the arrow is given by 2

$$y = x \tan \theta - \frac{gx^2}{5000} \sec^2 \theta$$

ii) In the 1992 Olympic Games in Barcelona, paralympian Antonio Rebello lit the Olympic cauldron in a most unique manner. From a horizontal distance of 60 metres from the base of the cauldron he fired a lit arrow across the top of the cauldron. The top of the cauldron was 21 metres higher than him. He had to shoot the arrow to within 2 metres above the cauldron to ignite the rising gas.

Using $g = 10$, find the range of angles from the horizontal that Antonio Rebello could aim through to successfully light the Olympic flame. 3



Name: _____

Multiple Choice Answer Sheet

Sample: $2 + 4 =$ (A) 2 (B) 6 (C) 8 (D) 9
A ☐ B ☒ C ☐ D ☐

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A  B  C  D 

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word **correct** and drawing an arrow as follows.

A  B  C  D 

correct

- | | | | | | | | | |
|-----|---|-----------------------|---|-----------------------|---|-----------------------|---|-----------------------|
| 1) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 2) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 3) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 4) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 5) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 6) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 7) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 8) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 9) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |
| 10) | A | ☐ | B | ☐ | C | ☐ | D | ☐ |

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$

013 CHS MATHS EXT 1 TRIAL HSC

Section 1

- 1) B 2) C 3) A 4) D 5) D
- 6) B 7) A 8) A 9) A 10) C


Section 2

Question 11

- a) -3
- b) 2x

$$\sqrt{1-x^4}$$

- c) $\frac{4}{3}$



- d) $u = 4 - x^3$ $u = 1$ $u = 3$

$$du = -3x^2 dx$$

$$\int_{-1}^1 x^2 \sqrt{4-x^3} dx = \int_3^1 \sqrt{u} \cdot \frac{-du}{3}$$

$$= -\frac{1}{3} \int_3^1 u^{\frac{1}{2}} du$$

$$= -\frac{1}{3} \left[\frac{2}{3} u^{\frac{3}{2}} \right]_3^1$$

$$= -\frac{2}{9} \left[\sqrt{3} - \sqrt{27} \right]$$

$$= \frac{2}{9} \left[\sqrt{27} - \sqrt{3} \right]$$

$$= \frac{2}{9} \left[3\sqrt{3} - \sqrt{3} \right]$$

$$= \frac{2}{9} \left[2\sqrt{3} \right]$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

- f) critical values

$$x=4$$

$$x=5$$

$$x=6$$

$$x=7$$

$$x=8$$

$$x=9$$

$$x=10$$

$$x=11$$

$$x=12$$

$$x=13$$

$$x=14$$

$$x=15$$

$$x=16$$

$$x=17$$

$$x=18$$

$$x=19$$

$$x=20$$

$$x=21$$

$$x=22$$

$$x=23$$

$$x=24$$

$$x=25$$

$$x=26$$

$$x=27$$

$$x=28$$

$$x=29$$

$$x=30$$

$$x=31$$

$$x=32$$

$$x=33$$

$$x=34$$

$$M = M_0 e^{-kt}$$

$$\frac{1}{2} M_0 = M_0 e^{-kt}$$

$$k = \frac{\ln(2)}{5580}$$

$$k \approx 1.2422 \times 10^{-4}$$

$$0.4 M_0 = M_0 e^{-kt}$$

$$t = \frac{\ln(0.4)}{k}$$

$$t \approx 7376 \text{ years}$$

$$t \approx 7376 \text{ years}$$

Question 13

$$a) \cos 2x = 2 \cos^2 x - 1$$

$$\text{so } 2 \cos^2 x - 1 = \sqrt{3} \cos x - 1$$

$$2 \cos^2 x - \sqrt{3} \cos x = 0$$

$$\cos x (2 \cos x - \sqrt{3}) = 0$$

$$\cos x = 0 \text{ or } \cos x = \frac{\sqrt{3}}{2}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$i) y = p^2 - ap^2$$

$$x = \frac{y + ap^2}{p}$$

$$i) \frac{y + ap^2}{p} = \frac{y + ap^2}{p}$$

$$y + ap^2 = py + ap^2$$

$$py - ap^2 = py - pa$$

$$py - pa = py - pa$$

$$y(p-p) = -a(p-p)$$

$$y = -a$$

$$y = -a$$

$$d) \sqrt{2} \sin \theta + \cos \theta = R \sin(\theta + \alpha)$$

$$R \cos \alpha = \sqrt{2} \quad R \sin \alpha = 1$$

$$R = \sqrt{3} \quad \tan \alpha = \frac{1}{\sqrt{2}}$$

$$\alpha = 0.615$$

$$\text{so } \sqrt{3} \sin(\theta + 0.615) = 1$$

$$\theta + 0.615 = \sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\text{as } 0 \leq \theta \leq 2\pi \quad \theta = 0, 1.911$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

$$\theta = 0.615 - 0.615 \text{ or } \theta = (\pi - 0.615) - 0$$

013 CHS MATHS EXT 1 TRIAL HSC

Section 1

- 1) B 2) C 3) A 4) D 5) D
- 6) B 7) A 8) A 9) A 10) C

Section 2

Question 11

- a) -3
- b) 2x

$$\sqrt{1-x^4}$$

- c) $\frac{4}{3}$



- d) $u = 4 - x^3$ $u = 1$ $u = 3$

$$du = -3x^2 dx$$

$$\int_{-1}^1 x^2 \sqrt{4-x^3} dx = \int_3^1 \sqrt{u} \cdot \frac{-du}{3}$$

$$= -\frac{1}{3} \int_3^1 u^{\frac{1}{2}} du$$

$$= -\frac{1}{3} \left[\frac{2}{3} u^{\frac{3}{2}} \right]_3^1$$

$$= -\frac{2}{9} \left[\sqrt{3} - \sqrt{27} \right]$$

$$= \frac{2}{9} \left[\sqrt{27} - \sqrt{3} \right]$$

$$= \frac{2}{9} \left[3\sqrt{3} - \sqrt{3} \right]$$

$$= \frac{2}{9} \left[2\sqrt{3} \right]$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$= \frac{4\sqrt{3}}{9}$$

$$f) M = M_0 e^{-kt}$$

$$\frac{1}{2} M_0 = M_0 e^{-kt}$$

$$k = \frac{\ln(2)}{5580}$$

$$k \approx 1.2422 \times 10^{-4}$$

$$0.4 M_0 = M_0 e^{-kt}$$

$$t = \frac{\ln(0.4)}{k}$$

$$t \approx 7376 \text{ years}$$

$$t \approx 7376 \text{ years}$$

Question 13

$$a) \cos 2x = 2 \cos^2 x - 1$$

$$\text{so } 2 \cos^2 x - 1 = \sqrt{3} \cos x - 1$$

$$2 \cos^2 x - \sqrt{3} \cos x = 0$$

$$\cos x (2 \cos x - \sqrt{3}) = 0$$

$$\cos x = 0 \text{ or } \cos x = \frac{\sqrt{3}}{2}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$x = \frac{\pi}{2} \text{ or } x = \frac{\pi}{6}$$

$$i) y = p^2 - ap^2$$

$$x = \frac{y + ap^2}{p}$$

$$i) \frac{y + ap^2}{p} = \frac{y + ap^2}{p}$$

$$y + ap^2 = py + ap^2$$

$$py - ap^2 = py - pa$$

$$py - pa = py - pa$$

$$y(p-p) = -a(p-p)$$

$$y = -a$$

$$y = -a$$

$$d) \sqrt{2} \sin \theta + \cos \theta = R \sin(\theta + \alpha)$$


$$R \cos \alpha = \sqrt{2} \quad R \sin \alpha = 1$$

$$R = \sqrt{3} \quad \tan \alpha = \frac{1}{\sqrt{2}}$$

$$\alpha = 0.615$$

$$\text{so } \$$

9)



i) $V = \pi \int_0^{\pi} [\cos x]^2 dx$

x	0	$\frac{\pi}{2}$	π
y	1	0	1

$h = \frac{1}{2}$
 $\frac{1}{3} = \frac{1}{6}$

$V = \frac{\pi}{6} \left[\left(\pi^2 + \frac{\pi^2}{4} + 4 \times \frac{4\pi^2}{9} \right) + \left(\frac{\pi^2}{4} + 0 + 4 \times \frac{\pi^2}{9} \right) \right]$
 $= \frac{67\pi^3}{108} \quad (19.24)$

ii) $n=1 \quad 7^2 - 3 = 42 \therefore \text{true for } n=1$

Assume true for $n=k$

$7^{2k} - 3 = 11M$

Prove true for $n=k+1$

$7^{2k+2} - 3 = 7^2 [7^{2k} - 3] + 7^2 \cdot 3 - 3$
 $= 7^2 \cdot 11M + 3(7^2 - 3)$
 $= 11(49M + 2 \times 3^2)$

$\therefore$ If true for $n=k$, then also true for $n=k+1$.

As it is true for $n=1$, then by mathematical induction it is true for

all $n=1, 2, 3, \dots$

e) $x = 50 \cos \theta$

$t = \frac{x}{50 \cos \theta}$

$y = \frac{-9}{2} \left(\frac{x}{50 \cos \theta} \right)^2 + 50 \left(\frac{x}{50 \cos \theta} \right) \sin \theta$
 $= \frac{-9x^2}{5000} \sec^2 \theta + x \tan \theta$

ii) When $x=60 \quad 2.1 \leq y \leq 2.3$

$2.3 = 60 \tan \theta - \frac{10 \times 60^2}{5000} (1 + \tan^2 \theta)$

$\frac{36}{5} \tan^2 \theta - 60 \tan \theta + (23 + \frac{36}{5}) = 0$

$A = \frac{36}{5} \quad B = -60 \quad C = (23 + \frac{36}{5})$

$\tan \theta = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}$

$0 < \theta < 45$

$\tan \theta = 0.5381$

$\theta = 28.3^\circ \quad (28^\circ 17')$

using $C = (21 + \frac{36}{5})$

$\tan \theta = \frac{1}{2}$

$\theta = 26.6^\circ \quad (26^\circ 34')$

$\therefore$ Angle must be between 26.6° and 28.3°