

ST IGNATIUS COLLEGE RIVERVIEW



TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

2000

MATHEMATICS

3/4 UNIT COMMON

Time allowed: Two hours
(plus 5 minutes reading time)

Instructions to Candidates

- Attempt all questions
- All questions are of equal value.
- Show all necessary working. Marks may be deducted for missing or poorly arranged work.
- Standard integrals are provided
- Board approved calculators may be used.
- Each question attempted must be returned in a *separate* writing booklet clearly marked Question 1, Question 2 etc, on the cover
- Each booklet must have your student number and the name of your Class Teacher.

Students are advised that this is a trial examination only and cannot in any way guarantee the content or format of the 2000 3/4 unit Mathematics Higher School Certificate Examination


Question 1 (12 marks) Start a new booklet

- (a) Solve $|x-3| > 5$ 2 marks
- (b) Find the exact value of $\cos^{-1}\left(\frac{-1}{\sqrt{2}}\right)$ 1 mark
- (c) Differentiate with respect to x : $e^{-\ln x}$ 2 marks
- (d) Show that $\int_{-2}^{2\sqrt{3}} \frac{dx}{\sqrt{16-x^2}} = \frac{\pi}{2}$ 2 marks
- (e) Find the coefficient of x^5 in the expansion of $\left(x + \frac{1}{x}\right)^{13}$ 2 marks
- (f) (i) Sketch $y = \frac{1}{x}$. 1 mark
- (ii) Hence or otherwise find the values of x for which $\frac{1}{x} > x$ 2 marks

Question 2 (12 marks) Start a new booklet

- (a) Find $\int_e^{e^2} \frac{\ln x}{x} dx$ using the substitution $u = \ln x$ 3 marks
- (b) (i) Prove that $\frac{1-\cos x}{\sin x} = \tan \frac{x}{2}$ 2 marks
- (ii) Hence sketch $y = \frac{1-\cos x}{\sin x}$ for $-\pi < x < \pi$ 2 mark

- (c) The graphs of $y=x^3-x$ and $y=x^2+1$ intersect at $P(x,y)$ as shown in the diagram.



- (i) Show that $1 < x < 2$. 2marks
- (ii) Taking $x = 1.8$ as a first approximation to the x -value of P , use one application of Newton's method to find a closer value for x . 3marks

Question 3 (12 marks) Start a new booklet

(a)




Diagram
not to scale

RS is a diameter. PQ is perpendicular to PR .
Prove that Q , T and S are collinear.

3marks

(b)

- (i) State the domain and range of

$$y = 2 \sin^{-1} 3x.$$

2marks

- (ii) Sketch $y = 2 \sin^{-1} 3x$.

1mark

- (iii) The graph of $y = 2 \sin^{-1} 3x$ is rotated about the y -axis. Show that the volume generated is $\frac{\pi^2}{9} \text{ units}^3$.

4marks

(c)

Julian has 10 different pairs of socks where the left sock and right sock of each pair are indistinguishable.

Find the number of odd pairs of socks (ie a pair which do not match) that Julian can wear. Explain your reasoning.

2marks

Question 4 (12 marks) Start a new booklet

- (a) Find all solutions to $\cos x = \frac{\sqrt{3}}{2}$.

2marks

- (b) (i) Show that $\frac{1}{(x+1)(x+2)} = \frac{1}{x+1} - \frac{1}{x+2}$

1mark

- (ii) Hence evaluate $\int_0^1 \frac{dx}{(x+1)(x+2)}$

2marks

(c)

Tangents from the point $T(x_0, y_0)$ touch the parabola $x^2 = 4y$ at $P(x_1, y_1)$ and $Q(x_2, y_2)$.

- (i) State the equation of the chord of contact.

1 mark

- (ii) Show that the x -values of P and Q are given by the roots of the equation $x^2 - 2x_0x + 4y_0 = 0$

2marks

- (iii) Hence or otherwise prove that the midpoint M of QP is
 $\left(x_0, \frac{1}{2}x_0^2 - y_0\right)$. 2marks

- (iv) If T moves on the line $y=x-1$ find the equation of the locus of M .
2marks

Question 5 (12 marks) Start a new booklet

- (a) Prove by Mathematical Induction that the expression $5^n - 1$ is divisible by 4 for all positive integers n .
4marks

- (b) Metal Fatigue is a phenomenon where a piece of steel will fail when repeatedly subjected to a force F . The endurance limit is the force below which the steel will not break even if subjected to an infinite number of applications of that force. Let the number of applications be n .

The force and the number of applications are related by the differential equation

$$\frac{dF}{dn} = -k(F - F_0) \quad \text{where } k \text{ and } F_0 \text{ are constants.}$$

- (i) Show that $F = 275e^{-k(n-1)} + F_0$ is a solution to
 $\frac{dF}{dn} = -k(F - F_0)$ 1 mark
- (ii) If $F=350$ when $n=1$, find the value of F_0 .
1mark
- (iii) Find the endurance limit. 1 mark
- (iv) Find the value of k if $F=80$ when $n=200$.
2marks

(c)

In today's society, statistics show that 28% of Australian women will never have children. Three women are selected at random. Find the probability that

- (i) they will all have children 1marks
- (ii) at least one of them will have children 2marks

Question 6 (12 marks) Start a new booklet

- (a) Consider the function $f(x) = e^{-x^2}$

- (i) Show that the function is even. 1mark
- (ii) Find the stationary point of $y=f(x)$. 1mark
- (iii) Show that $\frac{d^2y}{dx^2} = -2e^{-x^2}(1 - 2x^2)$ and hence find any points of inflexion.
2marks
- (iv) Sketch the curve of $y=f(x)$, $x \geq 0$. 1 mark
- (v) Sketch the inverse function $f^{-1}(x)$ of $f(x) = e^{-x^2}$ $x \geq 0$. 1mark
- (vi) Find the equation of $f^{-1}(x)$ and state its domain. 3marks

(b)

Malaysia has invented a minirail system which is completely automated, running to precision timing (ignoring passenger boarding and alightment). The journey between two stations A and B , where A is to the west of B , can be modelled by the equation $v^2 = 2(8x - x^2 - 7)$ where v is velocity in km/h and x is displacement in km from the central automated control office.

- (i) Show that the motion is simple harmonic. 1mark
- (ii) Find the distance between the two stations. 1mark
- (iii) Where is the control office in relation to A and B ? 1mark

Question 7 (12 marks) Start a new booklet

(a)




Diagram not to scale

The cross-section of a light fabric structure for a stadium roof is described by the equation $y = \frac{100}{x^2 + 100}$.

Dimensions are in metres.


- If Q is the point $(0, \frac{1}{4})$ find the value of k . 1 mark
- Show that the shaded area is $\frac{5(4\pi - 3\sqrt{3})}{3}$ square metres. 2 marks
- By considering the integral $\int_{-k}^k \frac{100}{x^2 + 100} dx$ or otherwise show that the area of the cross-section will never exceed 10π square metres. 2 marks

(b)

By considering the coefficient of x^{n+1} on both sides of the identity $(x+1)^n (x+1)^n = (x+1)^{2n}$ prove that

$${}^nC_0 {}^nC_1 + {}^nC_1 {}^nC_2 + {}^nC_2 {}^nC_3 + \dots + {}^nC_{n-1} {}^nC_n = \frac{(2n)!}{(n-1)!(n+1)!}$$
3 marks

(c)



One of the great historic problems which prompted the development of calculus was whether a cannonball would reach a target. Using the origin as shown and assuming $\ddot{x} = 0$ and $\ddot{y} = -10$, if a cannonball is fired at an angle of 45 degrees at a velocity of 30 m/s,

- show that $x = 15t\sqrt{2}$ and $y = -5t^2 + 15t\sqrt{2} + 6$. 2 marks
- Hence determine whether or not the ball will reach its target. 2 marks

Question 1

a) $|x - 3| > 5$
 $x - 3 > 5$ or $x - 3 < -5$
 $x > 8$ or $x < -2$

b) $\cos^{-1}\left(-\frac{1}{\sqrt{2}}\right) = \pi - \cos^{-1}\left(\frac{1}{\sqrt{2}}\right)$
 $= \pi - \frac{\pi}{4}$
 $= \frac{3\pi}{4}$

c) $\frac{d}{dx} e^{-\ln x} = -\frac{1}{x} e^{-\ln x}$
 $= -\frac{1}{x} e^{\ln \frac{1}{x}} \quad \left(e^{\ln \frac{1}{x}} = \frac{1}{x}\right)$
 $= -\frac{1}{x} \cdot \frac{1}{x}$
 $= -\frac{1}{x^2}$

Most students didn't get past this step to simplify answer. $\therefore$ second mark was not awarded to them.


d) $\int_{-2}^{2\sqrt{3}} \frac{dx}{\sqrt{16 - x^2}} = \left[\sin^{-1} \frac{x}{4} \right]_{-2}^{2\sqrt{3}}$
 $= \left(\sin^{-1} \frac{\sqrt{3}}{2} \right) - \left(\sin^{-1} -\frac{1}{2} \right)$
 $= \frac{\pi}{3} + \frac{\pi}{6}$
 $= \frac{\pi}{2}$

e) $\left(x + \frac{1}{x}\right)^{13}$
 $T_{r+1} = {}^{13}C_r (x^r) (x^{-1})^{13-r}$
 $= {}^{13}C_r (x^r) (x^{r-13})$
 $= {}^{13}C_r x^{2r-13}$

Since we are finding the coefficient of x^5 , let
 $2r - 13 = 5$

$2r = 18$
 $r = 9$
 $\therefore$ coefficient is ${}^{13}C_9 = 715$

f) i)



ii) Easier to use graph to solve $\frac{1}{x} > x$ than to solve algebraically.

$\therefore \frac{1}{x} > x$ for $0 < x < 1$ and $x < -1$

Question 2

$$\int_e^{e^2} \frac{\log_e x}{x} dx \text{ using the substitution } u = \log_e x$$

$$\frac{u}{x} \cdot x du$$

$$= \left[\frac{u^2}{2} \right]_1^2$$

get to limits in terms of u.

$$= 2 - \frac{1}{2}$$

$$= \frac{3}{2}$$

$$\frac{du}{dx} = \frac{1}{x}$$

$$dx = x du$$

$$\text{when } x = e^2, u = 2$$

$$\text{when } x = e, u = 1$$

ii) Sketch $y = \frac{1 - \cos x}{\sin x}$ for $-\pi < x < \pi$ ($x \neq 0$)


Using the result in (i), sketch $y = \tan \frac{x}{2}$ ($x \neq 0$)

$$y = \tan \frac{x}{2}$$

$$\text{Period} = \frac{\pi}{\frac{1}{2}}$$

$$= \frac{\pi}{1/2}$$

$$= 2\pi$$



i) Prove that $\frac{1 - \cos x}{\sin x} = \tan \frac{x}{2}$

$$\left\{ \begin{array}{l} \frac{x}{2} = t \\ x = \frac{1 - t^2}{1 + t^2} \\ x = \frac{2t}{1 + t^2} \end{array} \right.$$

$$\text{LHS} = 1 - \frac{1 - t^2}{1 + t^2}$$

$$\frac{2t}{1 + t^2}$$

$$= \frac{1 + t^2 - 1 + t^2}{1 + t^2} \times \frac{1 + t^2}{2t}$$

$$= \frac{2t^2}{2t}$$

$$= t$$

$$= \tan \frac{x}{2}$$

c) i) Solve $y = x^2 + 1$ and $y = x^3 - x$ simultaneously to find the point of intersection.

$$x^2 + 1 = x^3 - x$$

$$x^3 - x^2 - x - 1 = 0, \text{ Let } f(x) = x^3 - x^2 - x - 1$$

$$f(1) = -2$$

$$f(2) = 1$$

Since $f(1)$ and $f(2)$ are opposite in sign, then the root of $x^3 - x^2 - x - 1 = 0$ lies between $x=1$ and $x=2$.

ii) $f(x) = x^3 - x^2 - x - 1$

$$f'(x) = 3x^2 - 2x = 1.8 - \frac{-0.208}{5.12}$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$


$$\text{let } x_1 = 1.8$$

$$x_2 = 1.8 - \frac{f(1.8)}{f'(1.8)}$$

$$= 1.840625$$

Question 3

(a)



Construction

Join O to T,

T to S,

T to R.

$\angle RTS = 90^\circ$ (angle standing on a diameter)

$\angle QTR = 90^\circ$ (opposite angles of a cyclic quadrilateral are supplementary)

$\angle QTS = \angle RTS + \angle QTR$ (adjacent angles)
 $= 90^\circ + 90^\circ$
 $= 180^\circ$

O, T and S are collinear


(b)

i) $y = 2 \sin^{-1} 3x$

Domain: $-1 \leq 3x \leq 1$
 $-\frac{1}{3} \leq x \leq \frac{1}{3}$

Range: $-\frac{\pi}{2} \leq \sin^{-1} 3x \leq \frac{\pi}{2}$
 $-\pi \leq 2 \sin^{-1} 3x \leq \pi$
 $-\pi \leq y \leq \pi$

ii)



(Many students had curve in the wrong quadrants or concavity of curve was incorrect)

iii) $V = \pi \int_a^b x^2 dy$

$= \pi \int_{-\pi}^{\pi} \frac{1}{9} \sin^2 \frac{y}{2} dy$

$= 2\pi \cdot \frac{1}{9} \int_0^{\pi} \sin^2 \frac{y}{2} dy$

$= \frac{2\pi}{9} \int_0^{\pi} \frac{1}{2} (1 - \cos y) dy$

$= \frac{\pi}{9} [y - \sin y]_0^{\pi}$

$= \frac{\pi}{9} [(\pi - \sin \pi) - (0 - 0)]$

$= \frac{\pi^2}{9}$

$y = 2 \sin^{-1} 3x$

$\frac{y}{2} = \sin^{-1} 3x$

$3x = \sin \frac{y}{2}$

$x = \frac{1}{3} \sin \frac{y}{2}$

$x^2 = \frac{1}{9} \sin^2 \frac{y}{2}$

$\sin^2 \frac{y}{2} = \frac{1}{2} (1 - \cos y)$

This is from $\sin^2 A = \frac{1}{2} (1 - \cos 2A)$

iv)

A A

B B

C C

D D

E E

F F

G G

H H

I I

J J

10 choices for 1st sock ($^{10}C_1$)

9 choices for 2nd sock (9C_1)

number of odd pairs = 10×9

= 90

(Note: left sock and right sock are indistinguishable)

Alternate method:

Total combinations = $^{10}C_2$

= 190

subtract 10 pairs of matching socks

$190 - 10 = 180$

Question 4 (12 marks)

Comments

$$\cos x = \frac{\sqrt{3}}{2}$$

$$x = \frac{\pi}{6} \text{ (acute)} \quad (2)$$

$$x = 2n\pi \pm \frac{\pi}{6}$$

$$\frac{1}{(x+1)(x+2)} = \frac{1}{x+1} - \frac{1}{x+2}$$

$$\text{RHS} = \frac{1}{x+1} - \frac{1}{x+2}$$

$$= \frac{x+2 - (x+1)}{(x+1)(x+2)} \quad (1)$$

$$= \frac{1}{(x+1)(x+2)}$$

$$= \text{LHS}$$

$$\int_0^1 \frac{dx}{(x+1)(x+2)}$$

$$= \int_0^1 \left(\frac{1}{x+1} - \frac{1}{x+2} \right) dx \quad (2)$$

$$= \left[\ln(x+1) - \ln(x+2) \right]_0^1$$

$$= \left[\ln \left(\frac{x+1}{x+2} \right) \right]_0^1$$

$$\ln \frac{2}{3} - \ln \frac{1}{2}$$

$$\ln \frac{4}{3}$$

Most students did not know the general solution formula for $\cos^{-1}x$.

part (i) was generally done well

most students were able to find the indefinite integral as a log function BUT

many made errors in evaluating the definite integral.

(c) (i) Chord of Contact: $xx_0 = 2(y+y_0)$ (1)

(ii) $xx_0 = 2(y+y_0)$ — (1)

$$x^2 = 4y$$

$$y = \frac{x^2}{4} \quad \text{--- (2)}$$

Solve simultaneously for x

$$xx_0 = 2\left(\frac{x^2}{4} + y_0\right)$$

$$xx_0 = \frac{x^2}{2} + 2y_0 \quad (2)$$

$$2xx_0 = x^2 + 4y_0$$

$$x^2 - 2x_0x + 4y_0 = 0 \quad (A)$$

(iii) Use the quadratic formula to solve (A)

$$x = \frac{2x_0 \pm \sqrt{4x_0^2 - 4(4y_0)}}{2}$$

$$= \frac{2x_0 \pm 2\sqrt{x_0^2 - 4y_0}}{2}$$

$$= x_0 \pm \sqrt{x_0^2 - 4y_0}$$

Midpoint of PA = average of roots

$$x = \frac{x_0 + \sqrt{x_0^2 - 4y_0} + x_0 - \sqrt{x_0^2 - 4y_0}}{2}$$

$$x = x_0$$

Sub into (1) to find y .

(i) very few students could remember this formula.

(ii) Not knowing the formula made part virtually impossible

(iii) Some students use the sum of roots method

$$\text{sum of roots} = -\frac{b}{a}$$

$$= 2x_0 \text{ from}$$

$$\text{Average of roots} = \frac{\text{sum of roots}}{2}$$

$$= x_0$$

etc to find y_m .

QUESTION 5

COMMENTS

- 1) When $n=1$, $5^n - 1 = 4$ which is divisible by 4. $\therefore$ statement true when $n=1$
 Assume ^{result} true when $n=k$
 ie Assume $5^k - 1 = 4M$, $M \in \mathbb{I}$
 Now prove that the result holds when $n=k+1$
 ie Prove that $5^{k+1} - 1 = 4N$, $N \in \mathbb{I}$ ✓
 Now $5^{k+1} - 1$
 $= 5 \cdot 5^k - 1$ ✓
 $= (4+1)5^k - 1$
 $= 4 \cdot 5^k + 5^k - 1$ ✓ but $5^k - 1 = 4M$
 $= 4 \cdot 5^k + 4M$
 $= 4(5^k + M)$ ✓ but $5^k + M$ is an integer
 $= 4N$, $N \in \mathbb{I}$
 $\therefore$ result holds when $n=k+1$
 $\therefore$ since the result is true for $n=1$, it is also true for $n=2$ and hence for $n=3$ and so on for all $n \in \mathbb{I}^+$.

you do not assume $n=k$ you do not prove that $n=k+1$

$$5 \cdot 5^k - 1 \neq 5(5^k - 1)$$

$$\begin{aligned} &= 5 \cdot 5^k - 1 \\ &= 5(4M+1) - 1 \\ &= 20M + 4 = 4(5M+1) \\ &= 4N \end{aligned}$$

[4]

2) $\frac{dF}{dn} = -k(F - F_0)$ if $F = 275e^{-k(n-1)} + F_0$

(i) LHS = $\frac{dF}{dn} = \frac{d(275e^{-k(n-1)} + F_0)}{dn}$

$$= -k \cdot 275 e^{-k(n-1)}$$

$$\begin{aligned} \text{RHS} &= -k(F - F_0) = -k(275e^{-k(n-1)} + F_0 - F_0) \\ &= -k \cdot 275 e^{-k(n-1)} \end{aligned}$$

$$\text{LHS} = \text{RHS}$$

$$\therefore F = 275e^{-k(n-1)} + F_0 \text{ is a solution to } \frac{dF}{dn} = -k(F - F_0)$$

similar to showing that a point is on a line.
 It is best set up as an identity

Poorly done. You got away with very poor solutions because it was worth only 1 mark.

[1]

(ii) $F = 275e^{-k(n-1)} + F_0$ if $F=350$ when $n=1$
 $350 = 275e^{-k(1-1)} + F_0$
 $F_0 = 75$ ✓ [1]

$$350 = 275 + F_0$$

not division not $F_0 =$

(iii) $F = 275e^{-k(n-1)} + 75$
 $\lim_{n \rightarrow \infty} (275e^{-k(n-1)} + 75) = 0 + 75 = 75$

$\therefore$ the endurance limit is 75 ✓ [1]

(iv) if $F=80$ when $n=200$
 $80 = 275e^{-k(200-1)} + 75$
 $5 = 275e^{-199k}$
 $e^{-199k} = \frac{5}{275}$
 $-199k = \ln\left(\frac{5}{275}\right)$

$$k = -\ln\left(\frac{5}{275}\right) \div 199 = 0.02013735 \text{ (7 s.f.)} \quad \checkmark [2]$$

this question is mostly well done

(c) $P(\text{no children}) = .28$
 $P(\text{children}) = .72$

be careful of the "not"s

(i) $P(\text{ccc}) = .72^3 = 0.373248 \checkmark [1] \quad 37\%$

(ii) $P(\text{at least one child})$
 $= 1 - P(\text{none has children})$
 $= 1 - .28^3 \checkmark$
 $= .978048 \checkmark$

if you add these up separately don't forget 3C_1 .

[2] 98%

(iv) $T(x_0, y_0)$ moves on
the line $y = x - 1$
 $\therefore y_0 = x_0 - 1$ — (1)

$$M(x_0, \frac{1}{2}x_0^2 - y_0) \quad (2)$$

$$x = x_0 \quad \text{--- (1)}$$

$$y = \frac{1}{2}x_0^2 - y_0 \quad \text{--- (2)}$$

$$\begin{aligned} \text{A): } y &= \frac{1}{2}x_0^2 - (x_0 - 1) \\ &= \frac{1}{2}x_0^2 - x_0 + 1 \\ y &= \frac{1}{2}x^2 - x + 1 \end{aligned}$$

Question 6 (12 marks)

$$f(x) = e^{-x^2}$$

$$\begin{aligned} f(a) &= e^{-a^2} \\ f(-a) &= e^{-(-a)^2} \\ &= e^{-a^2} \\ &= f(a) \end{aligned}$$

$$\begin{aligned} \therefore f(x) \text{ is even since} \\ f(a) &= f(-a) \end{aligned}$$

It is important
under examination
conditions to realise
that parts (iii) and (iv)
of this question could
have been attempted
independently of
parts (i) and (ii)

$\rightarrow$ learn to SCROUNGE
FOR MARKS

(i) generally well done
HOWEVER

it is not good enough
to show that

$$f(1) = f(-1) \text{ etc}$$

ie for only one value
of x

$\rightarrow$ you must show
 $f(x) = f(-x)$ for
any value of x
e.g. $x = a$

$$\begin{aligned} \text{(a) (ii) } f(x) &= e^{-x^2} \\ f'(x) &= -2x e^{-x^2} \end{aligned}$$

$$\begin{aligned} \text{t.p. when } f'(x) &= 0 \quad (1) \\ -2x e^{-x^2} &= 0 \\ x &= 0 \\ y &= 1 \end{aligned}$$

x	-1	0	1
$f'(x)$	+	0	-

Max at (0,1)


$$\begin{aligned} \text{(iii) } f''(x) &= -2x - 2x e^{-x^2} + -2 \cdot e^{-x^2} \\ &= -2e^{-x^2} (1 - x^2) \end{aligned}$$

$$\begin{aligned} \text{infl. pt may occur when } f''(x) &= 0 \\ \text{ie } -2e^{-x^2} (1 - x^2) &= 0 \quad (2) \\ x^2 &= 1 \\ x &= \pm \frac{1}{\sqrt{2}} \end{aligned}$$

$$\text{infl. pts } \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{e}}\right) \text{ and } \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{e}}\right)$$

x	-2	$-\frac{1}{\sqrt{2}}$	0	$\frac{1}{\sqrt{2}}$	2
$f''(x)$	+	0	-	0	+

$\therefore$ change of concavity through I.P.'s



(ii) Many students could
not differentiate e^{-x^2}
 $\rightarrow -2x e^{-x^2}$ (common)


$\rightarrow$ should always classify
the t.p.'s
 $\rightarrow$ marks were not
deducted since this was
only worth 1 mark

(iii) When asked to find
either t.p.'s or I.
you should always
find the y-co-ord
 $\rightarrow$ many students just
found $x = 0$ part (i)
 $x = \pm \frac{1}{\sqrt{2}}$ part (ii)

$\rightarrow$ it is good practice to
test the concavity.
 $f''(x) = 0$ DOES NOT
PROVE change in
concavity !!

(iv) students were not
penalised for sketching
the full bell shape
nor for omitting the
inflection point

1) Question 6 (continued)



(1)

$$y = e^{-x^2}$$

$$x = e^{-y^2}$$

$$\ln x = -y^2$$

$$y^2 = -\ln x$$

$$= \ln \frac{1}{x}$$

(3)

use $\ln$ $y = \sqrt{\ln(\frac{1}{x})}$ take $+\sqrt{\quad}$ since range is $y \geq 0$

$Df^{-1}: 0 < x \leq 1$

$$(i) \quad v^2 = 2(8x - x^2 - 7)$$

$$\frac{1}{2}v^2 = 8x - x^2 - 7$$

$$\frac{d}{dx}(\frac{1}{2}v^2) = 8 - 2x$$

$$\ddot{x} = -2(x - 4)$$

HM Since it is in the form $\ddot{x} = -n^2x$ where $n = \sqrt{2}$, and centre of oscillation is $x = 4$ km from control office.

(iv) This was poorly done
 $\rightarrow$ most students had difficulty in finding the inverse shape
 $\rightarrow$ reflection in $y = x$
 $\rightarrow$ interchange $x \leftrightarrow y$

(vi) students picked up marks here, even though they were unable to sketch the inverse they were able to perform the inverse algebraic operations.


(i) Many students forgot about the formula

$$\frac{d}{dx}(\frac{1}{2}v^2) = \ddot{x}$$

$\therefore$ They struggled to prove the motion was S.H.M.

$\rightarrow$ i.e. that acceleration is proportional to displacement.

(b) (ii) $v^2 = 2(8x - x^2 - 7)$
 max displacement when $v = 0$
 $\therefore 2(8x - x^2 - 7) = 0$
 $x^2 - 8x + 7 = 0$
 $(x-7)(x-1) = 0$ (1)
 $x = 1$
 $x = 7$



(iii) Control office is 1 km West of Station (A) (1)

(ii) generally well done by those who attempted this part

(iii) Students thought the control office would be at the centre of the motion $\rightarrow$ question stated the distance was measured FROM THE CONTROL OFFICE $(x=0)$.

$$y = \frac{100}{x^2+100}$$

1) when $y = \frac{1}{4}$, $\frac{1}{4} = \frac{100}{x^2+100}$
 $x^2+100 = 400$

$$x^2 = 300 \quad x = \pm 10\sqrt{3}$$

From diagram, $k > 0 \therefore k = 10\sqrt{3}$ ✓

$$A = 2 \int_0^{10\sqrt{3}} \frac{100}{x^2+100} dx - \frac{1}{4} \times 20\sqrt{3}$$
 ✓

$$= 2 \times 100 \times \frac{1}{10} \left[\tan^{-1} \frac{x}{10} \right]_0^{10\sqrt{3}} - 5\sqrt{3}$$
 ✓

$$= 20 \left[\tan^{-1} \sqrt{3} - \tan^{-1} 0 \right] - 5\sqrt{3}$$

$$= 20 \cdot \frac{\pi}{3} - 5\sqrt{3} = 5 \left(\frac{4\pi}{3} - \sqrt{3} \right)$$

$$= \frac{5(4\pi - 3\sqrt{3})}{3}$$

$$\text{Area} = 2 \int_0^k \frac{100}{100+x^2} dx - 2k \times \frac{100}{k^2+100}$$

$$= 20 \left[\tan^{-1} \left(\frac{x}{10} \right) \right]_0^k - \frac{200k}{k^2+100}$$

$$= 20 \tan^{-1} \left(\frac{k}{10} \right) - \frac{200k}{k^2+100}$$
 ✓

now as $k \rightarrow \infty$

$$\tan^{-1} \left(\frac{k}{10} \right) \rightarrow \frac{\pi}{2} \quad \text{and} \quad \frac{200k}{k^2+100} \rightarrow 0$$
 ✓

$$\therefore \text{limit of Area is } 20 \times \frac{\pi}{2} = 10\pi$$

$$\therefore \text{area never exceeds } 10\pi \text{ m}^2$$


[1]

$\frac{1}{4} \times 20\sqrt{3}$ is area of rectangle below shaded area from $(-k, 0)$ to $(k, 0)$

$$\text{or } 2 \int_0^{10\sqrt{3}} \left(\frac{100}{x^2+100} - \frac{1}{4} \right) dx$$

[2]

$$\text{or } 2 \int_0^k \left(\frac{100}{100+x^2} - \frac{1}{4} \right) dx$$



$$\lim_{k \rightarrow \infty} \frac{200k}{k^2+100}$$

$$= \lim_{k \rightarrow \infty} \frac{200}{1 + \frac{100}{k^2}} = \frac{0}{1+0} = 0$$

[2]

$$\text{LHS} = (x+1)^n (x+1)^n$$

$$= \left[\binom{n}{0} x^n + \binom{n}{1} x^{n-1} + \binom{n}{2} x^{n-2} + \dots + \binom{n}{n-2} x^2 + \binom{n}{n-1} x + \binom{n}{n} \right]$$

$$\times \left[\binom{n}{0} x^n + \binom{n}{1} x^{n-1} + \binom{n}{2} x^{n-2} + \dots + \binom{n}{n-2} x^2 + \binom{n}{n-1} x + \binom{n}{n} \right]$$

learn how to set it up.

In this product the coefficient of x^{n+1} will be

$$\binom{n}{0} \binom{n}{n-1} + \binom{n}{1} \binom{n}{n-2} + \binom{n}{2} \binom{n}{n-3} + \dots + \binom{n}{n-2} \binom{n}{1} + \binom{n}{n-1} \binom{n}{0}$$
 ✓

But $\binom{n}{n-k} = \binom{n}{k} \therefore \binom{n}{n-1} = \binom{n}{1}, \binom{n}{n-2} = \binom{n}{2}, \dots, \binom{n}{0} = \binom{n}{n}$ ✓

$\therefore$ coefficient can be written as

$$\binom{n}{0} \binom{n}{1} + \binom{n}{1} \binom{n}{2} + \binom{n}{2} \binom{n}{3} + \dots + \binom{n}{n-2} \binom{n}{n-1} + \binom{n}{n-1} \binom{n}{n}$$

$$\text{LHS} = (x+1)^{2n}$$

$$= \binom{2n}{0} x^{2n} + \binom{2n}{1} x^{2n-1} + \dots + \binom{2n}{n-1} x^{n+1} + \dots + \binom{2n}{2n}$$

Here the coefficient of x^{n+1} is $\binom{2n}{n-1}$

but $\binom{2n}{n-1} = \frac{(2n)!}{(n-1)!(2n-(n-1))!}$


$$= \frac{(2n)!}{(n-1)!(n+1)!}$$
 ✓

$$\therefore \binom{n}{0} \binom{n}{1} + \binom{n}{1} \binom{n}{2} + \dots + \binom{n}{n-1} \binom{n}{n} = \frac{(2n)!}{(n-1)!(n+1)!}$$

[3]

QED

c)



Initially

$$\dot{y} = V \sin \theta = 30 \sin 45^\circ = 30 \times \frac{1}{\sqrt{2}} = 15\sqrt{2}$$

$$\dot{x} = V \cos \theta = 30 \cos 45^\circ = 15\sqrt{2}$$

$$i) \ddot{x} = 0$$

$$\dot{x} = c_1 \text{ but } \dot{x} = 15\sqrt{2} \text{ initially}$$

$$\therefore \dot{x} = 15\sqrt{2}$$

$$x = 15\sqrt{2}t + c_2 \text{ but } x = 0 \text{ when } t = 0 \therefore c_2 = 0$$

$$\therefore x = 15\sqrt{2}t$$

$$\ddot{y} = -10$$

$$\dot{y} = -10t + c_3 \text{ but } \dot{y} = 15\sqrt{2} \text{ when } t = 0 \therefore c_3 = 15\sqrt{2}$$

$$\therefore \dot{y} = -10t + 15\sqrt{2}$$

$$y = -5t^2 + 15\sqrt{2}t + c_4 \text{ but } y = 6 \text{ when } t = 0 \therefore c_4 = 6$$

$$\therefore y = -5t^2 + 15\sqrt{2}t + 6$$

Target is at (100, 4) $\therefore$ for the ball to reach target
 $y \geq 4$ when $x = 100$

$$\text{when } x = 100$$

$$100 = 15\sqrt{2}t$$

$$t = \frac{100}{15\sqrt{2}} = \frac{100\sqrt{2}}{30} = \frac{10\sqrt{2}}{3} \checkmark$$

$$\text{when } t = \frac{10\sqrt{2}}{3}$$

$$y = -5\left(\frac{10\sqrt{2}}{3}\right)^2 + 15\sqrt{2}\left(\frac{10\sqrt{2}}{3}\right) + 6$$

$$= \frac{-5 \times 200}{9} + 100 + 6$$

$$= -5\frac{1}{9}$$

$\therefore$ the ball will not reach the target. $\checkmark$

COMMENTS

set initial $\dot{x}$ & $\dot{y}$ up
before you start

Integrating wrt t


[1] must set up $\dot{x}$ & $\dot{y}$
find constants
all the way
through.

[1]

"hit" is not the same as
"reach".

(would hit only if $y = 4$
when $x = 100$)

OR set $y = 0$, find t ,
find x ($= 95.65$)



[2]