



NORMANHURST BOYS HIGH SCHOOL

MATHEMATICS ADVANCED

2024 Year 11 Course Assessment Task 2

Friday, 28 June 2024

General instructions

- Working time – 1 hour.
(plus 5 minutes reading time)
- Write using blue or black pen. Where diagrams are to be sketched, these may be done in pencil.
- NESA approved calculators may be used.
- Attempt **all** questions.

SECTION I

- Mark your answers on the answer grid provided (on page 2)

SECTION II

- All necessary working should be shown in every question. Marks may be deducted for illegible or incomplete working.

NESA STUDENT #:

Class (please ✓)

☐ 11MAA.1 – Ms Quek

Marker's use only.

QUESTION	1-3	4-5	6-7	8-9	10-11	12-15	16	Total	%
MARKS	3	9	13	7	8	14	6	60	

Section I

3 marks

Attempt Question 1 to 3

Allow approximately 4 minutes for this section

Mark your answers on the answer grid.

Questions

Marks

1. Which of the following is equivalent to 1

$$\frac{\cos(90^\circ - \theta)}{\sin(90^\circ - \theta)}$$


- (A) 1 (B) $\cot \theta$ (C) $\tan \theta$ (D) $-\tan \theta$

2. Evaluate $\lim_{x \rightarrow 1} \frac{x^2 + 2x - 3}{x - 1}$ 1

- (A) undefined (B) 0 (C) 1 (D) 4

3. Which of the following points on the graph of $y = f(x)$ has the following properties simultaneously: 1

- $f'(x) = 0$
- $f''(x) = 0$



- (A) A (B) B (C) C (D) D

Answer grid for Section I

- 1** – ☐ A ☐ B ☐ C ☐ D
- 2** – ☐ A ☐ B ☐ C ☐ D
- 3** – ☐ A ☐ B ☐ C ☐ D

Examination continues overleaf...

Section II

57 marks

Attempt Question 4 to 16

Allow approximately 56 minutes for this section

Write your answers in the space provided.

Question 4 (6 marks)

Differentiate with respect to x

(a) $x^2\sqrt{x}$ **2**

.....

.....

.....

(b) $(2x^2 + 1)^6$ **2**

.....

.....

.....

.....

.....

.....

(c) $\frac{1 + 7x}{x^3}$ **2**

.....

.....

.....

.....

.....

.....

.....

Question 5 (3 marks)

If $f(x) = \frac{1}{x-1}$, show that $f'(x) = -\frac{1}{(x-1)^2}$ by differentiation from first principles. **3**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Examination continues overleaf...

Question 6 (5 marks)

- (a) **By using the product rule**, show that the derivative of $y = (3x^3 + x)(x - 1)$ is **2**

$$\frac{dy}{dx} = 12x^3 - 9x^2 + 2x - 1$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence, find the equation of the normal to the curve $y = (3x^3 + x)(x - 1)$ at the point where $x = 1$. Express your answer in general form $ax + by + c = 0$. **3**

.....

.....

.....

.....

.....

.....


.....

.....

.....

Question 7 (8 marks)

The graph of $y = x^3 + x^2 - x + 2$ is sketched below.



The points A and B are the turning points.

- (a) Find the coordinates of A and B .

4

.....

.....

.....

.....

.....

.....

.....

- (b) For what values of x is the curve concave up?

2

Show all mathematical reasons for your answer.

.....

.....

.....

- (c) For what value(s) of k does the equation $x^3 + x^2 - x + 2 = k$ only has one real solution?

2

.....

Examination continues overleaf...

Question 8 (4 marks)

Given $\log_7 2 = 0.36$ and $\log_7 5 = 0.83$, find the values of the following.

(a) $\log_7 0.4$ **2**

.....

.....

.....

(b) $\log_7 50$ **2**

.....

.....

.....

.....

.....

Question 9 (3 marks)

In the space below, sketch the graph of $y = \log_5(5x - 5)$, showing all key features including: **3**

- any x intercept(s)
- any asymptote(s)

Examination continues overleaf...

Question 10 (4 marks)

- (a) Solve the following equation for x , correct to 2 decimal places. **2**
Full working out must be shown to gain full marks.

$$0.4^{x+6} = 128$$

.....

.....

.....

.....

.....

.....

- (b) By considering the graph of $y = 0.4^{x+6}$ and also part (a), find the values of x such that **2**

$$0.4^{x+6} < 128$$

(*Hint: draw a rough sketch of the graph of $y = 0.4^{x+6}$*)

.....

.....

.....

.....

Examination continues overleaf...

Question 11 (4 marks)

A drug is used to control a medical condition. It is known that the quantity Q of drug remaining in the body after t hours satisfies an equation of the form

$$Q = Q_0 e^{-kt} \quad \text{where } Q_0 \text{ and } k \text{ are constants}$$

The initial dose is a 6 milligrams and after 15 hours the amount remaining in the body is half the initial dose.

- (a) Find the value of Q_0 . **1**

.....

.....

.....

- (b) Show that the value of k is 0.046, correct to 3 decimal places. **1**

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Show (with full working out) that it takes approximately 45 hours 12 minutes for one-eighth of the initial dose to remain. **2**

.....

.....

.....

.....

.....

.....

.....

.....

Examination continues overleaf...

Question 12 (4 marks)

Sydney is 694 nautical miles (M) south and 761 M west of Noumea. **4**

A plane must fly to make a trip from Noumea to Sydney. Calculate the distance and the true bearing of the flight to the nearest nautical miles and degree respectively.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

Examination continues overleaf...

Question 13 (3 marks)

The diagram shows $\triangle ABC$ with sides $AB = 7\text{cm}$, $BC = 5\text{cm}$ and $AC = 10\text{cm}$.



- (a) Show that $\cos A = \frac{31}{35}$ **1**

.....

.....

.....

.....

.....

.....

.....

- (b) Hence by finding the exact value of $\sin A$, determine the exact area of $\triangle ABC$. **2**

.....

.....

.....

.....

.....

.....

.....

Examination continues overleaf...

Question 14 (2 marks)

Given $\sin \theta = \frac{3}{4}$ and $\tan \theta < 0$, find the exact value of $\cos \theta$

2

.....

.....

.....

Question 15 (5 marks)

(a) Prove that $\frac{2 \cos^2 \theta}{1 - \sin \theta} = 2 + 2 \sin \theta$

2

.....

.....

.....

.....

.....

.....


(b) Hence draw a sketch of a function $y = \frac{2 \cos^2 \theta}{1 - \sin \theta}$ in the domain $-90^\circ < \theta < 90^\circ$

3

Examination continues overleaf...

Question 16 (6 marks)

A person is standing at point A , which is due south of a tower OT that is h metres tall. The person walks 50 metres east to point B . The angle of elevation at points A and B to the top of the tower are 45° and 30° respectively.



- (a) Express OA and OB in terms of h . **2**

.....

.....

.....

.....

- (b) Show that $h = 25\sqrt{2}$ metres. **2**

.....

.....

.....

.....

- (c) Find the bearing of B from the base of the tower O , correct to the nearest minute. **2**

.....

.....

.....

.....

.....

.....

End of paper.

Additional writing space



NORMANHURST BOYS HIGH SCHOOL

MATHEMATICS ADVANCED

2024 Year 11 Course Assessment Task 2

Friday, 28 June 2024

General instructions

- Working time – 1 hour.
(plus 5 minutes reading time)
- Write using blue or black pen. Where diagrams are to be sketched, these may be done in pencil.
- NESA approved calculators may be used.
- Attempt **all** questions.

SECTION I

- Mark your answers on the answer grid provided (on page 2)

SECTION II

- All necessary working should be shown in every question. Marks may be deducted for illegible or incomplete working.

NESA STUDENT #: ...Sample...solution...

Class (please ✓)

☐ 11MAA.1 – Ms Quek

Marker's use only.

QUESTION	1-3	4-5	6-7	8-9	10-11	12-15	16	Total	%
MARKS	3	9	13	7	8	14	6	60	

Section I

3 marks

Attempt Question 1 to 3

Allow approximately 4 minutes for this section

Mark your answers on the answer grid.

Questions

Marks

1. Which of the following is equivalent to

1

$$\frac{\cos(90^\circ - \theta)}{\sin(90^\circ - \theta)} = \frac{\sin \theta}{\cos \theta} = \tan \theta$$

- (A) 1 (B) $\cot \theta$ (C) ☒ $\tan \theta$ (D) $-\tan \theta$

2. Evaluate $\lim_{x \rightarrow 1} \frac{x^2 + 2x - 3}{x - 1} = \lim_{x \rightarrow 1} \frac{(x+3)(x-1)}{x-1} = \lim_{x \rightarrow 1} x+3 = 4$


1

- (A) undefined (B) 0 (C) 1 (D) ☒ 4

3. Which of the following points on the graph of $y = f(x)$ has the following properties simultaneously:

1

- $f'(x) = 0 \rightarrow$ stationary point
- $f''(x) = 0 \rightarrow$ Horizontal point of inflection



- (A) A (B) ☒ B (C) C (D) D

Answer grid for Section I

- | | | | | |
|-----|-------------------------|------------------------------------|------------------------------------|------------------------------------|
| 1 – | ☐ A | ☐ B | ☒ C | ☐ D |
| 2 – | ☐ A | ☐ B | ☐ C | ☒ D |
| 3 – | ☐ A | ☒ B | ☐ C | ☐ D |

Examination continues overleaf...

Section II

57 marks

Attempt Question 4 to 16

Allow approximately 56 minutes for this section

Write your answers in the space provided.

Question 4 (6 marks)

-1 Setting out overall.

Differentiate with respect to x

(a) $x^2\sqrt{x} = x^{2+\frac{1}{2}} = x^{\frac{5}{2}}$ ① $\frac{d}{dx}(x^{\frac{5}{2}}) = \frac{5}{2}x^{\frac{3}{2}}$ ① 2

(b) $(2x^2 + 1)^6$ $\frac{d}{dx}(2x^2 + 1)^6 = 6(2x^2 + 1)^5 \times 4x$ ① 2
 $= 24x(2x^2 + 1)^5$ ①

(c) $\frac{1+7x}{x^3} = \frac{1}{x^3} + \frac{7x}{x^3}$ 2
 $= x^{-3} + 7x^{-2}$ ①
 $\frac{d}{dx}(x^{-3} + 7x^{-2}) = -3x^{-4} + 7x^{-2} \times -2x^{-3}$
 $= -\frac{3}{x^4} - \frac{14}{x^3}$ ①

Question 5 (3 marks)

If $f(x) = \frac{1}{x-1}$, show that $f'(x) = -\frac{1}{(x-1)^2}$ by differentiation from first principles.

3

$$f(x+h) = \frac{1}{x+h-1}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

① for keeping $\lim_{h \rightarrow 0}$ till the end

$$= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h-1} - \frac{1}{x-1}}{h}$$

①

$$= \lim_{h \rightarrow 0} \frac{\frac{x-1 - (x+h-1)}{(x+h-1)(x-1)}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x-1-x-h+1)}{h(x+h-1)(x-1)}$$

$$= \lim_{h \rightarrow 0} \frac{-h}{h(x+h-1)(x-1)}$$

$$= -\frac{1}{(x-1)(x-1)}$$

$$= -\frac{1}{(x-1)^2}$$

①

Examination continues overleaf...

Question 6 (5 marks)

- (a) By using the product rule, show that the derivative of $y = (3x^3 + x)(x - 1)$ is **2**

$$\frac{dy}{dx} = 12x^3 - 9x^2 + 2x - 1$$

$$u = 3x^3 + x \quad v = x - 1$$

$$u' = 9x^2 + 1 \quad v' = 1$$

$$\frac{dy}{dx} = (9x^2 + 1)(x - 1) + (3x^3 + x) \quad \textcircled{1}$$

$$= 9x^3 - 9x^2 + x - 1 + 3x^3 + x$$

$$= 12x^3 - 9x^2 + 2x - 1 \quad \textcircled{1}$$

- (b) Hence, find the equation of the normal to the curve $y = (3x^3 + x)(x - 1)$ at the point where $x = 1$. Express your answer in general form $ax + by + c = 0$. **3**

$$\text{at } x=1, \quad y = (3x^3 + x)(x - 1) = 0$$

$$m_{\text{tangent}} = 12(1)^3 - 9(1)^2 + 2(1) - 1 = 4$$

$$\therefore m_{\text{normal}} = -\frac{1}{4}$$

$$\text{Equation of normal: } y - 0 = -\frac{1}{4}(x - 1)$$


$$-4y = x - 1 \quad \textcircled{1}$$

$$\therefore x + 4y - 1 = 0 \quad \textcircled{1}$$

- For leaving answer in general form.

Question 7 (8 marks)

The graph of $y = x^3 + x^2 - x + 2$ is sketched below.



The points A and B are the turning points.

- (a) Find the coordinates of A and B .

4

$$\frac{dy}{dx} = 3x^2 + 2x - 1$$

$$\text{When } \frac{dy}{dx} = 0;$$

$$3x^2 + 2x - 1 = 0 \quad (1)$$

$$(3x - 1)(x + 1) = 0$$

$$x = \frac{1}{3}, -1. \quad (1)$$

$$\text{When } x = \frac{1}{3}, y = \left(\frac{1}{3}\right)^3 + \left(\frac{1}{3}\right)^2 - \left(\frac{1}{3}\right) + 2 = \frac{49}{27}$$

$$\text{When } x = -1, y = 3 \quad (1) \text{ - For finding both } y \text{ values}$$

$$\therefore A(-1, 3) \text{ \& } B\left(\frac{1}{3}, \frac{49}{27}\right) \quad (1)$$

- (b) For what values of x is the curve concave up?

2

Show all mathematical reasons for your answer.

$$\text{Concave up when } \frac{d^2y}{dx^2} > 0$$

$$\frac{d^2y}{dx^2} = 6x + 2 \quad \therefore 6x + 2 > 0 \quad (1)$$

$$x > -\frac{1}{3} \quad (1)$$

- (c) For what value(s) of k does the equation $x^3 + x^2 - x + 2 = k$ only has one real solution?

2

$$k > 3 \quad (1) \text{ or } k < \frac{49}{27} \quad (1)$$

Examination continues overleaf...

Question 8 (4 marks)

Given $\log_7 2 = 0.36$ and $\log_7 5 = 0.83$, find the values of the following.

(a) $\log_7 0.4$

2

$$\begin{aligned} \log_7 0.4 &= \log_7 \frac{2}{5} \\ &= \log_7 2 - \log_7 5 \quad (1) \\ &= 0.36 - 0.83 \\ &= -0.47 \quad (1) \end{aligned}$$

(b) $\log_7 50$

2

$$\begin{aligned} \log_7 50 &= \log_7 (25 \times 2) \\ &= \log_7 5^2 + \log_7 2 \quad (1) \\ &= 2 \log_7 5 + \log_7 2 \\ &= 2 \times 0.83 + 0.36 \\ &= 2.02 \quad (1) \end{aligned}$$

Question 9 (3 marks)

In the space below, sketch the graph of $y = \log_5(5x - 5)$, showing all key features including:


3

- any x intercept(s)
- any asymptote(s)

asymptote;
when $5x - 5 = 0$
 $x = 1$

when $x = 2$;
 $y = \log_5(5 \times 2 - 5)$
 $= 1$

x -intercept;
when $5x - 5 = 1$
 $5x = 6$
 $x = \frac{6}{5}$



Examination continues overleaf...

Question 10 (4 marks)

- (a) Solve the following equation for
- x
- , correct to 2 decimal places.

2*Full working out must be shown to gain full marks.*

$$0.4^{x+6} = 128$$

$$\log 0.4^{x+6} = \log 128$$

$$(x+6) \log 0.4 = \log 128$$

$$x+6 = \frac{\log 128}{\log 0.4}$$


$$x = \frac{\log 128}{\log 0.4} - 6$$

$$= -11.30 \text{ (2dp)}$$

- (b) By considering the graph of
- $y = 0.4^{x+6}$
- and also part (a), find the values of
- x
- such that

2

$$0.4^{x+6} < 128$$

(Hint: draw a rough sketch of the graph of $y = 0.4^{x+6}$)

$$x > -11.30$$

Examination continues overleaf...

Question 11 (4 marks)

A drug is used to control a medical condition. It is known that the quantity Q of drug remaining in the body after t hours satisfies an equation of the form

$$Q = Q_0 e^{-kt} \quad \text{where } Q_0 \text{ and } k \text{ are constants}$$

The initial dose is a 6 milligrams and after 15 hours the amount remaining in the body is half the initial dose.

- (a) Find the value of Q_0 .

1

when $t=0$, $Q=6$

$$6 = Q_0 e^{-k \times 0}$$

$$\therefore Q_0 = 6$$

- (b) Show that the value of k is 0.046, correct to 3 decimal places.

1

when $t=15$, $Q = \frac{1}{2} Q_0$

$$\frac{1}{2} Q_0 = Q_0 e^{-15k}$$

$$\ln \frac{1}{2} = \ln e^{-15k}$$

$$\therefore -15k = \ln \frac{1}{2}$$

$$k = \frac{\ln \frac{1}{2}}{-15}$$

$$= 0.046 \text{ (3dp)}$$

- (c) Show (with full working out) that it takes approximately 45 hours 12 minutes for one-eighth of the initial dose to remain.

2

$t=?$ $Q = \frac{1}{8} Q_0$

$$\frac{1}{8} Q_0 = Q_0 e^{-0.046t}$$

$$\ln \frac{1}{8} = \ln e^{-0.046t}$$

$$-0.046t = \ln \frac{1}{8}$$

$$t = \frac{\ln \frac{1}{8}}{-0.046}$$

$$= 45^{\circ} 12' 18.9''$$

$$\therefore \text{takes approx. 45hrs 12mins.}$$


Examination continues overleaf...

Question 12 (4 marks)

Sydney is 694 nautical miles (M) south and 761 M west of Noumea.

4

A plane must fly to make a trip from Noumea to Sydney. Calculate the distance and the true bearing of the flight to the nearest nautical miles and degree respectively.



$$d = \sqrt{694^2 + 761^2} \quad (d > 0) \rightarrow \textcircled{1}$$

$$= 1029.93 \dots$$

$$= 1030 \text{ M}$$

①

$$\tan \alpha = \frac{694}{761}$$

$$\alpha = 42^\circ 21' 48.6''$$

$$= 42^\circ \text{ (nearest degree)}$$

$$\therefore \theta = 90^\circ - 42^\circ$$

$$= 48^\circ$$

①

$\therefore$ bearing from Noumea to Sydney is


$$048^\circ \text{T}$$

①

Examination continues overleaf...

Question 13 (3 marks)

The diagram shows $\triangle ABC$ with sides $AB = 7\text{cm}$, $BC = 5\text{cm}$ and $AC = 10\text{cm}$.




- (a) Show that $\cos A = \frac{31}{35}$

1

$$\begin{aligned}\cos A &= \frac{7^2 + 10^2 - 5^2}{2 \times 7 \times 10} \quad (1) \\ &= \frac{124}{140} \\ &= \frac{31}{35}\end{aligned}$$

- (b) Hence by finding the exact value of $\sin A$, determine the exact area of $\triangle ABC$.

2

$$\begin{aligned}h &= \sqrt{35^2 - 31^2} \\ &= \sqrt{264} \\ &= 2\sqrt{66}\end{aligned}$$

$$\therefore \sin A = \frac{2\sqrt{66}}{35}$$

(1)


$$\begin{aligned}A_{\triangle ABC} &= \frac{1}{2} \times 7 \times 10 \times \frac{2\sqrt{66}}{35} \\ &= 2\sqrt{66} \text{ units}^2 \quad (1)\end{aligned}$$

Examination continues overleaf...

Question 14 (2 marks)

Given $\sin \theta = \frac{3}{4}$ and $\tan \theta < 0$, find the exact value of $\cos \theta$

2

**Question 15** (5 marks)

(a) Prove that $\frac{2 \cos^2 \theta}{1 - \sin \theta} = 2 + 2 \sin \theta$


2

$$\begin{aligned}
 \text{LHS} &= \frac{2 \cos^2 \theta}{1 - \sin \theta} \\
 &= \frac{2(1 - \sin^2 \theta)}{1 - \sin \theta} \quad \text{①} \\
 &= \frac{2(1 + \sin \theta)(1 - \sin \theta)}{1 - \sin \theta} \\
 &= 2 + 2 \sin \theta \quad \text{①} \\
 &= \text{RHS}
 \end{aligned}$$

(b) Hence draw a sketch of a function $y = \frac{2 \cos^2 \theta}{1 - \sin \theta}$ in the domain $-90^\circ < \theta < 90^\circ$

3


$$\begin{aligned}
 y &= \frac{2 \cos^2 \theta}{1 - \sin \theta} \\
 &= 2 + 2 \sin \theta \\
 a &= 2 \quad n = 1 \quad T =
 \end{aligned}$$



Examination continues overleaf...

Question 16 (6 marks)

A person is standing at point A , which is due south of a tower OT that is h metres tall. The person walks 50 metres east to point B . The angle of elevation at points A and B to the top of the tower are 45° and 30° respectively.



- (a) Express OA and OB in terms of h . 2

$$\begin{aligned} \tan 45^\circ &= \frac{h}{OA} & \tan 30^\circ &= \frac{h}{OB} \\ OA &= \frac{h}{\tan 45^\circ} & OB &= \frac{h}{\tan 30^\circ} \\ &= h \quad (1) & &= \sqrt{3}h \quad (1) \end{aligned}$$

- (b) Show that $h = 25\sqrt{2}$ metres. 2

$$\begin{aligned} \text{In } \triangle AOB, \quad OB^2 &= OA^2 + AB^2 \\ (\sqrt{3}h)^2 &= h^2 + 50^2 \quad (1) \\ 2h^2 &= 50^2 \\ h &= \sqrt{\frac{50^2}{2}} \quad h > 0 \quad \therefore h = \sqrt{1250} \\ &= 25\sqrt{2} \text{ metres} \quad (1) \end{aligned}$$

- (c) Find the bearing of B from the base of the tower O , correct to the nearest minute. 2

$$\begin{aligned} \tan \angle AOB &= \frac{AB}{OA} \\ &= \frac{50}{25\sqrt{2}} \\ \angle AOB &= 54^\circ 44' \text{ (nearest mins)} \quad (1) \\ \therefore \text{bearing of } B \text{ from } O &= 125^\circ 16' \text{ (nearest min)} \quad (1) \end{aligned}$$

End of paper.