



Name: _____

Teacher: _____

Class: _____

FORT STREET HIGH SCHOOL

2011

HIGHER SCHOOL CERTIFICATE COURSE

ASSESSMENT TASK 3: TRIAL HSC

Mathematics

**TIME ALLOWED: 3 HOURS
(PLUS 5 MINUTES READING TIME)**

Outcomes Assessed	Questions	Marks
Chooses and applies appropriate mathematical techniques in order to solve problems effectively	1	
Manipulates algebraic expressions to solve problems from topic areas such as geometry, co-ordinate geometry, quadratics, trigonometry, probability and logarithms	3,4,7	
Demonstrates skills in the processes of differential and integral calculus and applies them appropriately	2,5,6,8	
Synthesises mathematical solutions to harder problems and communicates them in appropriate form	9,10	

Question	1	2	3	4	5	6	7	8	9	10	Total	%
Marks	/12	/12	/12	/12	/12	/12	/12	/12	/12	/12	/120	

Directions to candidates:

- Attempt all questions
- The marks allocated for each question are indicated
- All necessary working should be shown in every question. Marks may be deducted for careless or badly arranged work.
- Board – approved calculators may be used
- Each new question is to be started in a new booklet
- A table of standard integrals is supplied

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

Note $\ln x = \log_e x, \quad x > 0$

QUESTION 1 (12 marks) Start a NEW booklet.**Marks**(a) Evaluate $2e^2$ correct to three significant figures**2**(b) Factorise $2x^2 - x - 6$ **2**

(c) Simplify

$$\frac{2}{x+1} - \frac{3}{x}$$

2(d) Solve $|2x - 1| = 9$ **2**

(e) Expand and simplify

$$(\sqrt{5} - 1)(2\sqrt{5} + 3)$$

2

(f) Find the sum of the first 12 terms of the arithmetic sequence

2

$$3+6+9+\dots\dots$$

QUESTION 2 (12 marks) Start a NEW booklet.

(a) Differentiate

(i) $(x^3 + 1)^7$

2

(ii) $x^4 \log_e x$

2

(iii) $\frac{\sin x}{x+1}$

2(b) Find the perpendicular distance from (6, -2) to the line
 $4x - 3y + 7 = 0$ **2**

(c) (i) Find

$$\int \frac{1}{x+7} dx$$

1


(ii) Evaluate

$$\int_0^{\frac{\pi}{8}} \sec^2 2x \, dx$$

3

QUESTION 3 (12 marks) Start a NEW booklet.

Marks



The diagram shows the straight line l which cuts the x-axis and y-axis at $Q(6,0)$ and $R(0,2)$ respectively.


- | | |
|---|---|
| (i) Copy the diagram and find the equation of the line l . | 2 |
| (ii) Draw the line k , passing through Q perpendicular to l , and find the equation of the line k . | 2 |
| (iii) Find the coordinates of T , where k cuts the y-axis. | 1 |
| (iv) Find the equation of the circle which has TR as a diameter. Verify that Q lies on this circle. | 2 |
| (v) $(8,m)$ also lies on the circle. Find two possible values of m . | 2 |
- (b) Find the equation of the straight line that passes through $(1,-2)$ and the intersection point of the lines $4x + y - 5 = 0$ and $3x - 2y - 12 = 0$
- 3

QUESTION 4 (12 marks) Start a NEW booklet.

Marks

(a)

Figure not
to scale




In the diagram AB is parallel to CD.
Find the value of θ . Give reasons.

2

(b)

Figure not
to scale



Find the value of x . Give reasons. All measurements are in cm.


2

QUESTION 4 (continued)

Marks

(c)

Figure not
to scale



In the diagram AD is parallel to BC and
angle DBC = angle ACB = x° .

(i) Show that $AE = DE$.

2

(ii) Prove that the triangles ABC and DCB are congruent.

3

(iii) Deduce angle ABD = angle DCA.

3

QUESTION 5 (12 marks) Start a NEW booklet.

Marks

- (a) The gradient of a curve is given by

$$\frac{dy}{dx} = \frac{1}{\sqrt{2x+1}}$$

3

The curve passes through (4, 5). What is the equation of the curve? (Answer in surd form)

- (b) Consider the geometric series

$$1 + 3x + 9x^2 + 27x^3 + \dots$$

- (i) For what values of x does this series have a limiting sum?

2

- (ii) The limiting sum of this series is 100.
Find the value of x

2

- (c) A 5kg mass of metal is placed in an acid and it starts to Dissolve. After t hours, the mass M kg of undissolved metal is given by

$$M = 5e^{-kt}.$$

- (i) Find k , given $M = 4.2$ when $t = 2$
(Answer in exact form)

2

- (ii) After how many hours will half of the initial mass remain undissolved? Note: This known as the half-life of the metal in this acid. (Answer to 2 decimal places)

3

QUESTION 6 (12 marks) Start a NEW booklet.

Marks

- (a) A particle moves along a straight line so that its distance x , in metres, from a fixed point O is given by

$$x = 1 - 2\sin 2t,$$

where the time t is measured in seconds from $t = 0$.


- | | |
|--|---|
| (i) Where is the particle initially? | 1 |
| (ii) When and where, does the particle come to rest? | 3 |
| (iii) Where does the particle next come to rest? | 2 |
| (iv) What is the acceleration of the particle when
$t = \frac{\pi}{12}$? | 2 |
| (b) (i) Graph $y = \cos x$, for $0 \leq x \leq 2\pi$ | 1 |
| (ii) On your diagram shade the regions bounded by the curve $y = \cos x$, the x -axis and the lines $x = 0$ and $x = \pi$.
Calculate the total area of these regions. | 2 |
| (iii) Solve $\cos x = \frac{\sqrt{3}}{2}$ for $0 \leq x \leq 2\pi$
(Answer in terms of π) | 1 |

QUESTION 7 (12 marks) Start a NEW booklet**Marks****(a) Solve**

$$\log_e x - \frac{3}{\log_e x} = 2$$

3

(Answer in exact form)

(b)

A car travels at 45 km/h on a circular curve whose radius is 0.5km.

1(i) Find the distance l km, that the car travel in one minute.**2**(ii) Calculate the size of the angle θ through which the car turns in one minute. Give your answer to the nearest degree.**(c) Pat and Chris each threw a die****1**

(i) Find the probability that they throw the same number.

1

(ii) Find the probability that the number thrown by Chris is greater than the number thrown by Pat.

(d) The focus of a parabola is S (1, -2) and the directrix is the line $y = 5$.**1**

(i) Write down the co-ordinates of the vertex.

1

(ii) Find the focal length.

2

(iii) Write down the equation of the parabola in the form

$$(x - h)^2 = 4a(y - k)$$

and sketch the parabola

Marks**QUESTION 8 (12 marks) Start a NEW booklet**

(a) Consider the curve given by $y = 3x^2 - x^3$

4

(i) Find the stationary points and determine their nature.

(ii) Sketch the curve, indicating where it crosses the x-axis and showing stationary points.

2

2

(iii) Find the equation of the tangent to the curve at the point R (-1,4).

1

(b) (i) Differentiate $\log_e(\cos x)$ with respect to x.

(ii) Hence or otherwise, show

3

$$\int_0^{\frac{\pi}{4}} \tan x dx = \frac{1}{2} \log_e 2$$


QUESTION 9 (12 marks) Start a NEW booklet.

(a) Solve the equation

$$3^{2x} + 2 \times 3^x - 15 = 0$$

2

(b)



The shaded region in the diagram is bounded by the curve $y = x^4$, the y-axis, and the line $y = 16$

4

Calculate the volume of solid of revolution formed when this region is rotated about the y-axis. (Answer in terms of π)

(c) A woman contributes \$1500 each year into a superannuation fund for the first 25 years of her working life. For the next fifteen years (until retirement), she decides to increase this, and invest a total of \$5000 each year. Each contribution is paid at the beginning of the year.

6

If the investment earns 7% p.a. paid yearly over the whole year period, how much will her investment be upon retiring? (Answer to the nearest dollar)

QUESTION 10 (12 marks) Start a NEW booklet.**Marks**

- (a) The speed of a train was recorded at intervals of one minute. The times, in minute, and the corresponding speeds v , in kilometres per hour, are listed in the following table.

time (min)	0	1	2	3	4
v (km/h)	0	25	34	30	40

- (i) Explain why the distance x , in km, travelled by the train in these four minutes is given by

$$x = \int_0^{\frac{1}{15}} v dt$$

2

- (ii) Estimate x by using Simpson's Rule with five function values. (Answer to 1 decimal place)

3

- (b) A truck is to travel 1000 kilometres at a constant speed of v km/h.

When travelling at v km/h, the truck consumes fuel at the rate of

$$\left(6 + \frac{v^2}{50}\right) \text{ litres per hour.}$$

The truck company pays \$1.50/litre for fuel and pays each of 2 drivers \$36 per hour whilst the truck is travelling.

- (i) Let the total cost of fuel and the drivers' wages for the trip be C dollars. Show that

$$C = 30v + \frac{81000}{v}$$

3

- (ii) The truck must take no longer than 12 hours to complete the trip, and speed limits require that $v \leq 100$. At what speed v should the truck travel to minimize the cost C . (Answer to 2 decimal places)

4**END OF EXAMINATION**

(1)

2011 20 Trial HSC
Solutions

QUESTION ONE

(a) $2e^3 = 14.7781123...$ ✓
 $= 14.8$ (to 3 sig fig) ✓

(b) $2x^2 - x - 6$
 $= 2x^2 - 4x + 3x - 6$ ✓
 $= 2x(x-2) + 3(x-2)$
 $= (2x+3)(x-2)$ ✓

(c) $\frac{2}{x+1} - \frac{3}{x}$
 $= \frac{2x}{x(x+1)} - \frac{3(x+1)}{x(x+1)}$ ✓
 $= \frac{2x - 3x - 3}{x(x+1)}$
 $= \frac{-x-3}{x(x+1)}$ ✓

(d) $|2x-1| = 9$

either $2x-1 = 9$ or $2x-1 = -9$
 $2x = 10$ or $2x = -8$
 $x = 5$ ✓ or $x = -4$ ✓

(e) $(\sqrt{5}-1)(2\sqrt{5}+3)$ ✓
 $= \sqrt{5}(2\sqrt{5}+3) - 1(2\sqrt{5}+3)$
 $= 10 + 3\sqrt{5} - 2\sqrt{5} - 3$
 $= 7 + \sqrt{5}$ ✓

Some students did not show the initial answer from calculator.

Some students made mistakes with the minus sign.

Some students need to revise how to add and subtract surds.

(2)

(f) $S_n = \frac{n}{2} [2a + (n-1)d]$

$= \frac{12}{2} [2 \times 3 + (12-1) \times 3]$

$= 6 \times 39$

$= 234$

Some students did not remember the sum of AP or did not use it correctly.

(3)

QUESTION TWO

$$(a) (i) y = (x^3 + 1)^7$$

$$y' = 3x^2 \times 7 \times (x^3 + 1)^6$$

$$= 21x^2 (x^3 + 1)^6$$

$$(ii) y = x^4 \log_e x$$

$$y' = \frac{x^4}{x} + 4x^3 \log_e x$$

$$= x^3 + 4x^3 \log_e x$$

$$(iii) y = \frac{\sin x}{x+1}$$

$$y' = \frac{(x+1) \cos x - 1 \times \sin x}{(x+1)^2}$$

$$= \frac{(x+1) \cos x - \sin x}{(x+1)^2}$$

$$(b) \text{ perp dist} = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

$$= \frac{|4 \times 6 - 3 \times -2 + 7|}{\sqrt{4^2 + (-3)^2}}$$

$$= \frac{37}{5}$$

Students needed to use proper setting out involving y' , $\frac{dy}{dx}$ or $\frac{d}{dx}$

Many students did not know correct formula

(4)

$$(c) (i) \int \frac{1}{x+7} dx$$

$$= \log_e (x+7) + C$$

$$(ii) \int_0^{\frac{\pi}{8}} \sec^2 2x dx$$

$$= \left[\frac{1}{2} \tan 2x \right]_0^{\frac{\pi}{8}}$$


$$= \frac{1}{2} \tan \frac{\pi}{4}$$

$$= \frac{1}{2}$$

Remember constant in all indefinite integrals

(5)

QUESTION THREE



Some had the lines confused

gradient of $l = \frac{\text{rise}}{\text{run}} = -\frac{2}{6} = -\frac{1}{3}$ ✓

equation of l : $y = mx + b$

$y = -\frac{1}{3}x + 2$ ✓

should have been obvious that line l has a y int of 2! Yet some silly answers were given.

($b = y$ -intercept = 2)

See diagram for drawing of k

$m_l = -\frac{1}{3} \Rightarrow m_k = 3$ ✓
(neg reciprocal)

equation of k : $y - y_1 = m(x - x_1)$

$y - 0 = 3(x - 6)$

$\therefore y = 3x - 18$ ✓

(6)

(iii) k cuts the y -axis when $x = 0$

ie when $y = 3(0) - 18$

$= -18$

$\therefore T(0, -18)$ ✓

Need to learn to read the question. Many did not give the co-ordinates of T .

(iv) midpoint $TR = \left(\frac{0+0}{2}, \frac{2-18}{2}\right)$

$= (0, -8)$

$\therefore$ circle has centre $(0, -8)$
radius 10

Quite a few 'errors carried' here + hence many had wrong equation.

$\therefore$ eqn is $x^2 + (y + 8)^2 = 100$ ✓

sub $Q(6, 0)$ into

$x^2 + (y + 8)^2 = 100$

LHS = $6^2 + (0 + 8)^2$

$= 36 + 64$

$= 100$ ✓

$=$ RHS $\therefore Q(6, 0)$ lies on circle

For a 'show' question should start with LHS & work through to show it is equal to RHS. Those who had wrong circle eqn.

(v) $x^2 + (y + 8)^2 = 100$

sub $(8, m)$ $\Rightarrow$

$8^2 + (m + 8)^2 = 100$

$(m + 8)^2 = 36$

$m + 8 = \pm 6$ ✓

$m = \pm 6 - 8$

$\therefore m = -2$ or $m = -14$ ✓

obviously had a problem $\therefore$ should have realised the had made an error. (No mark awarded)

(vi) Some used ' y '

instead of ' m '
- need 2 values of m

(7)

(b) Using 'k' method

$$4x + y - 5 + k(3x - 2y - 12) = 0$$

sub $(1, -2)$

$$4 - 2 - 5 + k(3 + 4 - 12) = 0$$

$$-3 + -5k = 0$$

$$k = -\frac{3}{5}$$

$$\therefore 4x + y - 5 - \frac{3}{5}(3x - 2y - 12) = 0$$

$$20x + 5y - 25 - 9x + 6y + 36 = 0$$

$$11x + 11y + 11 = 0$$

$$x + y + 1 = 0$$

$$y = -x - 1$$

OR

$$4x + y - 5 = 0 \quad (1)$$

$$3x - 2y - 12 = 0 \quad (2)$$

$$8x + 2y - 10 = 0 \quad (3) = (1) \times 2$$

$$11x - 22 = 0 \quad (2) + (3)$$

$$x = 2 \quad y = -3$$

$\therefore$ Pt. of intersection

of (1) & (2) is $(2, -3)$

grad from $(2, -3)$ to $(1, -2)$

$$= \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 + 2}{2 - 1} = -1$$

$\therefore$ required eqn: $y - y_1 = m(x - x_1)$

$$y + 2 = -1(x - 1)$$

$$y + 2 = -x + 1$$

$$y = -x - 1$$

If this method was used some errors with 'k'

Most used this method with reasonable success

(8)

QUESTION FOUR

(a) $\angle EDC = \angle BAD$
(alternate angles $AD \parallel EC$)
 $= 60^\circ$

$$\therefore \angle ECD = \theta$$

$$= 180^\circ - 70^\circ - 60^\circ$$

$$\text{(angle sum of } \triangle ECD \text{ is } 180^\circ)$$

$$= 50^\circ$$

well done

(b)

$$\frac{x}{7} = \frac{5}{9} \quad (\text{ratio of intercepts})$$

$$x = \frac{7 \times 5}{9}$$

$$= \frac{35}{9}$$

$$= 3\frac{8}{9}$$

Some students did not provide a reason

(c) (i) $\angle ADE = \angle CBE$ $AD \parallel BC$
 $= x^\circ$ alternate $\angle$ s

Similarly $\angle DAE = x^\circ$

$AE = DE$ (sides opposite equal $\angle$ s $\angle ADE, \angle DAE$)

(ii) In $\triangle BEC$

$EB = EC$ (opposite equal $\angle$ s $\angle EBC, \angle ECB$)

But $AC = AE + EC$

and $DB = DE + EB$

$$\therefore AC = DB$$

In $\triangle ABC, \triangle DCB$

$AC = DB$ (as above)

BC is common

Some students need to provide clearer sequential geometrical proofs

(9)

$$\angle ACB = \angle DBC \text{ (given)} \checkmark$$

$$\therefore \triangle ABC \cong \triangle DCB \text{ (SAS)} \checkmark$$

$$(iii) \angle ABC = \angle DCB$$

(Corresponding $\angle$ s in congruent $\triangle$ s) $\checkmark$

$$\angle ABD + \angle DBC = \angle DCA + \angle ACB$$

But $\angle DBC = \angle ACB$ (given)

$$\therefore \angle ABD = \angle DCA \checkmark$$

To prove
 $\angle ABD = \angle DCA$
 some students also
 proved
 $\triangle AEB \cong \triangle DEC$
 This was also
 acceptable

(10)

QUESTION FIVE

$$(a) \frac{dy}{dx} = \frac{1}{\sqrt{2x+1}}$$

$$\frac{dy}{dx} = (2x+1)^{-\frac{1}{2}}$$

$$\therefore y = \frac{(2x+1)^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} + C \checkmark$$

$$y = \sqrt{2x+1} + C$$

$$\text{Sub } (4, 5) \Rightarrow 5 = \sqrt{2(4)+1} + C$$

$$5 = 3 + C \checkmark$$

$$C = 2$$

$$\therefore \text{equation is } y = \sqrt{2x+1} + 2 \checkmark$$

$$(b) (i) \text{ For limiting sum}$$

$$-1 < 3x < 1 \checkmark$$

$$-\frac{1}{3} < x < \frac{1}{3} \checkmark$$

$$(iii) S_{\infty} = \frac{a}{1-r}$$

$$100 = \frac{1}{1-3x} \checkmark$$

$$1-3x = \frac{1}{100}$$

$$-3x = -\frac{99}{100}$$

$$x = \frac{33}{100} \checkmark$$

Well done

Confusion in solving
 this simple equation

(11)

(e) $M = Se^{-kt}$

(e) when $M = 4.2$, $t = 3$

$$4.2 = Se^{-2k}$$

$$e^{-2k} = 0.84$$

$$\ln(0.84) = -2k$$

$$k = \frac{\ln(0.84)}{-2}$$

$$= 0.087176693$$

(Calc)

(ii) we want to find t

when $M = \frac{1}{2} \times 5 = 2.5$

$$2.5 = Se^{-0.087176693t}$$

$$0.5 = e^{-0.087176693t}$$

$$\ln(0.5) = -0.087176693t$$

$$t = \frac{\ln(0.5)}{-0.087176693}$$

$$= 7.95 \text{ hours}$$

(to 2 dec pls)

well done

(12)

QUESTION SIX

(a) (i) $x = 1 - 2 \sin 2\omega$

at $t = 0$, $x = 1 - 2 \sin 0 = 1$

i 1 metre on right-hand side of O

(ii) $v = \frac{dx}{dt} = -4 \cos 2t$

comes to rest means $v = 0$

$$-4 \cos 2t = 0$$

$$\cos 2t = 0$$

$$2t = \frac{\pi}{2} \text{ (1st time)}$$

$$t = \frac{\pi}{4}$$

$$t = \frac{\pi}{4} \quad x = 1 - 2 \sin 2 \times \frac{\pi}{4}$$

$$= 1 - 2 \sin \frac{\pi}{2}$$

$$= 1 - 2$$

$$= -1$$

i first comes to rest 1 metre on the left-hand side of O

(iii) $v = 0$ $\cos 2t = 0$

$$2t = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$t = \frac{\pi}{4}, \frac{3\pi}{4}$$

For $t = \frac{3\pi}{4}$

$$x = 1 - 2 \sin (2 \times \frac{3\pi}{4})$$

$$= 1 - 2 \sin \frac{3\pi}{2}$$

$$= 1 - 2 \times -1$$

$$= 1 + 2 = 3$$

i 3 metres on right-hand side of O

Some students answered when a particle come to rest, not where.


Some students answered when a particle come to a rest next, not where

(13)

$$(ii) \quad a = \frac{dv}{dt} = \frac{d}{dt} (-4 \cos 2t) \\ = 8 \sin 2t \quad \checkmark$$

$$t = \frac{\pi}{12} \quad a = 8 \sin \left(2 \times \frac{\pi}{12} \right) \\ = 8 \sin \frac{\pi}{6} \\ = 8 \times \frac{1}{2} \\ = 4$$

i.e. acceleration is $4 \text{ m s}^{-2} \quad \checkmark$



(a) See above, for shaded regions

$$A = \int_0^{\pi/2} \cos x dx + \left| \int_{\pi/2}^{\pi} \cos x dx \right| \\ = \left[\sin x \right]_0^{\pi/2} + \left| \left[\sin x \right]_{\pi/2}^{\pi} \right|$$

$$= \sin \frac{\pi}{2} - \sin 0 + \left| \sin \pi - \sin \frac{\pi}{2} \right|$$

$$= 1 - 0 + |0 - 1|$$

$$= 1 + 1$$

$$\therefore \text{Area} = 2 \text{ units}^2 \quad \checkmark$$

Some students did not take the absolute of the area below the x-axis

(14)

$$\text{OR} \quad A = 2 \int_0^{\pi} \cos x dx \quad \checkmark \\ = 2 (\sin x) \Big|_0^{\pi}$$

$$= 2 \left(\sin \frac{\pi}{2} - \sin 0 \right)$$

$$= 2 (1 - 0) \quad \checkmark$$

$$\text{Area} = 2 \text{ units}^2$$

$$(iii) \quad \cos x = \frac{\sqrt{3}}{2} \quad 0 \leq x \leq 2\pi \quad \text{well done} \\ x = \frac{\pi}{6}, \quad 2\pi - \frac{\pi}{6} \\ x = \frac{\pi}{6}, \quad \frac{11\pi}{6} \quad \checkmark$$

(15)

QUESTION SEVEN

$$(a) \log_e x - \frac{3}{\log_e x} = 3$$

$$\text{let } u = \log_e x$$

$$u - \frac{3}{u} = 3$$

$$u^2 - 3 = 3u$$

$$u^2 - 3u - 3 = 0 \quad \checkmark$$

$$(u-3)(u+1) = 0$$

$$u = 3 \quad \text{or} \quad u = -1 \quad \checkmark$$

$$\log_e x = 3 \quad \log_e x = -1$$

$$x = e^3 \quad x = e^{-1} \quad \checkmark$$

$$(b) (i) \begin{aligned} &45 \text{ km in } 1 \text{ hour} \\ &= 45 \text{ km in } 60 \text{ minutes} \\ &= \frac{3}{4} \text{ km in } 1 \text{ min} \quad \checkmark \end{aligned}$$

$$(ii) \begin{aligned} l &= r \theta \\ \frac{3}{4} &= 0.5 \times \theta \quad \checkmark \\ \theta &= \frac{3}{4} \times 2 \\ &= 1.5 \text{ radians} \\ &= 1.5 \times \frac{180^\circ}{\pi} \end{aligned}$$

$$\theta = 85.94366927$$

$$= 86^\circ \text{ (nearest deg)} \quad \checkmark$$


$$(c) (i) \text{ Probability} \\ = P[(1,1) \text{ or } (2,2) \text{ or } (3,3) \text{ or } (4,4) \text{ or } (5,5) \text{ or } (6,6)] \\ = \frac{1}{36} + \frac{1}{36} + \frac{1}{36} + \frac{1}{36} + \frac{1}{36} + \frac{1}{36} \\ = \frac{6}{36} = \frac{1}{6} \quad \checkmark \\ \text{or Probability (Chris's throw same as Pat's)} \\ = \frac{1}{6} \quad \checkmark$$

(16)

(ii) Let Pat be first, Chris be second relevant pairs are

(1,2) (1,3) (1,4) (1,5) (1,6)
 (2,3) (2,4) (2,5) (2,6)
 (3,4) (3,5) (3,6) $\checkmark$
 (4,5) (4,6)
 (5,6)

$$P(\text{Chris' no} > \text{Pat's no}) = \frac{15}{36} = \frac{5}{12} \quad \checkmark$$



$$(i) \text{ Vertex} = \text{midpt } SX = \left(\frac{1+1}{2}, \frac{5-3}{2} \right) = \left(1, \frac{3}{2} \right) \quad \checkmark$$

$$(ii) \text{ Focal length} = \text{dist } XV = 5 - \frac{3}{2} = \frac{7}{2} \quad \checkmark$$

$$(iii) \begin{aligned} (x-h)^2 &= 4a(y-k) \\ (x-1)^2 &= -4 \times \frac{7}{2} \left(y - \frac{3}{2} \right) \\ (x-1)^2 &= -14 \left(y - \frac{3}{2} \right) \\ x^2 - 2x + 1 &= -14y + 21 \\ x^2 - 2x + 14y - 20 &= 0 \quad \checkmark \end{aligned}$$

See above for sketch

Confusion with the focal length and vertex.

(17)

QUESTION EIGHT

(a) (i) $y = 3x^2 - x^3$ ✓

$$\frac{dy}{dx} = 6x - 3x^2 = 0$$

for stationary points

$$2x - x^2 = 0$$

$$x(2-x) = 0$$

$$x = 0 \text{ or } 2$$
 ✓

$$\frac{d^2y}{dx^2} = 6 - 6x$$

at $x = 0$ $\frac{d^2y}{dx^2} > 0$

∴ minimum at $(0, 0)$ ✓


at $x = 2$ $\frac{d^2y}{dx^2} < 0$

∴ maximum at $(2, 4)$ ✓(ii) x intercepts at $y = 0$

$$3x^2 - x^3 = 0$$

$$x^2(3-x) = 0$$

$$x = 0 \text{ or } 3$$

 $(2, 4)$ max $(0, 0)$

M-17

well done and
well set out.

(18)

(iii) $\frac{dy}{dx} = 6x - 3x^2$

at $x = -1$ $\frac{dy}{dx} = -6 - 3 = -9$ ✓

eqn of tangent

$$y - 4 = -9(x + 1)$$

$$y - 4 = -9x - 9$$

$$y = -9x - 5$$

$$9x + y + 5 = 0$$
 ✓

(b) (i) $y = \log_e(\cos x)$

$$= -\frac{\sin x}{\cos x}$$

$$= -\tan x$$

(ii) $\int_0^{\frac{\pi}{4}} \tan x = -[\log_e \cos x]_0^{\frac{\pi}{4}}$ ✓

$$= -\left[\log_e \frac{1}{\sqrt{2}} - \log_e 1\right]$$

$$= -\log_e \frac{1}{\sqrt{2}}$$
 ✓

$$= -\log_e 2^{\frac{1}{2}}$$
 ✓

$$= \frac{1}{2} \log_e 2$$

Students need
log laws rather
than a calculator.

(19)

QUESTION NINE

$$(a) 3^{2x} + 2 \times 3^x - 15 = 0$$

well done

$$\text{Let } u = 3^x, \quad u^2 = 3^{2x}$$

$$\text{Then } u^2 + 2u - 15 = 0$$

$$(u+5)(u-3) = 0$$

$$u = -5 \text{ or } u = 3$$

$$3^x = -5 \text{ or } 3^x = 3$$

no soln

$$x = 1$$

$$(b) V = \pi \int_0^{16} x^3 dy$$

well done

$$= \pi \int_0^{16} y^{\frac{1}{3}} dy$$

$$(y = x^{\frac{4}{3}} \text{ so } y^{\frac{1}{3}} = x^{\frac{1}{3}})$$

$$V = \pi \left[\frac{2y^{\frac{1}{3}}}{\frac{1}{3}} \right]_0^{16}$$

$$= \pi \left[\frac{2 \times 16^{\frac{1}{3}}}{\frac{1}{3}} - \frac{2 \times 0^{\frac{1}{3}}}{\frac{1}{3}} \right]$$

$$= \pi \times \frac{128}{3}$$

$$\text{Volume} = \frac{128}{3} \pi \text{ units}^3$$

(c) Treat the investment in 2 parts

Part 1 \$1500 invested each year for

40 years

$$\text{1st } \$1500 = \$1500 \times 1.07^{40}$$

$$\text{2nd } \$1500 = \$1500 \times 1.07^{39}$$

↓

$$\text{last } \$1500 = \$1500 \times 1.07^1$$

(20)

Part 2: For the last 15 years, the extra \$3500 (to make up the \$5000 to be invested each year for 15 years)

$$\text{1st } \$3500 = \$3500 \times 1.07^{15}$$

$$\text{2nd } \$3500 = \$3500 \times 1.07^{14}$$

↓

$$\text{last } \$3500 = \$3500 \times 1.07^1$$

Some students had difficulty establishing the G.P. for the two separate investments

$$\text{Total} = 1500 (1.07 + 1.07^2 + \dots + 1.07^{40}) \text{ plus}$$

$$3500 (1.07 + 1.07^2 + \dots + 1.07^{15})$$

$$= 1500 \times 1.07 \frac{(1.07^{40} - 1)}{1.07 - 1} + 3500 \times 1.07 \frac{(1.07^{15} - 1)}{1.07 - 1}$$

$$= \$320414.35 + \$94108.19$$

$$= \$414523 \text{ (nearest dollar)}$$

(3 marks for correctly calculating one part only)

QUESTION TEN

$$(a) \quad (i) \quad v = \frac{dx}{dt}$$

$$x = \int_a^b v dt \quad \checkmark$$

Since the initial time is 0, $a = 0$

The unit for velocity was km/h

The final time was 4 mins or $\frac{1}{15}$ h

$$\therefore b = \frac{1}{15} \quad \checkmark$$

$$\therefore x = \int_0^{\frac{1}{15}} \frac{1}{15} v dt$$

(ii) Using the formula:

$$\int_0^{\frac{1}{15}} f(t) dt$$

$$= \frac{h}{3} (y_0 + y_4 + 4 \times (y_1 + y_3) + 2y_2)$$

$$\text{where } h = \frac{1}{4} \left(\frac{1}{15} - 0 \right) \text{ i.e. } \frac{1}{60} \quad \checkmark$$

$$x = \frac{1}{3} \times \frac{1}{60} (0 + 0 + 4 \times (25 + 30) + 2 \times 34) \quad \checkmark$$

$$= \frac{1}{180} (228)$$

$$= 1.8 \text{ km (to 1 dp)} \quad \checkmark$$

(b) (i) Using speed = $\frac{\text{distance}}{\text{time}}$

$$\therefore \text{time} = \frac{\text{distance}}{\text{speed}} \quad \checkmark$$

$$\therefore \text{time} = \frac{1000}{v} \quad \checkmark$$

Very leniently marked
should have been
stating v is the
rate of change of
displacement w.r.t
time $\therefore$ to get
position we need
to integrate.

generally well
explained that
velocity was in km/h
 $\therefore$ time needs to
be given in hours
4 mins = $\frac{1}{15}$ h

Some had difficulty
calculating h .
Some had students
still getting formula
incorrect.

(22)

But cost per hour

$$= \text{driver cost/h} + \text{fuel cost/h}$$

$$= 2 \times 36 + \left(6 + \frac{v^2}{50} \right) \times 1.5$$

$$= 72 + 9 + \frac{1.5 v^2}{50}$$

$$= 81 + \frac{30 v^2}{100} \quad \checkmark$$

$$\therefore \text{Total Cost} = \left(81 + \frac{30 v^2}{100} \right) \times \frac{1000}{v} \quad \checkmark$$

$$= 30 v + \frac{81000}{v}$$

$$(ii) \quad \frac{dC}{dv} = -\frac{81000}{v^2} + 30 = 0 \quad \checkmark$$

$$30 v^2 = 81000$$

$$v^2 = \frac{81000}{30}$$

$$= 2700$$

$$v = \sqrt{2700}$$

$$= 51.96 \text{ km/h} \quad \checkmark$$

(to 2 decpl)

$$\frac{d^2C}{dv^2} = \frac{162000}{v^3} > 0$$

$$\therefore \text{min at } v = 51.96 \quad \checkmark$$

however the trip time must be

less than 12 hours

$$\frac{1000}{v} \leq 12 \quad \checkmark$$

$$v \geq \frac{1000}{12} \geq 83 \frac{1}{3}$$

Hence the soln found to $\frac{dC}{dv} = 0$ does not
satisfy the time constraint. Since cost
increases the further you are away from
51.96 km/h, the speed to minimise cost
is $v \geq 83.33 \text{ km/h (to 1 dp)}$

Some had difficulty
showing C

Many did not test
using the 2nd
derivative or attentive

Some ignored time
& speed constraints

Some ignored using
calculus at all
- penalised.