

Student Number:.....

Class Teacher:.....

St George Girls High School

Trial Higher School Certificate Examination

2016



Mathematics Extension 1

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black pen.
- Write your student number and your class teacher's name on each booklet.
- Board-approved calculators may be used.
- The mark allocated for each question is listed at the side of the question.
- Marks may be deducted for careless or poorly presented work.
- A table of standard integrals is provided.
- A multiple choice answer sheet is provided for Section I.

Total marks – 70

Section I – Pages 2 to 4 10 marks

- Attempt questions 1 to 10
- Allow about 15 minutes for this section.
- Answer on the sheet provided.

Section II – Pages 5 to 10 60 marks

- Attempt questions 11 – 15.
- Allow about 1 hour 45 minutes for this section.
- Begin each question in a new booklet.
- Show all necessary working in questions 11 - 15.

Students are advised that this is a trial Examination only and does not necessarily reflect the content or format of the Higher School Certificate Examination.

Section I:

Multiple Choice (Each question is worth 1 mark)

Answer this section on the multiple choice answer sheet provided.

1. The acute angle between the lines $x - 2y + 1 = 0$ and $2x - y - 1 = 0$ is closest to:

A. 37°

B. 45°

C. 90°

D. 143°

2. The point P divides the interval from A (-2, 2) to B (8, -3) internally in the ratio 3:2. What is the x coordinate of P?

A. 4

B. 2

C. 0

D. -1

3. Which of the following is the correct expression for $\int \frac{dx}{9+25x^2}$?

A. $\frac{1}{15} \tan^{-1} \frac{3x}{5} + C$

B. $\frac{1}{25} \tan^{-1} \frac{3x}{5} + C$

C. $\frac{1}{25} \tan^{-1} \frac{5x}{3} + C$

D. $\frac{1}{15} \tan^{-1} \frac{5x}{3} + C$

4. A curve has parametric equations $x = t - 3$ and $y = t^2 + 2$. What is the Cartesian equation of this curve?
- A. $y = x^2 - x - 1$
- B. $y = x^2 + x - 1$
- C. $y = x^2 - 6x + 11$
- D. $y = x^2 + 6x + 11$
5. A particle is moving in a straight line with $v^2 = 36 - 4x^2$ and undergoing simple harmonic motion. If the particle is initially at the origin, which of the following is the correct equation for its displacement in terms of t ? (v is the velocity and x is the displacement of the particle).
- A. $x = 2 \sin(3t)$
- B. $x = 3 \sin(2t)$
- C. $x = 2 \sin(9t)$
- D. $x = 3 \sin(4t)$
6. Solve the inequality $\frac{x^2 - 4}{x} \geq 0$
- A. $-2 \leq x < 0$ or $x \geq 2$
- B. $-2 \geq x > 0$ or $x \leq 2$
- C. $-4 \leq x < 0$ or $x \geq 4$
- D. $-4 \geq x > 0$ or $x \leq 4$
7. What is the value of $\int_0^1 \frac{4x}{2x+1} dx$? Use the substitution $u = 2x + 1$.
- A. $2 - \log_e 2$
- B. $2 - \log_e 3$
- C. $4 - 2\log_e 2$

- D. $4 - 2\log_e 3$
8. What is the correct expression for the indefinite integral $\int (\cos^2 x + 2 \sec^2 x) dx$?
- A. $\frac{1}{2}x + \frac{1}{4}\sin 2x + \tan x + C$
- B. $\frac{1}{2}x - \frac{1}{4}\sin 2x + \tan x + C$
- C. $\frac{1}{2}x + \frac{1}{4}\sin 2x + 2 \tan x + C$
- D. $\frac{1}{2}x - \frac{1}{4}\sin 2x + 2 \tan x + C$
9. What is the term independent of x in the expansion of $\left(x^2 - \frac{2}{x}\right)^9$?
- A. ${}^9C_3(-2)^3$
- B. ${}^9C_6(-2)^6$
- C. ${}^9C_3(2)^3$
- D. ${}^9C_6-(2)^6$
10. A particle moves in a straight line with a displacement of x and velocity of v . When $t = 0$ the acceleration is $3x^2$, velocity is $-\sqrt{2}$ and displacement is 1. Which of the following is the correct equation for x as a function of t ?
- A. $x = \frac{-2}{(t+\sqrt{2})^2}$
- B. $x = \frac{-2}{(t-\sqrt{2})^2}$
- C. $x = \frac{2}{(t+\sqrt{2})^2}$
- D. $x = \frac{2}{(t-\sqrt{2})^2}$

End of Section I

Section II:

Answer each question in a SEPARATE writing booklet.

In Questions 11, 12, 13, 14 and 15 your responses should include relevant mathematical reasoning and/or calculations.

Marks

Question 11 (12 marks) Start a new booklet

- a) Use Newton's method to find a second approximation to the positive root of $-2 \sin x = 0$. Take $x = 1.6$ as the first approximation.

2

- b) Solve:

$$|2x - 1| = |x|$$

2

- c) A particle moves in a straight line and its position (x) at any time (t) is given by:

$$x = 1 + \sqrt{3} \cos 4t + \sin 4t$$

- (i) Prove the motion is simple harmonic.

2

- (ii) When does the particle first reach maximum speed? ($t > 0$)

2

- d) $P(2p, p^2)$ and $Q(2q, q^2)$ are two points on the parabola $x^2 = 4y$. M is the midpoint of PQ .

- (i) Prove the identity $(p - q)^2 = 2(p^2 + q^2) - (p + q)^2$

1

- (ii) If P and Q move on the parabola so that $p - q = 4$, show that the locus of M is the parabola $x^2 = 4y - 16$

2

- (iii) What is the focus of the locus of M ?

1


Marks

Question 12 (12 marks) Start a new booklet

- a) The polynomial equation $8x^3 - 36x^2 + 22x + 21 = 0$ has roots which form an arithmetic progression. Find the roots.

3

- b) A is the top of a vertical radio mast AB standing on level ground. Two points C and D are on ground level such that C is due East of B and D is 500 metres due North of C .



The angle of elevation of A from C is $11^\circ 13'$ and the angle of elevation of A from D is $8^\circ 14'$. Calculate the height of the tower to the nearest metre.

3

- c) The velocity V of a particle decreases according to the equation:

$$\frac{dV}{dt} = -k(V - P)$$

where t is the time in seconds and k is a positive constant. The initial velocity of the particle is 0 ms^{-1} and the terminal velocity P is 60 ms^{-1} .

- (i) Show that $V = P + Ae^{-kt}$, where A is a constant, satisfies the given differential equation.

1


- (ii) Find the value of k if the velocity of the particle after 10 seconds is 35 ms^{-1} . Give your answer correct to two significant figures.

2

Marks

Question 12 continued

- d) $ABCD$ is a cyclic quadrilateral. EAF is a tangent at A to the circle. CA bisects $\angle BCD$.



NOT DRAWN TO SCALE

Show that $EAF \parallel DB$.

3

Marks

Question 13 (12 marks) Start a new booklet

a) (i) Show:

$$\sin x - \cos x = \sqrt{2} \sin \left(x - \frac{\pi}{4} \right)$$

2

(ii) Hence sketch the graph of $y = \sin x - \cos x$, for $0 \leq x \leq 2\pi$.

2

(iii) Show that $x = \frac{\pi}{2}$ is a solution to $\sin x - \cos x = 1$ and hence solve
 $\sin x - \cos x > 1$ for $0 \leq x \leq 2\pi$.

2

b) Prove:

$$\sin^{-1} \frac{1}{\sqrt{5}} + \sin^{-1} \frac{1}{\sqrt{10}} = \frac{\pi}{4}.$$

3


c) Use mathematical induction to show that the expression $7^n + 5$ is divisible by 6 for all positive integer values of n.

3

Marks

Question 14 (12 marks) Start a new booklet

- a) Water is pumped into a conical vessel at a constant rate of 24 cm^3 per second. The depth of the water is ' h ' cm at any time ' t ' seconds



What is the rate of increase in the area of the surface ' S ' of the water when the depth is 16 cm.

4

- b) In the expansion of $(1 + ax)^n$ in ascending powers of x , the first three terms are:

$$1 + 6x + 16x^2 + \dots$$

- (i) By comparing coefficients write two equations in a and n .

2

- (ii) Hence find the values of a and n .

2

- c) A particle is performing Simple Harmonic Motion in a straight line. At time t seconds it has displacement x metres from a fixed point O on the line, velocity $v \text{ ms}^{-1}$ and acceleration in ms^{-2} given by $\ddot{x} = -4(x - 1)$. When the particle is at the centre of its motion it has speed 6 ms^{-1} .

- (i) Show that $v^2 = -4x^2 + 8x + 32$.

2

- (ii) Find the period and amplitude of the motion.

2

Marks

Question 15 (12 marks) Start a new booklet

- a) State the largest possible (natural) domain of the function:

2

$$y = \log_e(\sin^{-1} x)$$

- b) A stone is projected from the top of a 80 metre high vertical cliff with an initial velocity of V ms at an angle of projection of θ . It reaches its greatest height after 3 seconds and hits the ground at a horizontal distance of 320 metres from the foot of the cliff. Assume $g = 10\text{ms}^{-2}$.

- (i) Using the top of the cliff as the origin, the horizontal position of the stone is given as $x = V \cos \theta t$. Show that the vertical component of the parametric equations of the path of the stone is:

$$y = -5t^2 + V \sin \theta t$$

2

- (ii) Show that $V \sin \theta = 30$.

1

- (iii) Find how long it takes for the stone to reach the ground.

1

- (iv) Show that $V \cos \theta = 40$

1

- (v) Find the value of V and the angle of projection.

1

- c) (i) Show that for all positive integers n ,

$$x[(1+x)^{n-1} + (1+x)^{n-2} + \dots + (1+x)^2 + (1+x) + 1] = (1+x)^n - 1$$

1

- (ii) Hence show that for $1 \leq k \leq n$

$$\binom{n-1}{k-1} + \binom{n-2}{k-1} + \binom{n-3}{k-1} + \dots + \binom{k-1}{k-1} = \binom{n}{k}$$


1

- (iii) Show that $n\binom{n-1}{k} = (k+1)\binom{n}{k+1}$

2

SECTION 1

Question	Solution	Criteria
1.	$x - 2y + 1 = 0$ $y = \frac{1}{2}x + \frac{1}{2}$ $m_1 = \frac{1}{2}$ $\tan \theta = \left \frac{m_1 - m_2}{1 + m_1 m_2} \right $ $= \left \frac{\frac{1}{2} - 2}{1 + \frac{1}{2} \times 2} \right $ $= \frac{3}{4}$ $2x - y - 1 = 0$ $y = 2x - 1$ $m_2 = 2$ $\theta = 36.86989765...$ $\approx 37^\circ$	A
2.	$\left(\frac{m \cdot x_2 + n \cdot x_1}{m+n}, \frac{m \cdot y_2 + n \cdot y_1}{m+n} \right)$ $m=3, n=2,$ $x_1=-2, x_2=8$ $y_1=2, y_2=-3$ $= \left(\frac{3 \times 8 + 2 \times -2}{3+2}, \frac{3 \times -3 + 2 \times 2}{3+2} \right)$ $= \left(\frac{24-4}{5}, \frac{-9+4}{5} \right)$ $= (4, -1)$	A
3.	$\int \frac{dx}{25 \left(\frac{9}{25} + x^2 \right)} \quad a = \frac{3}{5}$ $= \frac{5}{3} \times \frac{1}{25} \tan^{-1} \frac{5x}{3} + C$ $= \frac{1}{15} \tan^{-1} \frac{5x}{3} + C$	D
4.	$x = t - 3$ $t = x + 3$ $y = t^2 + 2$ $y = (x+3)^2 + 2$ $= x^2 + 6x + 11$	D
5.	$v^2 = 36 - 4x^2$ $= 2^2(9 - x^2) \quad [= n^2(a^2 - x^2)]$ $a^2 = 9, n = 2 \text{ and } x = 0 \text{ (initially at the origin)}$ $a = 3$ $x = a \sin(nt + \alpha)$ $= 3 \sin 2t$	B

Question	Solution	Criteria
6	$x^2 \times \frac{x^2-4}{x} \geq 0 \times x^2 \quad x \neq 0$ $x(x^2-4) \geq 0$ $x(x-2)(x+2) \geq 0$ $-2 \leq x \leq 0 \text{ or } x \geq 2$ 	A
7	$u = 2x+1 \quad x=1 \Rightarrow u=3$ $du = 2 dx \quad x=0 \Rightarrow u=1$ $\int_0^1 \frac{4x}{2x+1} dx = \int_1^3 \frac{2(u-1)}{u} \cdot \frac{1}{2} du$ $= \int_1^3 \left(1 - \frac{1}{u}\right) du$ $= [u - \log_e u]_1^3$ $= 3 - \log_e 3 - 1 + \log_e 1$ $= 2 - \log_e 3$	B
8	$\int (\cos^2 x + 2 \sec^2 x) dx = \int \left(\frac{1}{2}(1 + \cos 2x) + 2 \sec^2 x\right) dx$ $= \frac{1}{2}x + \frac{1}{4}\sin 2x + 2 \tan x + C$	C
9	$T_{r+1} = {}^9C_r (x^2)^{9-r} \left(-\frac{2}{x}\right)^r$ $= {}^9C_r x^{18-2r} (-2)^r x^{-r}$ $= {}^9C_r x^{18-3r} (-2)^r$ Term independent of x $18-3r=0$ $r=6$ $T_7 = {}^9C_6 (-2)^6$	B
10	$a = 3x^2$ $v^2 = 2 \int (3x^2) dx$ $= 2x^3 + C$ When $x=1$, $v=-\sqrt{2}$ Then $C=0$ $v = -\sqrt{2x^3} \quad (v < 0 \text{ when } x=1)$ $\frac{dv}{dt} = -\sqrt{2x^3}$ $\frac{dt}{dx} = -\frac{1}{\sqrt{2}} x^{-\frac{3}{2}}$ $t = \frac{2}{\sqrt{2}} x^{-\frac{1}{2}} + C_1$ When $t=0$, $x=1$ then $C_1 = -\sqrt{2}$ $x^{\frac{1}{2}} = \frac{t + \sqrt{2}}{\sqrt{2}}$ $x^{\frac{1}{2}} = \frac{\sqrt{2}}{t + \sqrt{2}}$ $x = \frac{2}{(t + \sqrt{2})^2}$	C

MATHEMATICS EXTENSION I - QUESTION //

19/1

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
a) $f(x) = x - 2\sin x$ $f'(x) = 1 - 2\cos x$ $f(1.6) = 1.6 - 2\sin 1.6$ $f'(1.6) = 1 - 2\cos 1.6$ $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$ $= 1.6 - \frac{1.6 - 2\sin 1.6}{1 - 2\cos 1.6}$ $= 1.977123551 \dots$ ≈ 1.98 or any other correct rounding off	1	many students forgot to use radians.
b) $2x-1 = x$ $x < 0$ $x \geq 0$ $2x-1 = -x$ $2x-1 = x$ $3x = 1$ $x = 1$ ✓ $x = \frac{1}{3}$ ✓ or: $(2x-1)^2 = (x)^2$ $(2x-1)^2 = x^2$ $4x^2 - 4x + 1 = x^2$ $3x^2 - 4x + 1 = 0$ $(3x-1)(x-1) = 0$ $x = 1$ or $x = \frac{1}{3}$ ✓ ✓	1	generally well done only mistakes being student only gave one soln.

MATHEMATICS EXTENSION I – QUESTION 11

pg 2

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
c) i) when $\ddot{x} = -v^2(x)$ SHM. $x = 1 + \sqrt{3} \cos 4t + \sin 4t$ $\dot{x} = -\sqrt{3} 4 \sin 4t + 4 \cos 4t$ $\ddot{x} = -\sqrt{3} \times 4^2 \cos 4t - 4^2 \sin 4t$ $= -4^2 (\sqrt{3} \cos 4t + \sin 4t)$ $= -4^2 (x - 1) \checkmark \text{ SHM. } *$	1	marks lost if the best statement was not written. * and transcript errors.
ii) Max speed when $\ddot{x} = 0$ or at the centre of motion $\ddot{x} = -4^2 (\sqrt{3} \cos 4t + \sin 4t)$ $0 = -4^2 (\sqrt{3} \cos 4t + \sin 4t)$ $\sqrt{3} \cos 4t + \sin 4t = 0$ $\tan 4t = -\sqrt{3} \cos 4t$ $\sin 4t = -\sqrt{3}$ $4t = \frac{2\pi}{3}, \frac{5\pi}{3}$ $t = \frac{2\pi}{12}, \frac{5\pi}{12}$ $= \frac{\pi}{6}, \frac{5\pi}{12}$ particle first reaches max. speed when $t = \frac{\pi}{6}$.	1	transcript error caused major error in method student used.

MATHEMATICS EXTENSION I – QUESTION 11

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
d) (i) $RHS = 2(p^2 + q^2) - (p+q)^2$ $= 2p^2 + 2q^2 - p^2 - 2pq - q^2$ $= p^2 + q^2 - 2pq$ $= (p-q)^2$ $= RHS$	/	well done
(ii) coordinates M $\left(\frac{2p+2q}{2}, \frac{p^2+q^2}{2}\right)$ $\left(p+q, \frac{p^2+q^2}{2}\right)$ using part (i) and $p-q=4$ $(p-q)^2 = 2(p^2+q^2) - (p+q)^2$ $x = p+q, y = \frac{p^2+q^2}{2}$	/	generally mistake when made finding the coordinates of midpoint!
$4^2 = 2 \times 2y - x^2$ $x^2 = 4y - 16$ locus of M is the parabola $x^2 = 4y - 16$	/	
(iii) $x^2 = 4y - 16$ $= 4(y-4)$ focal length = 1. Vertex (0, 4) Focus (0, 5)	/	Mark given if correct vertex indicated and focus.
		a=1 not good enough

MATHEMATICS EXTENSION I - QUESTION 12

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
(a) For $8x^3 - 36x^2 + 22x + 21 = 0$		
Let the roots be $\alpha - d, \alpha, \alpha + d$ (where d is the common difference)		This method was the
Sum of roots one at a time: $\alpha - d + \alpha + \alpha + d = \frac{36}{8}$		better one to use to
$3\alpha = \frac{9}{2}$		find the roots and
$\alpha = \frac{3}{2}$	1	for this reason,
Product of the roots: $\alpha(\alpha - d)(\alpha + d) = -\frac{21}{8}$		most of the students
$\alpha(\alpha^2 - d^2) = -\frac{21}{8}$		(who used this method)
Sub $\alpha = \frac{3}{2}$, $\frac{3}{2}[(\frac{3}{2})^2 - d^2] = -\frac{21}{8}$		answered this question
$\frac{9}{4} - d^2 = -\frac{7}{4}$		well.
$d^2 = \frac{9}{4} + \frac{7}{4}$		
$= 4$		
$d = \pm 2$	1	
$\therefore$ roots are: $\frac{3}{2}, \frac{3}{2} - 2, \frac{3}{2} + 2$		
$= \frac{3}{2}, -\frac{1}{2}, \frac{7}{2}$	1	Refer to next page
Alternatively you could use Sum of roots two at a time to find d .		for the alternate
ie $\alpha(\alpha - d) + \alpha(\alpha + d) + (\alpha - d)(\alpha + d) = \frac{22}{8}$		solution where the
$\alpha^2 - \alpha d + \alpha^2 + \alpha d + \alpha^2 - d^2 = \frac{22}{8}$		roots were taken as
$3\alpha^2 - d^2 = \frac{22}{8}$		$\alpha, \alpha + d, \alpha + 2d$.
$d^2 = 4$		
$d = \pm 2$	(3)	

MATHEMATICS EXTENSION I - QUESTION 12

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
a) Alternate solution to 12(a) Let roots be $\alpha, \alpha+d, \alpha+2d$ $\Sigma \alpha: \alpha + \alpha + d + \alpha + 2d = \frac{36}{8}$ $3\alpha + 3d = \frac{9}{2}$ $3(\alpha + d) = \frac{9}{2}$ $\alpha + d = \frac{3}{2}$ $\alpha = \frac{3}{2} - d \quad \text{--- (1)}$ $\Sigma \alpha\beta: \alpha(\alpha+d) + \alpha(\alpha+2d) + (\alpha+d)(\alpha+2d) = \frac{22}{8}$ $\alpha^2 + \alpha d + \alpha^2 + 2\alpha d + \alpha^2 + 3\alpha d + 2d^2 = \frac{22}{8}$ $3\alpha^2 + 6\alpha d + 2d^2 = \frac{22}{8} \quad \text{--- (2)}$ Sub (1) in (2): $3[\frac{3}{2} - d]^2 + 6[(\frac{3}{2} - d)d] + 2d^2 = \frac{22}{8}$ $3(\frac{9}{4} - 3d + d^2) + 9d - 6d^2 + 2d^2 = \frac{22}{8}$ $\frac{27}{4} - 9d + 3d^2 + 9d - 6d^2 + 2d^2 = \frac{22}{8}$ $d^2 = 4$ $d = \pm 2 \quad \text{sub } d=2 \text{ in (1)}$ $\alpha = \frac{3}{2} - 2 = -\frac{1}{2} \quad (\text{or } d=-2)$ and $\alpha + 2d = -\frac{1}{2} + 4 = 3\frac{1}{2}$ and $\alpha + d = \frac{3}{2}$ $\therefore$ roots are $-\frac{1}{2}, \frac{3}{2}, 3\frac{1}{2}$	1 1 1	Many students who used this method struggled with the algebra and the simplification of terms.

MATHEMATICS EXTENSION I – QUESTION 12


SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

12

6)



From $\triangle ABC$, Let $AB = h$

$$\tan 11^\circ 13' = \frac{h}{BC}$$

$$\therefore BC = \frac{h}{\tan 11^\circ 13'} \quad \text{--- (1)}$$

From $\triangle ABD$

$$\tan 8^\circ 14' = \frac{h}{BD}$$

$$BD = \frac{h}{\tan 8^\circ 14'} \quad \text{--- (2)}$$

In $\triangle BCD$, $BD^2 = BC^2 + 500^2$

$$\text{and } BD^2 - BC^2 = 500^2 \quad \text{--- (3)}$$

Sub (1) and (2) in (3)

$$\frac{h^2}{\tan^2 8^\circ 14'} - \frac{h^2}{\tan^2 11^\circ 13'} = 500^2 \quad \text{--- (4)}$$

$$h^2 \left(\frac{1}{\tan^2 8^\circ 14'} - \frac{1}{\tan^2 11^\circ 13'} \right) = 500^2$$

$$h^2 = \frac{500^2}{\frac{1}{\tan^2 8^\circ 14'} - \frac{1}{\tan^2 11^\circ 13'}}$$

$$h = \sqrt{\frac{500^2}{\frac{1}{\tan^2 8^\circ 14'} - \frac{1}{\tan^2 11^\circ 13'}}}$$

$$h = 105.8$$

$\therefore$ height of tower is 106m to the nearest minute

1

1

1/2

1/2

(3)

A few students were not confident in applying the particular method to find the height AB.

Using Pythagoras Theorem was the best formula to use as $\triangle BCD$ was a right-angled $\triangle$. Some students used the cosine rule but did not substitute the correct

angles and sides. Should be:
 $BD^2 = BC^2 + CD^2 - 2(BC)(CD)\cos \angle C$
 $\frac{h^2}{\tan^2 8^\circ 14'} = \frac{h^2}{\tan^2 11^\circ 13'} + 500^2 - 2(BC)(CD)\cos 90^\circ = 0$
 etc.

PTD

MATHEMATICS EXTENSION I – QUESTION 12

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
12b) Examiners Comments: Cont'd. From Equation (4) students found a common denominator making the question more difficult and hence allowed for more errors.		
From (4) $h^2 \frac{(\tan^2 11^\circ 13') - \tan^2 8^\circ 14')}{(\tan^2 11^\circ 13')(\tan^2 8^\circ 14')} = 500^2$	1	
$h^2 = \frac{500^2 \tan^2 11^\circ 13' \tan^2 8^\circ 14'}{\tan^2 11^\circ 13' - \tan^2 8^\circ 14'}$		
$h = \frac{\sqrt{500^2 \tan^2 11^\circ 13' \tan^2 8^\circ 14'}}{\sqrt{\tan^2 11^\circ 13' - \tan^2 8^\circ 14'}}$		
$h = \frac{500 \tan 11^\circ 13' \tan 8^\circ 14'}{\sqrt{\tan^2 11^\circ 13' - \tan^2 8^\circ 14'}}$ $= 106$		

MATHEMATICS EXTENSION I – QUESTION 12

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
Alternate solution 12 b) (But harder and <u>NOT</u> preferred)		
$\tan 11^\circ 13' = \frac{h}{BC}$ and $\tan 8^\circ 14' = \frac{h}{BD}$		
$h = BC \tan 11^\circ 13' \text{ --- (1)}$ and $h = BD \tan 8^\circ 14' \text{ --- (2)}$		
(1) = (2)		
$BC \tan 11^\circ 13' = BD \tan 8^\circ 14'$	1	
$BC = \frac{BD \tan 8^\circ 14'}{\tan 11^\circ 13'}$		
In $\triangle BCD$, $500^2 + BC^2 = BD^2$		
$BD^2 = 500^2 + \frac{BD^2 \tan^2 8^\circ 14'}{\tan^2 11^\circ 13'}$	1	
$BD^2 - \frac{BD^2 \tan^2 8^\circ 14'}{\tan^2 11^\circ 13'} = 500^2$		
$BD^2 (\tan^2 11^\circ 13' - \tan^2 8^\circ 14') = 500^2 \tan^2 11^\circ 13'$		
$BD^2 = \frac{500^2 \tan^2 11^\circ 13'}{\tan^2 11^\circ 13' - \tan^2 8^\circ 14'}$		
$BD = \frac{500 \tan 11^\circ 13'}{\sqrt{\tan^2 11^\circ 13' - \tan^2 8^\circ 14'}}$	1 1/2	
Sub in (2) $h = \frac{500 \tan 11^\circ 13' \times \tan 8^\circ 14'}{\sqrt{\tan^2 11^\circ 13' - \tan^2 8^\circ 14'}}$		
≈ 106	1/2	

MATHEMATICS EXTENSION I – QUESTION 12


SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
12 c) i) If $V = P + A e^{-kt}$ ——— (1)		
then $V - P = A e^{-kt}$ ——— (2)		This question was
From (1) $\frac{dV}{dt} = -k A e^{-kt}$	1	very well done.
$= -k(V - P)$ from (2)		
ii) Initially $t=0$, $V=0$, $P=60$	(1)	
To find A, $V = P + A e^{-kt}$		
$0 = 60 + A e^{-k \times 0}$		
$A = -60$		
Also $t=10$ and $V=35$		
To find k, $35 = 60 + 60 e^{-k \times 10}$	1	
$60 e^{-10k} = 25$		
$e^{-10k} = \frac{25}{60}$		
$-10k = \ln \frac{25}{60}$		
$k = \frac{-1}{10} \ln \frac{25}{60}$	1	
$= \frac{1}{10} \ln \frac{60}{25}$	(2)	
≈ 0.88		

MATHEMATICS EXTENSION I - QUESTION 12

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS



Solution 1.

Let $\angle BAF = \alpha$

$\angle BCA = \angle BAF$ (angle between a tangent and a chord is equal to the angle in the alternate segment).

Also

$\angle BCA = \angle DCA$ (given that AC bisects $\angle BCD$)

$\therefore \angle DCA = \alpha$

Now $\angle DBA = \angle DCA$ (angles in the same segment)

i.e. $\angle DBA = \alpha$

$\therefore \angle BAF = \angle DBA$ (both equal to α)

The alternate angles $\angle BAF$ and $\angle DBA$ are equal

$\therefore$ the two lines are parallel, $\downarrow$

$\therefore EAF \parallel BD$

Note: 1 mark was awarded when 2 angles made a connection with another angle with the correct reason in a logical manner. Max of 2 marks for this]

Note: Nearly all students gave the incorrect reason to this final statement.

They gave the converse to alternate angle theorem which is "alternate angles on parallel lines are equal" However not penalised for this.

This question was not well done.

When proving in Geometry it is important to communicate in a logical sequential manner. Marks were not awarded when this did not occur.

Also, the only given facts were that AC bisects $\angle DCB$ and that ABCD was a cyclic Quadrilateral.

PTO $\rightarrow$

3

MATHEMATICS EXTENSION I – QUESTION 12

SUGGESTED SOLUTIONS

QUESTIONS	MARKS
1. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
2. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
3. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
4. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
5. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
6. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
7. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
8. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
9. The following are the marks obtained by 10 students in a test. Find the mean marks.	10
10. The following are the marks obtained by 10 students in a test. Find the mean marks.	10

MARKER'S COMMENTS

Q12 d) Alternate solution 2

Let $\angle EAD = \theta$.

$\angle ABD = \angle EAD$ (angle between tangent and
= θ a chord is equal to the angle in the alternate
segment)

$$\angle DCA = \angle ABD \text{ (angles in the same segment)}$$

Also $\angle BCA = \angle DCA$ (CA bisects $\angle BCD$)
 $= 40^\circ$

Now $\angle DCB = 2\theta$

Also $\angle DAB = 180 - \angle DCB$ (opposite angles in a cyclic quadrilateral)
 $= 180 - 20$

$$\begin{aligned}\text{Now } \angle BAF &= 180 - \angle EAD - \angle DAB \quad (\text{angle sum of straight line}) \\ &= 180 - \theta - (180 - 2\theta) \quad (\text{shown above}) \\ &= \theta\end{aligned}$$


$\therefore \angle ADB = \angle BAF$ (angles in the alternate segment)

$$\therefore \angle EAD = \angle ADR \quad (\text{both equal to } \theta)$$

Alternate angles $\angle EAO$ and $\angle ADB$ are equal in

$\therefore$ two lines are parallel

ie $EF \parallel BD$.



Many students made assumptions from the

information that

CA bisects $\angle BCD$.

They assumed:

- AC is a diameter

$$-\angle EAC = 90^\circ$$

(the angle between tangent and radius is 90°)

$$\underline{-AC \perp DC}$$

and $\angle ACB \cong \angle DAB$

- AC bisects $\angle DIB$

Although these may seem to be true, you would need to prove them first, then it would be OK to use.

By making these assumptions, no marks were awarded.

3


MATHEMATICS EXTENSION I – QUESTION 13

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
i Let $\sin x - \cos x = R \sin(x - \alpha)$ $= R \sin x \cos \alpha - R \cos x \sin \alpha$		One mark each for finding R and α .
$\therefore R \cos \alpha = 1$ $R^2 \cos^2 \alpha = 1$		Simply stating " $R = \sqrt{2}$ " or " $\alpha = \frac{\pi}{4}$ " earned no marks, as this information is provided in the question
$R \sin \alpha = 1$ $R^2 \sin^2 \alpha = 1$		
So $R^2 \cos^2 \alpha + R^2 \sin^2 \alpha = 1 + 1$ $R^2 (\sin^2 \alpha + \cos^2 \alpha) = 2$ $R^2 = 2$ $R = \sqrt{2} \quad (R > 0)$	1	
$\therefore \sqrt{2} \cos \alpha = 1$ $\cos \alpha = \frac{1}{\sqrt{2}}$		
$\therefore \sqrt{2} \sin \alpha = 1$ $\sin \alpha = \frac{1}{\sqrt{2}}$		
$\therefore \alpha = \frac{\pi}{4}$	1	
$\therefore \sin x - \cos x = \sqrt{2} \sin \left(x - \frac{\pi}{4}\right)$		
<u>OR</u> RHS = $\sqrt{2} \sin \left(x - \frac{\pi}{4}\right)$ $= \sqrt{2} \left(\sin x \cos \frac{\pi}{4} - \cos x \sin \frac{\pi}{4}\right)$ $= \sqrt{2} \left(\sin x \cdot \frac{1}{\sqrt{2}} - \cos x \cdot \frac{1}{\sqrt{2}}\right)$ $= \sqrt{2} \times \frac{1}{\sqrt{2}} (\sin x - \cos x)$ $= \sin x - \cos x$ $= \text{LHS}$		

MATHEMATICS EXTENSION I – QUESTION 13 (continued)

a)

ii



$$\begin{aligned} \text{iii} \quad \sin x - \cos x &= 1 \\ \therefore \sqrt{2} \sin\left(x - \frac{\pi}{4}\right) &= 1 \\ \sin\left(x - \frac{\pi}{4}\right) &= \frac{1}{\sqrt{2}} \\ \therefore x - \frac{\pi}{4} &= \frac{\pi}{4}, \frac{3\pi}{4} \dots \\ \therefore x &= \frac{\pi}{2}, \pi, \dots \quad \therefore \frac{\pi}{2} \text{ is a solution} \end{aligned}$$

$$\begin{aligned} \underline{\text{OR}} \quad \text{when } x &= \frac{\pi}{2}, \text{ LHS} = \sin\left(\frac{\pi}{2}\right) - \cos\left(\frac{\pi}{2}\right) \\ &= 1 - 0 \\ &= 1 \\ &= \text{RHS} \quad \therefore \frac{\pi}{2} \text{ is a solution} \end{aligned}$$

$$\therefore \frac{\pi}{2} < x < \pi$$

($\frac{1}{2}$ mark for $\frac{\pi}{2} \leq x \leq \pi$)

MARKS

MARKER'S COMMENTS

$\frac{1}{2}$

Intercepts correct and labelled

$\frac{1}{2}$

Amplitude

$\frac{1}{2}$

Correct shift / phase

$\frac{1}{2}$

Period

For a "show" question, anything you are given in the question is worth zero marks.

1

Eg. simply stating $\sin\left(\frac{\pi}{2}\right) - \cos\left(\frac{\pi}{2}\right) = 1$ does not show the result – the truth of this statement is provided in the question itself.

1

Also note that a rough sketch is insufficient to show this result – do it algebraically

MATHEMATICS EXTENSION I – QUESTION 13

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
b) Let $X = \sin^{-1} \frac{1}{\sqrt{5}}$ and $Y = \sin^{-1} \frac{1}{\sqrt{10}}$		
then		
$\sin X = \frac{1}{\sqrt{5}}$ and $\sin Y = \frac{1}{\sqrt{10}}$	1	
		
$\sin(X+Y) = \sin X \cos Y + \cos X \sin Y$		
$= \frac{1}{\sqrt{5}} \times \frac{3}{\sqrt{10}} + \frac{2}{\sqrt{5}} \times \frac{1}{\sqrt{10}}$	1	
$= \frac{5}{5\sqrt{2}}$		
$= \frac{1}{\sqrt{2}}$		
$X+Y = \frac{\pi}{4}$	1	
$\therefore \sin^{-1} \frac{1}{\sqrt{5}} + \sin^{-1} \frac{1}{\sqrt{10}} = \frac{\pi}{4}$		

MATHEMATICS EXTENSION I – QUESTION 13 (continued)

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

c) Prove the statement true for $n=1$

$$7^1 + 5 = 12$$

$= 6(2)$ which is divisible by 6

1

Correct proof for $n=1$

Let k be a positive integer for which the statement is true

i.e. $7^k + 5 = 6M$, where M is an integer

$$7^k = 6M - 5$$

1

Correct assumption, including the essential requirement that M be an integer

Prove that if the statement is true for $n=k$, then it is true for $n=k+1$

i.e. $7^{k+1} + 5$ is divisible by 6

$$7^{k+1} + 5 = 7(7^k) + 5$$

$$= 7(6M - 5) + 5, \text{ by the induction hypothesis}$$

$$= 42M - 35 + 5$$

$$= 42M - 30$$

$$= 6(7M - 5) \text{ which is divisible by 6}$$

1

Many students made their working needlessly complicated by attempting to prove $7^{k+1} + 5 = 6P$ without knowing what P would be (or worse, $7^{k+1} + 5 = 6M$, which is impossible given the assumption).

Therefore if the statement is true for $n=k$ it is also true for $n=k+1$. Since it is true for $n=1$, it is true for $n=2$ and so on. Therefore $7^n + 5$ is divisible by 6 for all positive integer values of n

EXTENSION 1

MATHEMATICS - QUESTION NO: 14

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

$$14(a) \quad V = \frac{1}{3} \pi r^2 h \quad \text{with } h = r \text{ (by similarity)}$$

$$\therefore V = \frac{1}{3} \pi r^3$$

$$\frac{dV}{dr} = \pi r^2$$

$$\frac{dV}{dt} = \frac{dV}{dr} \frac{dr}{dt}$$

$$24 = \pi r^2 \frac{dr}{dt}$$

$$\therefore \frac{dV}{dt} = \frac{24}{\pi r^2}$$

$$S = \pi r^2$$

$$\frac{dS}{dr} = 2\pi r$$

$$\frac{dS}{dt} = \frac{dS}{dr} \frac{dr}{dt}$$

$$= 2\pi r \frac{24}{\pi r^2} \text{ (from above)}$$

$$= \frac{48}{r}$$

$$\text{When } r = 16$$

$$\frac{dS}{dt} = 3 \text{ cm}^2 \text{ s}^{-1}$$

4

There were a multitude of variations to the solution of this question. I read carefully through all the solutions, and allocated marks by subtracting 1 for each error ^{and} or omission of steps. No $\frac{1}{2}$ marks were awarded. The most common errors were of the type

$$V = \frac{1}{3} \pi r^2 h \quad \left. \begin{array}{l} \text{WRONG} \\ \frac{dV}{dr} = \frac{1}{3} \pi r^2 \end{array} \right\} \quad \left. \begin{array}{l} V = \frac{1}{3} S \\ \frac{dV}{dr} = \frac{1}{3} \end{array} \right\} \text{WRONG}$$

$$\text{(or)} \quad S = \pi r^2 + \pi r h \quad \left. \begin{array}{l} \text{WRONG} \\ \frac{dS}{dr} = 2\pi r + \pi r \end{array} \right\} \text{WRONG}$$

(many more variations)

STRESS TO STUDENTS that

if h and r are variables related to one another, not constants, must be differentiated with differentiations

MATHEMATICS - QUESTION NO: 14

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

14 (b)(i) $(1+ax)^n = 1 + {}^nC_1 ax + {}^nC_2 a^2 x^2 + \dots$
 $= 1 + 6x + 16x^2 + \dots$

2

A number of students thought the line marked * was the solution to part (a) but did ① and ② in part (b), no penalty. Marks were awarded by deduction for errors and/or omissions, no $\frac{1}{2}$ marks were given.

$${}^nC_1 a = 6$$

$$\therefore na = 6 \text{ ①}$$

$${}^nC_2 a^2 = 16$$

$$\frac{n(n-1)a^2}{2} = 16$$

$$n(n-1)a^2 = 32 \text{ ②}$$

* ←

1 mark awarded for this line

(ii) From (i) $a = \frac{6}{n}$

$$\therefore n(n-1) \frac{36}{n^2} = 32$$

$$\therefore (n-1) \cdot 36 = 32n$$

$$36n - 36 = 32n$$

$$4n = 36$$

$$n = 9$$

Sub in ① $a = \frac{2}{3}$

2

MATHEMATICS - QUESTION NO: 14

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

14 (c) (i) $\ddot{x} = -4(x-1)$

2

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = -4(x-1)$$

$$\frac{1}{2} v^2 = -2(x-1)^2 + c$$

When $x = 1, v = 6, c = 18$

$$\therefore \frac{1}{2} v^2 = -2(x-1)^2 + 18$$

$$v^2 = -4(x-1)^2 + 36 = -4x^2 + 8x - 4 + 36$$

$$v^2 = -4x^2 + 8x + 32 \text{ as required}$$

Loss of 1 mark
for using $x=0$
 $v=6$.



A lot of students had $x=0, v=6$ and found 'c' incorrectly, then manipulated to get 32 incorrectly

(ii) $v^2 = -4(x^2 - 2x - 8)$
 $= -4[(x-1)^2 - 9]$

2

$$= 4[9 - (x-1)^2]$$

of the form

$$v^2 = n^2[a^2 - (x-b)^2] \text{ where } x=b \text{ is the centre of motion}$$

$$n = 2 \quad T = \frac{2\pi}{n} = \pi$$

$$a^2 = 9 \quad a = 3 \text{ since } a > 0$$

Many students did this correctly, but those who got 'a' incorrect lost 1 mark even if some of the working was correct. No $\frac{1}{2}$ marks were awarded. Most got $T = \pi$

MATHEMATICS EXTENSION I – QUESTION 15

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
a) $\sin^{-1}x > 0$ as $\log_e k$ defined only for $k > 0$ $\sin^{-1}x \leq \frac{\pi}{2}$ $x \leq 1$ $\therefore$ Domain is $0 < x \leq 1$	2	Some did not look at where $\log_e x$ is defined. Others did not define the domain. Overall well done.
b) (i) Vertical $y = -10t$ $y = -10t + c_1$ when $t = 0$, $y = V \cos \theta$ $\therefore c_1 = V \cos \theta$ $y = -10t + V \cos \theta$ $y = \frac{-10t^2}{2} + V \cos \theta t + c_2$ when $t = 0$ $y = 0 \Rightarrow c_2 = 0$ $y = -5t^2 + V \cos \theta t$	2	Mostly done well. Some students did not give enough supporting evidence.
(ii) Greatest height when $y = 0$ and $t = 3$ $y = -10 \times 3 + V \sin \theta$ $0 = -30 + V \sin \theta$ $V \sin \theta = 30$	1	Most student did this well.

MATHEMATICS EXTENSION I – QUESTION 15

[illegible]

MATHEMATICS EXTENSION I – QUESTION 15

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
part c) (i) LHS = $x[1 + (1+x) + (1+x)^2 + \dots + (1+x)^{n-1}]$ GP with $a=1, r=1+x, n$ terms $S_n = \frac{1((1+x)^n - 1)}{1+x-1}$ $= \frac{(1+x)^n - 1}{x}$ $= x \times \left[\frac{(1+x)^n - 1}{x} \right]$ $= (1+x)^n - 1$ $= RHS$	1	Most students had difficulty with this question. They did not recognise the GP
(ii) RHS = $(1+x)^n - 1$ $= \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots + \binom{n}{k}x^k + \dots + \binom{n}{n}x^n - 1$ coefficient of x^k is $\binom{n}{k}$ LHS = $[x(1+x)^{n-1}] + [x(1+x)^{n-2}] + \dots + [x(1+x)^{k+1}] + \dots$ $= \left[\binom{n-1}{0}x + \binom{n-1}{1}x^2 + \dots + \binom{n-1}{k-1}x^k + \dots \right]$ $+ \left[\binom{n-2}{0}x + \binom{n-2}{1}x^2 + \dots + \binom{n-2}{k-1}x^k + \dots \right] + \dots$ $+ \left[\binom{k-1}{0}x + \binom{k-1}{1}x^2 + \dots + \binom{k-1}{k-1}x^k + \dots \right]$ coefficient of x^k is: $\binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{k-1}{k-1}$ $\therefore \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{k-1}{k-1} = \binom{n}{k}$	1	This question was quite poorly set out by many students.

MATHEMATICS - QUESTION NO: 14

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

$$14 (b)(i) (1+ax)^n = 1 + {}^nC_1 ax + {}^nC_2 a^2 x^2 + \dots$$

$$= 1 + 6x + 16x^2 + \dots$$

2

A number of students thought the line marked * was the solution to part (a) but did ① and ② in part (b), no penalty. Marks were awarded by deduction for errors and/or omissions, no $\frac{1}{2}$ marks were given.

$${}^nC_1 a = 6$$

$${}^nC_2 a^2 = 16$$

* ←

1 mark awarded for this line

$$\therefore na = 6 \text{ ①}$$

$$\frac{n(n-1)a^2}{2} = 16$$

$$n(n-1)a^2 = 32 \text{ ②}$$

$$(ii) \text{ From (i) } a = \frac{6}{n}$$

2

$$\therefore \frac{n(n-1)36}{n^2} = 32$$

$$\therefore (n-1) \cdot 36 = 32n$$

$$36n - 36 = 32n$$

$$4n = 36$$

$$n = 9$$

$$\text{Sub in ① } a = \frac{2}{3}$$

MATHEMATICS - QUESTION NO: 14

SUGGESTED SOLUTIONS

MARKS

MARKER'S COMMENTS

14 (c) (i) $\ddot{x} = -4(x-1)$

2

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = -4(x-1)$$

$$\frac{1}{2} v^2 = -2(x-1)^2 + c$$

When $x = 1, v = 6, c = 18$

$$\therefore \frac{1}{2} v^2 = -2(x-1)^2 + 18$$

$$v^2 = -4(x-1)^2 + 36 = -4x^2 + 8x - 4 + 36$$

$$v^2 = -4x^2 + 8x + 32 \text{ as required}$$

Loss of 1 mark
for using $x=0$
 $v=6$.



A lot of students had $x=0, v=6$ and found 'c' incorrectly, then manipulated to get 32 incorrectly

(ii) $v^2 = -4(x^2 - 2x - 8)$
 $= -4[(x-1)^2 - 9]$

2

$$= 4[9 - (x-1)^2]$$

of the form

$$v^2 = n^2[a^2 - (x-b)^2] \text{ where } x=b \text{ is the centre of motion}$$

$$n = 2 \quad T = \frac{2\pi}{n} = \pi$$

$$a^2 = 9 \quad a = 3 \text{ since } a > 0$$

Many students did this correctly, but those who got 'a' incorrect lost 1 mark even if some of the working was correct. No $\frac{1}{2}$ marks were awarded. Most got $T = \pi$

MATHEMATICS EXTENSION I – QUESTION 15

SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
a) $\sin^{-1}x > 0$ as $\log_e k$ defined only for $k > 0$ $\sin^{-1}x \leq \frac{\pi}{2}$ $x \leq 1$ $\therefore$ Domain is $0 < x \leq 1$	2	Some did not look at where $\log_e x$ is defined. Others did not define the domain. Overall well done.
b) (i) Vertical $y = -10t$ $y = -10t + c_1$ when $t = 0$, $y = V \cos \theta$ $\therefore c_1 = V \cos \theta$ $y = -10t + V \cos \theta$ $y = \frac{-10t^2}{2} + V \cos \theta t + c_2$ when $t = 0$ $y = 0 \Rightarrow c_2 = 0$ $y = -5t^2 + V \cos \theta t$	2	Mostly done well. Some students did not give enough supporting evidence.
(ii) Greatest height when $y = 0$ and $t = 3$ $y = -10 \times 3 + V \sin \theta$ $0 = -30 + V \sin \theta$ $V \sin \theta = 30$	1	Most student did this well.

MATHEMATICS EXTENSION I - QUESTION 15		
SUGGESTED SOLUTIONS	MARKS	MARKE R'S COMMENTS
b) (iii) Stone reaches the ground when $y = -80$ $y = -5t^2 + V \sin \theta t$ $V \sin \theta = 30$ $-80 = -5t^2 + 30t$ $t^2 - 6t - 16 = 0$ $(t-8)(t+2) = 0$ $\therefore t = 8$ since $t \geq 0$	1	Some students took ground level as $y=0$. (They need to read the question carefully)
(iv) Stone reaches the ground at a range of 320m after 8 seconds $x = V \cos \theta t$ $x = 320$, $t = 8$ $320 = 8V \cos \theta$ $V \cos \theta = 40$	1	Some students had $320 \div 6 = 40$ others had $320 \div 80 = 40$
(v) $V^2 = x^2 + y^2$ $\tan \theta = \frac{V \sin \theta}{V \cos \theta}$ $= (V \cos \theta)^2 + (V \sin \theta)^2$ $= 40^2 + 30^2$ $= 2500$ $V = 50 \text{ ms}^{-1}$	1	Very well done by most that attempted it.
		$\theta = 36^\circ 52'$ (nearest minute)

b) (iii) Stone reaches the ground when $y = -80$	Some students took
--	--------------------

$$y = -5t^2 + v \sin \theta t \quad v \sin \theta = 30 \quad | \quad \text{ground level as } y=0$$

$$-80 = -5t^2 + 30t$$

$$t^2 - 6t - 16 = 0$$

$$(t-8)(t+2) = 0$$

$\therefore t = 8$ since $t \geq 0$

(iv) Stone reaches the ground at a range of 320m after 8 seconds	1	Some students had $320 \div 6 = 40$
--	---	-------------------------------------

$x = V \cos \theta t$	$x = 320, t = 8$	others had
-----------------------	------------------	------------

$$320 = 8 \sqrt{\cos A} \quad \quad \quad 320 \div 80 = 40$$

$$V \cos \theta = 40$$

(v) $V^2 = x^2 + y^2$ $\tan \theta = \frac{V_{\sin \theta}}{V_{\cos \theta}}$ Very well done by

$$= (V \cos \theta)^2 + (V \sin \theta)^2$$

$$= 40^2 + 30^2 = 30$$

$$= 2500 \quad \theta = 36^\circ 52'$$

$V = 50 \text{ ms}^{-1}$ (nearest minute)

$\tan \theta = \frac{V \sin \theta}{V \cos \theta}$	Very well done by
---	-------------------

11	30	40	it.
----	----	----	-----

$$\theta = 36^\circ 52'$$

(nearest minute)	
------------------	--

Some students took ground level as $y=0$ (They need to read the question carefully)

Some students
had $320 \div 6 = 40$
others had
 $320 \div 80 = 40$

Very well done by most that attempted it.

MATHEMATICS EXTENSION I – QUESTION 15

MATHEMATICS EXTENSION I – QUESTION 15		
SUGGESTED SOLUTIONS	MARKS	MARKER'S COMMENTS
$c) (iii) \quad n \binom{n-1}{k} = \frac{n(n-1)!}{k!(n-1-k)!}$ $= \frac{n!}{k!(n-(k+1))!}$ $(k+1) \binom{n}{k+1} = (k+1) \frac{n!}{(k+1)!(n-(k+1))!}$ $= (k+1) \frac{n!}{(k+1)k!(n-(k+1))!}$ $= \frac{n!}{k!(n-(k+1))!}$ $\therefore n \binom{n-1}{k} = (k+1) \binom{n}{k+1}$	2	Done very well by most students that attempted it