



**BAULKHAM HILLS  
HIGH  
SCHOOL**

**2018**

**YEAR 11  
HALF YEARLY  
ASSESSMENTS**

# Mathematics Extension 1

---

## **General Instructions**

- Reading time – 5 minutes
- Working time – 2 hours 30 minutes
- Write using black or blue pen  
Black pen is preferred
- Board-approved calculators may be used
- In Questions 11 – 18, show relevant mathematical reasoning and/or calculations
- Marks may be deducted for careless or badly arranged work

---

## **Total marks: 130**

### **Section I – 10 marks** (pages 2 – 5)

- Attempt Questions 1 – 10
- Allow about 15 minutes for this section

### **Section II – 120 marks** (pages 6 – 18)

- Attempt Questions 11 – 18
- Allow about 2 hours 15 minutes for this section

## Section I

10 marks

Attempt Questions 1 – 10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1 – 10

---

1 Which one of the following expressions represents the factored form of  $8x^3 + 27$  ?

(A)  $(2x + 3)(4x^2 + 6x + 9)$


(B)  $(2x + 3)(4x^2 - 6x + 9)$

(C)  $(2x - 3)(4x^2 - 6x - 9)$


(D)  $(2x - 3)(4x^2 + 6x - 9)$

2 Which of the following is a function?


(A)




(B)



(C)



(D)



3  $\frac{1 + \sqrt{3}}{2 - \sqrt{3}} =$


(A)  $5 + 3\sqrt{3}$

(B)  $\frac{5 + 3\sqrt{3}}{7}$

(C)  $\sqrt{3} - 1$

(D)  $\frac{\sqrt{3} - 1}{7}$

4



In the diagram above;

- (A) The angle of elevation is  $\alpha$  and the angle of depression is  $\beta$
- (B) The angle of elevation is  $\beta$  and the angle of depression is  $\theta$
- (C) The angle of elevation is  $\theta$  and the angle of depression is  $\beta$
- (D) The angle of elevation is  $\alpha$  and the angle of depression is  $\theta$

5 Given  $\cos\theta = -\frac{3}{5}$  and  $\sin\theta < 0$ , the ratio for  $\tan\theta =$

- (A)  $-\frac{4}{5}$
- (B)  $-\frac{4}{3}$
- (C)  $\frac{4}{3}$
- (D)  $\frac{4}{5}$

- 6 In simplified form, the algebraic expression  $\frac{x+5}{(x-3)(x+1)} - \frac{x-1}{x^2-x-2}$  can be written as

(A)  $\frac{7x-13}{(x+1)(x-2)(x-3)}$

(B)  $\frac{-x-13}{(x+1)(x-2)(x-3)}$

(C)  $\frac{7x-7}{(x+1)(x-2)(x-3)}$

(D)  $\frac{-x-7}{(x+1)(x-2)(x-3)}$

7  $\frac{\sin(360^\circ - A)}{\sin(90^\circ - A)} =$

(A)  $-1$

(B)  $1$

(C)  $-\tan A$

(D)  $\tan A$

- 8 Two functions are defined as  $f(x) = 3x^2 - 4$  and  $g(y) = y^2 - 2y$ . How many solutions are there to the equation  $f(a) = g(2a)$  ?

(A)  $0$

(B)  $1$

(C)  $2$

(D)  $3$

- 9 The circle with equation  $x^2 + y^2 - 12x - 10y + k = 0$  meets the coordinate axes at exactly three points.

What is the value of  $k$  ?

- (A) 5  
(B) 6  
(C) 25  
(D) 36
- 10 Let  $r$  and  $s$  be integers, then  $\frac{6^{r+s} \times 12^{r-s}}{8^r \times 9^{r+2s}}$  is an integer if
- (A)  $r + s \leq 0$   
(B)  $s \leq 0$   
(C)  $r \leq 0$   
(D)  $r \geq s$

**END OF SECTION I**

## Section II

120 marks

Attempt Questions 11 – 18

Allow about 2 hours 15 minutes for this section

Answer each question on the appropriate answer sheet. Each answer sheet must show your name. Extra paper is available.

In Questions 11 to 18, your responses should include relevant mathematical reasoning and/or calculations.

---

**Question 11** (15 marks) Use a *separate* answer sheet

**Marks**

- (a) Write  $\sqrt[4]{x^5}$  in index form. 1
- (b) Evaluate  $\sqrt{\frac{4.81 \times 10^5}{7.36 \times 10^9}}$  correct to two significant figures. 2
- (c) If  $f(x) = 7 - 2x^2$ , find the value of  $f(-1)$ . 1
- (d) Simplify  $\sqrt{75} - 2\sqrt{27}$ . 2
- (e) Expand and simplify  $(3x - 4)(x - 2)(x + 2)$ . 2
- (f) Solve the inequation  $\frac{2}{x-1} \leq 1$ . 3
- (g) Factorise
- (i)  $2x^2 + 3x - 2$  1
- (ii)  $x^3 + 5x^2 + x + 5$  1
- (iii)  $4a^2(x^3 + 18ab^2) - (32a^5 + 9b^2x^3)$  2

**Question 12 (15 marks)** Use a *separate* answer sheet

(a) Express as a single fraction in simplest terms

(i)  $\frac{125a^3 - 8}{a^2 - 7a + 10} \times \frac{a - 5}{25a^2 - 4}$  2

(ii)  $\frac{2}{x^2 - 1} - \frac{1}{x^2 - x} + \frac{x - 1}{x^2 + x}$  3

(b) Solve

(i)  $\frac{3x - 2}{4} - \frac{2x - 1}{8} = 5$  2

(ii)  $|2x + 6| = 3x - 1$  3


(iii) Solve  $2x^2 - 6x \leq 0$  2

(iv)  $\frac{x + 1}{x^2 - 4} \geq 0$  3

**Question 13 (15 marks)** Use a *separate* answer sheet

- (a) The graph below shows the absolute value function  $y = |2x - 3|$  and the straight line  $y = \frac{9-x}{2}$ .

1



Use the graph to solve the inequation  $|2x - 3| \geq \frac{9-x}{2}$ .

- (b) Rationalise the denominator of  $\frac{10}{\sqrt{5} - \sqrt{3}}$ .

2

- (c) Express  $0.01\dot{2}\dot{3}$  as a fraction in its simplest form, without the aid of a calculator.

2

- (d) Make  $v$  the subject of the formula  $F = \frac{mv^2}{gr}$

2

- (e) Use the method of “completing the square” to solve the equation

2

$$x^2 - 8x + 9 = 0$$

**Question 13 continues on page 9**

**Question 13** (continued)

(f) For the equation  $(x + 2y)(2x - y) + (x - y)(3x + 4y) = 22$

(i) Verify that  $x = 2$  and  $y = 1$  is a possible solution 1

(ii) Find any other value(s) of  $x$  which makes the equation true when  $y = 1$  2

(g) Consider the function  $f(x) = x^2 - y^2 + x + 3y - 2$ .

(i) Show that  $f(x) = (x - y + 2)(x + y - 1)$  1

(ii) Hence, or otherwise, sketch the region  $x^2 - y^2 + x + 3y > 2$ . 2


**End of Question 13**

**Question 14 (15 marks)** Use a *separate* answer sheet

- (a) The diagram below can be described by the statement;

1

“B divides AP in the ratio 1:3”



Write down two different statements, other than the one given above, that would also describe the diagram

- (b) Consider the function
- $y = f(x)$

- (i) State the condition for the function to be even.

1

- (ii) Give an example of an even function.

1

- (c) The function
- $f(x)$
- is defined as

2

$$f(x) = \begin{cases} x^2 + 1 & : x > 3 \\ 3x & : -2 \leq x \leq 3 \\ 2 & : x < -2 \end{cases}$$

Find  $f(2) + f(5) - f(-2)$ .

- (d) Consider the function
- $y = \sqrt{x^2 - 16}$

- (i) Find the domain of the function.

1


- (ii) Find the range of the function.

1

**Question 14 continues on page 11**

**Question 14** (continued)

- (e) A circle, centred at the origin and an exponential of the form  $y = a^x$  are shown below.



$P$  and  $Q$  lie on the curves as shown,  $Q$  has coordinates  $(2,4)$

- |                                                                            |   |
|----------------------------------------------------------------------------|---|
| (i) Write down the coordinates of $P$ .                                    | 1 |
| (ii) Find the equation of the circle.                                      | 2 |
| (iii) Find the equation of the exponential function.                       | 1 |
| (iv) Give the three inequations which combine to define the shaded region. | 2 |
- 
- (f) Is the function  $f(x) = |x - 2| - |x + 2|$  odd, even or neither?  
Justify your answer.
- 2

**End of Question 14**

**Question 15 (15 marks)** Use a *separate* answer sheet

(a) It is known that  $\sin 15^\circ = \frac{\sqrt{3}-1}{2\sqrt{2}}$ .

Using this value, find the following;

(Note: your working *MUST* show how you used  $\sin 15^\circ$ )

(i)  $\cos 75^\circ$ . 1

(ii)  $\operatorname{cosec} 15^\circ$ . 1

(iii)  $\sin 195^\circ$ . 1

(b) Solve the simultaneous equations 2

$$\begin{aligned} 2x + y &= 4 \\ 5x + 2y &= 9 \end{aligned}$$

(c) Prove the relationship  $\sin^4 x - \cos^4 x \equiv 1 - 2\cos^2 x$ . 2

(d) A student is asked to solve  $\tan x = \sin x$ , giving all solutions in the range  $0^\circ \leq x \leq 360^\circ$ .  
Below is the student's working;

$$\begin{aligned} \tan x &= \sin x \\ \frac{\sin x}{\cos x} &= \sin x && \text{(write } \tan x \text{ as } \frac{\sin x}{\cos x}) \\ \sin x &= \sin x \cos x && \text{(multiply by } \cos x) \\ 1 &= \cos x && \text{(cancel } \sin x) \\ x &= 0^\circ, 360^\circ \end{aligned}$$

Whilst the student did find two correct answers, there is another answer that they did not find.

(i) What did the student do wrong in their working? 1

(ii) Find the missing answer. 1


(e) Solve, to the nearest degree where necessary, for  $0^\circ \leq \theta \leq 360^\circ$

(i)  $\sin \theta \tan \theta + 2 \sin \theta = 3 \cos \theta$ . 3

(ii)  $\sin(20^\circ - 2\theta) = \frac{1}{7}$ . 3

**Question 16 (15 marks)** Use a *separate* answer sheet


- (a) The diagram below represents a framework for supporting a roof. The outer beams TP and TS are each 4 metres in length, the inner beams TQ and TR are inclined at  $65^\circ$  to the horizontal and the height of the framework is 2.5 metres.



Calculate the;

- |                                                        |   |
|--------------------------------------------------------|---|
| (i) angle TPS, correct to the nearest degree           | 1 |
| (ii) width PS, correct to the nearest centimetre.      | 1 |
| (iii) length of TQ, correct to the nearest centimetre. | 1 |

- (b) A field is triangular, with two sides of 80 metres and 40 metres, enclosing an angle of  $130^\circ$ .




- |                                                                                                                    |   |
|--------------------------------------------------------------------------------------------------------------------|---|
| (i) Calculate the area of the field, correct to the nearest metre squared.                                         | 1 |
| (ii) Without calculating its length, how do you know that the third side must be the largest side of the triangle? | 1 |
| (ii) Use the cosine rule to calculate the length of the third side, correct to the nearest metre.                  | 1 |


**Question 16 continues on page 14**

**Question 16** (continued)

- (c) A and B are two points 1500 metres apart on a road running due west. A soldier at A observes that the bearing of an enemy's gun battery at G is  $296^\circ$  and heads towards B, where the bearing of the gun battery is  $302^\circ$ .



- (i) Copy this diagram onto your answer sheet, showing the given information. 1
  - (ii) Explain why  $\angle AGB = 6^\circ$  1
  - (iii) Show that  $BG = \frac{1500 \sin 26^\circ}{\sin 6^\circ}$  1
  - (iv) The range of the guns in the enemy's battery is 5 km. How far past B can the soldier travel before being in range of the enemy's guns? 3  
Give your answer correct to the nearest metre.
- (d) Inside a large semicircle with centre O and diameter AOB of length  $2R$ , a smaller semicircle is drawn with diameter AO and centre P. A small circle is drawn with centre Q and radius  $r$  which is tangent to the large semicircle at T and the small semicircle at S. Let  $\angle POQ = \theta$ . 3  
(Note: P, S and Q and O, Q and T are collinear)




Show that  $\frac{r}{R} = \frac{1 - \cos \theta}{3 - \cos \theta}$

**End of Question 16**

**Question 17 (15 marks)** Use a *separate* answer sheet

- (a) Given the points  $A(-2,4)$  and  $B(3,11)$ , find the coordinates of the point  $P$ , which divides the interval  $AB$  in the ratio 2:3. 2

- (b) The base of the triangular prism shown below is a rectangle 8 cm long and 6 cm wide. The vertical faces  $ABC$  and  $PQR$  are equilateral triangles.



Calculate, to the nearest degree;

- (i) the angle between the plane  $PBC$  and the base. 2
- (ii) the angle between the diagonal  $PC$  and the base. 2
- (c) Find the vertical and horizontal asymptotes of the graph of the function; 2

$$y = \frac{x^2}{x^2 + 5x + 6}$$

- (d) It is known that the difference of two squares factorisation is given by; 1

$$a^2 - b^2 = (a - b)(a + b)$$

This factorisation can be generalised for higher powers to;

$$a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + \dots + b^{n-1})$$

Using this generalisation, or otherwise, determine whether  $3^{2018} - 2^{2018}$  is a prime number.

**Question 17 continues on page 16**

**Question 17** (continued)

(e) The point  $K(x,y)$  divides the interval joining  $M(1,2)$  and  $N(-1,-4)$  in the ratio  $k:1$ .

(i) Write expressions for  $x$  and  $y$  in terms of  $k$ .


1

(ii) Find the value of  $k$ , such that the line  $3x - 4y - 5 = 0$  divides  $MN$  in the ratio  $k:1$ .

2

(f)

3




An aeroplane is sighted simultaneously from two stations A and B, B being one kilometre north-east of A. To the observer at A the aeroplane appears due north at an elevation of  $34^\circ$  ; to the observer at B it appears in a direction  $N72^\circ W$  .

Find the height of the aeroplane, correct to the nearest metre..

**End of Question 17**

**Question 18 (15 marks)** Use a *separate* answer sheet

- (a) A railway carriage compartment consists of two bench seats facing each other, with each bench being able to accommodate four people.




- (i) How many ways can a party of eight people be seated in this compartment? 1
- (ii) How many ways can the eight passengers take their seats if Aiden and Alex must sit next to each other? 2
- (b) There are thirteen boys and nine girls in a class. From this class, five students are randomly selected to form the class committee.
- (i) How many possibilities are there for the committee? 1
- (ii) How many of the committees would contain at least one boy and one girl? 2
- (iii) The committee is to be seated around a table. In how many ways could the committee be seated? 1
- (c) Consider the letters of **ZOOLOGICAL**.
- (i) How many distinct arrangements of all of the letters are possible? 1
- (ii) How many arrangements are possible if no letter **O** is placed next to another letter **O**? 2

**Question 18 continues on page 18**

**Question 18** (continued)

(d)

2





There are ten points in a plane, four of which are collinear. How many triangles can be found using these points?

- (e) If you have one white, two blue and three red flags, find how many signals you can make, each containing four flags arranged one above the other. 3


**End of paper**

**BAULKHAM HILLS HIGH SCHOOL**


**YEAR 11 MATHEMATICS EXTENSION HALF YEARLY ASSESSMENTS 2018 SOLUTIONS**

| Solution                                                                                                                                                                                                                                                                                                                 | Marks    | Comments                                                                             |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|--------------------------------------------------------------------------------------|
| <b>SECTION I</b>                                                                                                                                                                                                                                                                                                         |          |                                                                                      |
| <b>1. B</b> - $8x^3 + 27 = (2x)^3 + 3^3$<br>$= (2x + 3)((2x)^2 - (2x)(3) + 3^2)$<br>$= (2x + 3)(4x^2 - 6x + 9)$                                                                                                                                                                                                          | <b>1</b> |                                                                                      |
| <b>2. D</b> – the only graph that satisfies the “vertical line test”                                                                                                                                                                                                                                                     | <b>1</b> |                                                                                      |
| <b>3. A</b> - $\frac{1+\sqrt{3}}{2-\sqrt{3}} \times \frac{2+\sqrt{3}}{2+\sqrt{3}} = \frac{2+3\sqrt{3}+3}{4-3}$<br>$= 5 + 3\sqrt{3}$                                                                                                                                                                                      | <b>1</b> |                                                                                      |
| <b>4. D</b> – from the diagram;<br>Angle of elevation is $\alpha$<br>Angle of depression is $\theta$                                                                                                                                                                                                                     | <b>1</b> |                                                                                      |
| <b>5. C</b> - $\cos \theta < 0$ and $\sin \theta < 0 \Rightarrow$ quadrant 3, $\therefore \tan \theta > 0$<br><br>$\tan \theta = \frac{4}{3}$                                                                                                                                                                            | <b>1</b> |    |
| <b>6. A</b> - $\frac{x+5}{(x-3)(x+1)} - \frac{x-1}{x^2-x-2} = \frac{x+5}{(x-3)(x+1)} - \frac{x-1}{(x-2)(x+1)}$<br>$= \frac{(x+5)(x-2) - (x-1)(x-3)}{(x-3)(x-2)(x+1)}$<br>$= \frac{x^2 + 3x - 10 - x^2 + 4x - 3}{(x-3)(x-2)(x+1)}$<br>$= \frac{7x - 13}{(x-3)(x-2)(x+1)}$                                                 | <b>1</b> |                                                                                      |
| <b>7. C</b> - $\frac{\sin(360^\circ - A)}{\sin(90^\circ - A)} = \frac{-\sin A}{\cos A}$<br>$= -\tan A$                                                                                                                                                                                                                   | <b>1</b> |                                                                                      |
| <b>8. B</b> - $f(a) = g(2a)$<br>$3a^2 - 4 = (2a)^2 - 2(2a)$<br>$3a^2 - 4 = 4a^2 - 4a$<br>$a^2 - 4a + 4 = 0$<br>$(a - 2)^2 = 0$<br>$a = 2 \Rightarrow$ only one solution                                                                                                                                                  | <b>1</b> |                                                                                      |
| <b>9. C</b> - $x^2 + y^2 - 12x - 10y + k = 0$<br>$(x - 6)^2 + (y - 5)^2 = 61 - k \Rightarrow$ centre is (6,5)<br><br>For exactly three intercepts the radius = 6<br><br>$61 - k = 36$<br>$k = 25$                                                                                                                        | <b>1</b> |  |
| <b>10. B</b> - $\frac{6^{r+s} \times 12^{r-s}}{8^r \times 9^{r+2s}} = \frac{2^{r+s} \times 3^{r+s} \times 2^{2r-2s} \times 3^{r-s}}{2^{3r} \times 3^{2r+4s}}$<br>$= \frac{2^{3r-s} \times 3^{2r}}{2^{3r} \times 3^{2r+4s}}$<br>$= 2^{-s} \times 3^{-4s}$<br><br>Thus $s \leq 0$ in order for expression to be an integer | <b>1</b> |                                                                                      |


**SECTION II**  
**QUESTION 11**

| Solution                                                                                                                                                                                                                                                                        | Marks | Comments                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>11(a)</b><br>$\sqrt[4]{x^5} = x^{\frac{5}{4}}$                                                                                                                                                                                                                               | 1     | <b>1 mark</b><br>• Correct answer                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>11 (b)</b><br>$\sqrt{\frac{4.81 \times 10^5}{7.36 \times 10^9}} = 0.00808413637...$<br>$= 0.0081 \text{ correct to two significant figures}$                                                                                                                                 | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Performs the correct calculation                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>11(c)</b><br>$\begin{aligned} f(-1) &= 7 - 2(-1)^2 \\ &= 7 - 2 \\ &= 5 \end{aligned}$                                                                                                                                                                                        | 1     | <b>1 mark</b><br>• Correct answer                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>11 (d)</b><br>$\begin{aligned} \sqrt{75} - 2\sqrt{27} &= 5\sqrt{3} - 6\sqrt{3} \\ &= -\sqrt{3} \end{aligned}$                                                                                                                                                                | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Simplifies at least one surd correctly                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>11 (e)</b><br>$\begin{aligned} (3x-4)(x-2)(x+2) \\ &= (3x-4)(x^2-4) \\ &= 3x^3 - 12x - 4x^2 + 16 \\ &= 3x^3 - 4x^2 - 12x + 16 \end{aligned}$                                                                                                                                 | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Performs a binomial product expansion                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>11 (f)</b><br>$\frac{2}{x-1} \leq 1$<br>$\begin{aligned} x-1 &\neq 0 \\ x &\neq 1 \end{aligned}$<br>$\begin{aligned} 2 &= x-1 \\ x &= 3 \end{aligned}$<br><br>$x < 1 \text{ or } x \geq 3$ | 3     | <b>3 marks</b><br>• Correct graphical solution on number line or algebraic solution, with correct working<br><b>2 marks</b><br>• Bald answer<br>• Identifies the two correct critical points via a correct method<br>• Correct conclusion to their critical points obtained using a correct method<br><b>1 mark</b><br>• Uses a correct method<br>• Acknowledges a problem with the denominator.<br><b>0 marks</b><br>• Solves like a normal equation, with no consideration of the denominator. |
| <b>11 (g) (i)</b><br>$\begin{aligned} 2x^2 + 3x - 2 \\ &= (2x-1)(x+2) \end{aligned}$                                                                                                                                                                                            | 1     | <b>1 mark</b><br>• Correct answer                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>11 (g) (ii)</b><br>$\begin{aligned} x^3 + 5x^2 + x + 5 \\ &= x^2(x+5) + 1(x+5) \\ &= (x+5)(x^2+1) \end{aligned}$                                                                                                                                                             | 1     | <b>1 mark</b><br>• Correct answer                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>11 (g) (iii)</b><br>$\begin{aligned} 4a^2(x^3 + 18ab^2) - (32a^5 + 9b^2x^3) \\ &= x^3(4a^2 - 9b^2) - 8a^3(4a^2 - 9b^2) \\ &= (4a^2 - 9b^2)(x^3 - 8a^3) \\ &= (2a-3b)(2a+3b)(x-2a)(x^2+2ax+4a^2) \end{aligned}$                                                               | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Correctly uses $a^2 - b^2$ or $a^3 - b^3$ factorisation                                                                                                                                                                                                                                                                                                                                                                               |
| QUESTION 12                                                                                                                                                                                                                                                                     |       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>12 (a) (i)</b><br>$\begin{aligned} \frac{125a^3 - 8}{a^2 - 7a + 10} \times \frac{a-5}{25a^2 - 4} \\ &= \frac{(5a-2)(25a^2 + 10a + 4)}{(a-5)(a-2)} \times \frac{a-5}{(5a-2)(5a+2)} \\ &= \frac{25a^2 + 10a + 4}{(a-2)(5a+2)} \end{aligned}$                                   | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Factorises 2 out of the 3 non-linear expressions                                                                                                                                                                                                                                                                                                                                                                                      |

**QUESTION 12...continued.**

| Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Marks    | Comments                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>12 (a) (ii)</b></p> $\frac{2}{x^2-1} - \frac{1}{x^2-x} + \frac{x-1}{x^2+x}$ $= \frac{2}{(x-1)(x+1)} - \frac{1}{x(x-1)} + \frac{x-1}{x(x+1)}$ $= \frac{2x - (x+1) + (x-1)^2}{x(x+1)(x-1)}$ $= \frac{2x - x - 1 + x^2 - 2x + 1}{x(x+1)(x-1)}$ $= \frac{x^2 - x}{x(x+1)(x-1)}$ $= \frac{x(x-1)}{x(x+1)(x-1)}$ $= \frac{1}{x+1}$                                                                                                                                                                                                                          | <b>3</b> | <p><b>3 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Rewrites as a single fraction</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Finds the LCD</li> </ul>                                                                                                                                                                                                                                                              |
| <p><b>12 (b) (i)</b></p> $\frac{3x-2}{4} - \frac{2x-1}{8} = 5$ $2(3x-2) - (2x-1) = 40$ $6x - 4 - 2x + 1 = 40$ $4x = 43$ $x = \frac{43}{4}$                                                                                                                                                                                                                                                                                                                                                                                                                  | <b>2</b> | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Removes the fractions by multiplying by the LCD, or equivalent</li> </ul>                                                                                                                                                                                                                                                                                                                     |
| <p><b>12 (b) (ii)</b></p> $ 2x+6  = 3x-1$ <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <math display="block">2x+6 = 3x-1</math> <math display="block">x = 7</math> </div> <div style="width: 45%; text-align: right;"> <math display="block">-(2x+6) = 3x-1</math> <math display="block">5x = -5</math> <math display="block">x = -1</math> </div> </div> <p align="center"><b>OR</b></p> <p align="center"><b>NOT A SOLUTION</b></p> <p align="center"><math>\therefore x = 7</math></p>                         | <b>3</b> | <p><b>3 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Finds two “possible” answers</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Finds an answer without considering cases.</li> </ul>                                                                                                                                                                                                                                  |
| <p><b>12 (b) (iii)</b></p> $2x^2 - 6x \leq 0$ $2x(x-3) \leq 0$ $0 \leq x \leq 3$                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | <b>2</b> | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Establishes the two critical points of the solution.</li> </ul>                                                                                                                                                                                                                                                                                                                               |
| <p><b>12(b) (iv)</b></p> $\frac{x+1}{x^2-4} \geq 0$ <div style="display: flex; justify-content: space-between; margin-top: 10px;"> <div style="width: 45%;"> <math display="block">x^2 - 4 \neq 0</math> <math display="block">x \neq \pm 2</math> </div> <div style="width: 45%; text-align: right;"> <math display="block">x+1 = 0</math> <math display="block">x = -1</math> </div> </div>  <p align="center"><math>-2 &lt; x \leq -1 \text{ or } x &gt; 2</math></p> | <b>3</b> | <p><b>3 marks</b></p> <ul style="list-style-type: none"> <li>• Correct graphical solution on number line or algebraic solution</li> </ul> <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Bald answer</li> <li>• Identifies the three correct critical points via a correct method</li> <li>• Correct conclusion to their critical points obtained using a correct method</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Uses a correct method</li> <li>• Acknowledges a problem with denominator.</li> </ul> |
| <b>QUESTION 13</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <p><b>13 (a)</b> Using the graph, the question becomes;<br/> <i>“when is the line below the absolute value graph?”</i><br/>         As we are only concerned with the x values, the solution is</p> $x \leq -1 \text{ or } x \geq 3$                                                                                                                                                                                                                                                                                                                        | <b>1</b> | <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| <p><b>13 (b)</b></p> $\frac{10}{\sqrt{5}-\sqrt{3}} \times \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}+\sqrt{3}} = \frac{10(\sqrt{5}+\sqrt{3})}{5-3}$ $= \frac{10(\sqrt{5}+\sqrt{3})}{2}$ $= 5(\sqrt{5}+\sqrt{3})$                                                                                                                                                                                                                                                                                                                                                     | <b>2</b> | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Attempts to multiply by the conjugate of the denominator</li> </ul>                                                                                                                                                                                                                                                                                                                           |

**QUESTION 13...continued.**

| Solution                            |                                                                                                                                                                                                                                                                                                                         | Marks | Comments                                                                                                                                                                                                                                                                                 |
|-------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 13 (c)                              | <p>let <math>x = 0.01232323...</math> <b>OR</b> <math>0.01\dot{2}\dot{3} = \frac{123-1}{9900}</math></p> $100x = 1.23232323...$ $99x = 1.22$ $x = \frac{1.22}{99}$ $= \frac{122}{9900}$ $= \frac{61}{4950}$ <p><math>\therefore 0.01\dot{2}\dot{3} = \frac{61}{4950}</math></p>                                         | 2     | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Evidence of a valid manual calculation</li> </ul>                                                                               |
| 13 (d)                              | $F = \frac{mv^2}{gr}$ $mv^2 = Fgr$ $v^2 = \frac{Fgr}{m}$ $v = \pm \sqrt{\frac{Fgr}{m}}$                                                                                                                                                                                                                                 | 2     | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• makes <math>v^2</math> the subject of the formula</li> </ul>                                                                    |
| 13 (e)                              | <p><math>x^2 - 8x + 9 = 0</math> <b>OR</b> <math>x^2 - 8x + 9 = 0</math></p> $x^2 - 8x = -9$ $x^2 - 8x + 4^2 = 7$ $(x-4)^2 = 7$ $x-4 = \pm\sqrt{7}$ $x = 4 \pm \sqrt{7}$ <p><math>(x-4)^2 - 7 = 0</math></p> $(x-4-\sqrt{7})(x-4+\sqrt{7}) = 0$ $x = 4 \pm \sqrt{7}$                                                    | 2     | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Completes the square to obtain <math>(x-4)^2</math></li> <li>• Solves the equation for a different completed square.</li> </ul> |
| 13 (f) (i) when $x = 2$ and $y = 1$ | $(x+2y)(2x-y) + (x-y)(3x+4y) = (2+2)(4-1) + (2-1)(6+4)$ $= 12 + 10$ $= 22$                                                                                                                                                                                                                                              | 1     | <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Correct answer</li> </ul>                                                                                                                                                                                                  |
| 13 (f) (ii) when $y = 1$            | $(x+2)(2x-1) + (x-1)(3x+4) = 22$ $2x^2 + 3x - 2 + 3x^2 + x - 4 = 22$ $5x^2 + 4x - 28 = 0$ $(x-2)(5x+14) = 0$ $x = 2 \quad \text{or} \quad x = -\frac{14}{5}$ <p><math>\therefore</math> the only other possible value of <math>x</math> is <math>-\frac{14}{5}</math></p>                                               | 2     | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Establishes quadratic in terms of <math>x</math></li> </ul>                                                                     |
| 13 (g) (i)                          | $(x-y+2)(x+y-1)$ $= x^2 + xy - x - xy - y^2 + y + 2x + 2y - 2$ $= x^2 - y^2 + x + 3y - 2$ $= f(x)$                                                                                                                                                                                                                      | 1     | <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul>                                                                                                                                                                                                |
| 13 (g) (ii)                         | $x^2 - y^2 + x + 3y > 2$ $x^2 - y^2 + x + 3y - 2 > 0$ $(x-y+2)(x+y-1) > 0$ <p><math>\therefore</math> either <math>x-y+2 &gt; 0</math> and <math>x+y-1 &gt; 0</math> OR <math>x-y+2 &lt; 0</math> and <math>x+y-1 &lt; 0</math></p>  | 2     | <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Some correct regions indicated, with no more than one incorrect region.</li> </ul>                                              |

**QUESTION 14**



| Solution                                                                                                                                                                                                                            | Marks | Comments                                                                                                                                                                                                                        |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>14 (a)</b> some possibilities would include; <ul style="list-style-type: none"> <li>A divides BP externally in the ratio 1:4</li> <li>B divides PA in the ratio 3:1</li> <li>P divides BA externally in the ratio 3:4</li> </ul> | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Two different statements</li> </ul>                                                                                                                                        |
| <b>14 (b) (i)</b> $f(x)$ is even if $f(-x) = f(x)$                                                                                                                                                                                  | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct condition</li> </ul>                                                                                                                                               |
| <b>14 (b) (ii)</b> some examples would include; <ul style="list-style-type: none"> <li><math>y = x^2</math></li> <li><math>y =  x </math></li> <li><math>y = \cos x</math></li> </ul>                                               | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct example</li> </ul>                                                                                                                                                 |
| <b>14 (c)</b> $f(2) + f(5) - f(-2) = 3(2) + (5^2 + 1) - 3(-2)$<br>$= 6 + 26 + 6$<br>$= 38$                                                                                                                                          | 2     | <b>2 marks</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul> <b>1 mark</b> <ul style="list-style-type: none"> <li>finds at least two function values</li> </ul>                                            |
| <b>14 (d) (i)</b> $x^2 - 16 \geq 0$<br>$x \leq -4$ or $x \geq 4$                                                                                                                                                                    | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct answer</li> </ul>                                                                                                                                                  |
| <b>14 (d) (ii)</b> $y \geq 0$                                                                                                                                                                                                       | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct answer</li> </ul>                                                                                                                                                  |
| <b>14 (e) (i)</b> $P(0,1)$                                                                                                                                                                                                          | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct answer</li> </ul>                                                                                                                                                  |
| <b>14 (e) (ii)</b> $x^2 + y^2 = 2^2 + 4^2$<br>$= 20$<br>$\therefore$ circle has equation $x^2 + y^2 = 20$                                                                                                                           | 2     | <b>2 marks</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul> <b>1 mark</b> <ul style="list-style-type: none"> <li>Recognises that a circle's equation is of the form <math>x^2 + y^2 = k</math></li> </ul> |
| <b>14 (e) (iii)</b> $(2,4) : 4 = a^2$<br>$a = 2$<br>$\therefore$ exponential has the equation $y = 2^x$                                                                                                                             | 1     | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct answer</li> </ul>                                                                                                                                                  |
| <b>14 (e) (iv)</b> $x \geq 0$<br>$x^2 + y^2 \leq 20$<br>$y \geq 2^x$                                                                                                                                                                | 2     | <b>2 marks</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul> <b>1 mark</b> <ul style="list-style-type: none"> <li>At least two correct inequations</li> </ul>                                              |
| <b>14 (f)</b> $f(-x) =  -x-2  -  -x+2 $<br>$=  -1  x+2  -  -1  x-2 $<br>$=  x+2  -  x-2 $<br>$= -f(x)$<br>$\therefore$ function is odd                                                                                              | 2     | <b>2 marks</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul> <b>1 mark</b> <ul style="list-style-type: none"> <li>Recognises the condition for an odd function</li> </ul>                                  |

**QUESTION 15**





|                                                                                                                                |   |                                                                                                                                                                                              |
|--------------------------------------------------------------------------------------------------------------------------------|---|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>15 (a) (i)</b> $\cos 75^\circ = \sin(90^\circ - 75^\circ)$<br>$= \sin 15^\circ$<br>$= \frac{\sqrt{3} - 1}{2\sqrt{2}}$       | 1 | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul>                                                                                                             |
| <b>15 (a) (ii)</b> $\operatorname{cosec} 15^\circ = \frac{1}{\sin 15^\circ}$<br>$= \frac{2\sqrt{2}}{\sqrt{3} - 1}$             | 1 | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul>                                                                                                             |
| <b>15 (a) (iii)</b> $\sin 195^\circ = \sin(180^\circ + 15^\circ)$<br>$= -\sin 15^\circ$<br>$= -\frac{1 - \sqrt{3}}{2\sqrt{2}}$ | 1 | <b>1 mark</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul>                                                                                                             |
| <b>15 (b)</b> $2x + y = 4 \Rightarrow 4x + 2y = 8$<br>$5x + 2y = 9$<br>$\underline{5x + 2y = 9}$<br>$x = 1 \therefore y = 2$   | 2 | <b>2 marks</b> <ul style="list-style-type: none"> <li>Correct solution</li> </ul> <b>1 mark</b> <ul style="list-style-type: none"> <li>Finds the correct value for one pronumeral</li> </ul> |

| QUESTION 15...continued.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |       |                                                                                                                                                                                                                                                                                    |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Marks | Comments                                                                                                                                                                                                                                                                           |
| <b>15 (c)</b> $\sin^4 x - \cos^4 x = (\sin^2 x - \cos^2 x)(\sin^2 x + \cos^2 x)$<br>$= \sin^2 x - \cos^2 x$<br>$= (1 - \cos^2 x) - \cos^2 x$<br>$= 1 - 2\cos^2 x$                                                                                                                                                                                                                                                                                                                                                                    | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Uses a valid trig identity in a relevant manner.                                                                                                                                                                        |
| <b>15 (d) (i)</b> The student divided out a possible solution when canceling $\sin x$ .<br>When solving equations you can only divide by an unknown if there is no possibility that the unknown could equal zero.                                                                                                                                                                                                                                                                                                                    | 1     | <b>1 mark</b><br>• Correct explanation                                                                                                                                                                                                                                             |
| <b>15 (d) (ii)</b> The missing answer comes from the possibility that;<br>$\sin x = 0$<br>$x = 0^\circ, 180^\circ, 360^\circ$<br>So the missing answer is $x = 180^\circ$ .                                                                                                                                                                                                                                                                                                                                                          | 1     | <b>1 mark</b><br>• Correct solution                                                                                                                                                                                                                                                |
| <b>15 (e) (i)</b> $\sin \theta \tan \theta + 2\sin \theta = 3\cos \theta$<br>$\frac{\sin^2 \theta}{\cos \theta} + 2\sin \theta = 3\cos \theta$<br>$\sin^2 \theta + 2\sin \theta \cos \theta = 3\cos^2 \theta$<br>$3\cos^2 \theta - 2\sin \theta \cos \theta - \sin^2 \theta = 0$<br>$(3\cos \theta + \sin \theta)(\cos \theta - \sin \theta) = 0$<br>$\tan \theta = -3$ OR $\tan \theta = 1$<br>$\theta = 108^\circ, 288^\circ$ OR $\theta = 45^\circ, 225^\circ$<br>$\therefore \theta = 45^\circ, 108^\circ, 225^\circ, 288^\circ$ | 3     | <b>3 marks</b><br>• Correct solution<br><b>2 marks</b><br>• Finds the two possibilities for $\tan \theta$<br><b>1 mark</b><br>• Correctly manipulates terms into a quadratic equation                                                                                              |
| <b>15 (e) (ii)</b> $\sin(20^\circ - 2\theta) = \frac{1}{7}$<br>$\sin \alpha = \frac{1}{7}$<br>$\alpha = 8^\circ$<br>$(20^\circ - 2\theta) = 8^\circ, 172^\circ, 368^\circ, 532^\circ$<br>$-2\theta = -12^\circ, 152^\circ, 348^\circ, 512^\circ$<br>$\theta = 6^\circ, -76^\circ, -174^\circ, -256^\circ$<br>However solutions need to be $0^\circ \leq \theta \leq 360^\circ$ , so add $\pm 360^\circ$ to any answers not in required range<br>$\therefore \theta = 6^\circ, 104^\circ, 186^\circ, 284^\circ$                       | 3     | <b>3 marks</b><br>• Correct solution<br><b>2 marks</b><br>• Finds four consecutive answers for $(20^\circ - 2\theta)$<br>• Finds two answers for $\theta$<br><b>1 mark</b><br>• Finds four consecutive answers for $(20^\circ - 2\theta)$<br>• Finds two answers for $\pm 2\theta$ |
| QUESTION 16                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |       |                                                                                                                                                                                                                                                                                    |
| <b>16(a) (i)</b> $\sin \angle \text{TPS} = \frac{2.5}{4}$<br>$\angle \text{TPS} = 36.6821875\dots$<br>$= 36^\circ$ to nearest degree                                                                                                                                                                                                                                                                                                                                                                                                 | 1     | <b>1 mark</b><br>• Correct answer<br><i>Note: no rounding penalty</i>                                                                                                                                                                                                              |
| <b>16 (a) (ii)</b> $\left(\frac{1}{2}\text{PS}\right)^2 = 4^2 - 2.5^2$<br>$\frac{1}{2}\text{PS} = 3.1224998999\dots$<br>$\text{PS} = 6.244997998\dots$<br>$= 6.24$ metres to the nearest centimetre                                                                                                                                                                                                                                                                                                                                  | 1     | <b>1 mark</b><br>• Correct answer<br><i>Note: no rounding penalty</i>                                                                                                                                                                                                              |
| <b>16 (a) (iii)</b> $\frac{2.5}{\text{TQ}} = \sin 65^\circ$<br>$\text{TQ} = \frac{2.5}{\sin 65^\circ}$<br>$= 2.758444797\dots$<br>$= 276$ metres to the nearest centimetre                                                                                                                                                                                                                                                                                                                                                           | 1     | <b>1 mark</b><br>• Correct answer<br><i>Note: no rounding penalty</i>                                                                                                                                                                                                              |
| <b>16 (b) (i)</b> $\text{Area} = \frac{1}{2} \times 40 \times 80 \times \sin 130^\circ$<br>$= 1225.671109\dots$<br>$= 1226 \text{ m}^2$                                                                                                                                                                                                                                                                                                                                                                                              | 1     | <b>1 mark</b><br>• Correct answer<br><i>Note: no rounding penalty</i>                                                                                                                                                                                                              |
| <b>16 (b) (ii)</b> The largest side is always opposite the largest angle, and since the angle is obtuse, it must be the largest angle.                                                                                                                                                                                                                                                                                                                                                                                               | 1     | <b>1 mark</b><br>• Correct explanation                                                                                                                                                                                                                                             |
| <b>16 (b) (iii)</b> $x^2 = 40^2 + 80^2 - 2 \times 40 \times 80 \times \cos 130^\circ$<br>$x = 110.0628943\dots$<br>$x = 110$ metres to the nearest metre                                                                                                                                                                                                                                                                                                                                                                             | 1     | <b>1 mark</b><br>• Correct answer<br><i>Note: no rounding penalty</i>                                                                                                                                                                                                              |

**QUESTION 16...continued.**

| Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Marks | Comments                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>16 (c) (i)</b></p>                                                                                                                                                                                                                                                                                                                                                                                                | 1     | <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Correct diagram with <b>all</b> information labelled</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <p><b>16 (c) (ii)</b></p> $\angle GBW = 302^\circ - 270^\circ = 32^\circ$ $\angle GAB = 296^\circ - 270^\circ = 26^\circ$ $\angle GBW = \angle AGB + \angle GAB \quad (\text{exterior } \angle, \triangle BAG)$ $32^\circ = \angle AGB + 26^\circ$ $\angle AGB = 6^\circ$                                                                                                                                                                                                                                | 1     | <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Correct explanation</li> </ul> <p><i>Note: formal geometric explanation not required</i></p>                                                                                                                                                                                                                                                                                                                                                                                                             |
| <p><b>16 (c) (iii)</b></p> $\frac{BG}{\sin \angle GAB} = \frac{AB}{\sin \angle AGB}$ $\frac{BG}{\sin 26^\circ} = \frac{1500}{\sin 6^\circ}$ $BG = \frac{1500 \sin 26^\circ}{\sin 6^\circ}$                                                                                                                                                                                                                                                                                                               | 1     | <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Evidence of using sine rule in finding the correct expression, or similar merit.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                              |
| <p><b>16 (c) (iv)</b></p>  $RG^2 = BG^2 + BR^2 - 2 \times BG \times BR \times \cos \angle RBG$ $BR^2 - (2 \times BG \times \cos \angle RBG) BR + (BG^2 - RG^2) = 0$ $BR = \frac{2 \times BG \times \cos 32^\circ \pm \sqrt{(2 \times BG \times \cos 32^\circ)^2 - 4(BG^2 - RG^2)}}{2}$ $= 1608.235614... \text{ or } 9061.388854$ <p><math>\therefore</math> the soldier can travel 1608 metres</p>                    | 3     | <p><b>3 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Establishes a quadratic equation in terms of BR or similar merit</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Attempts to find RG using the cosine rule or similar merit.</li> </ul> <p><i>Notes:</i></p> <ul style="list-style-type: none"> <li>• no rounding penalty</li> <li>• OK for approximate BG value (6290.695373...) to be used in working and calculations</li> </ul> |
| <p><b>16 (d)</b> In <math>\triangle OPQ</math>;</p> $\cos \theta = \frac{OP^2 + OQ^2 - PQ^2}{2 \times OP \times OQ}$ $= \frac{\left(\frac{R}{2}\right)^2 + (R-r)^2 - \left(\frac{R}{2} + r\right)^2}{2 \left(\frac{R}{2}\right)(R-r)}$ $= \frac{\frac{R^2}{4} + R^2 - 2rR + r^2 - \frac{R^2}{4} - rR - r^2}{R(R-r)}$ $= \frac{R^2 - 3rR}{R(R-r)}$ $= \frac{R - 3r}{R - r}$ $(R-r)\cos \theta = R - 3r$ $r(3 - \cos \theta) = R(1 - \cos \theta)$ $\frac{r}{R} = \frac{1 - \cos \theta}{3 - \cos \theta}$ | 3     | <p><b>3 marks</b></p> <ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <p><b>2 marks</b></p> <ul style="list-style-type: none"> <li>• Establishes <math>\cos \theta = \frac{R - 3r}{R - r}</math>, or equivalent</li> </ul> <p><b>1 mark</b></p> <ul style="list-style-type: none"> <li>• Finds a relationship between <math>\theta</math>, <math>R</math> and <math>r</math>.</li> </ul>                                                                                                                                                          |

**QUESTION 17**

| Solution    |                                                                                                                                                                                                                                                                              | Marks | Comments                                                                                                                                                           |
|-------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 17 (a)      |  $P\left(\frac{6 - 6 \cdot \frac{2}{5}, 12 - \frac{22}{5}}{5}\right)$ $= (0, -2)$                                                                                                           | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Finds correct x or y value                                                                              |
| 17 (b) (i)  | $PX^2 = PR^2 - RX^2$ $= 6^2 - 3^2$ $= 27$ $PX = 3\sqrt{3}$ $\tan \alpha = \frac{3\sqrt{3}}{8}$ $\alpha = 33.0044916\dots$ $\alpha = 33^\circ$                                               | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Calculates PX<br>• Correctly identifies the angle to be found<br><i>Note: no rounding penalty</i>       |
| 17 (b) (ii) | $PC^2 = PR^2 + CR^2$ $= 6^2 + 8^2$ $= 100$ $PC = 10$ $\sin \beta = \frac{3\sqrt{3}}{10}$ $\beta = 31.30644625\dots$ $\beta = 31^\circ$                                                      | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Calculates PC or CX<br>• Correctly identifies the angle to be found<br><i>Note: no rounding penalty</i> |
| 17 (c)      | $y = \frac{x^2}{x^2 + 5x + 6}$ $= \frac{x^2 + 5x + 6}{x^2 + 5x + 6} - \frac{5x + 6}{x^2 + 5x + 6}$ $= 1 - \frac{5x + 6}{(x + 3)(x + 2)}$ <p>Thus;<br/>vertical asymptotes are <math>x = -3</math> and <math>x = -2</math><br/>horizontal asymptote is <math>y = 1</math></p> | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Finds both vertical asymptotes<br>• Finds the horizontal asymptote                                      |
| 17 (d)      | $3^{2018} - 2^{2018} = (3^2)^{1009} - (2^2)^{1009}$ $= (3^2 - 2^2)(3^{2016} + 3^{2014} \times 2^2 + 3^{2012} \times 2^4 + \dots + 2^{2016})$ <p>As neither factor is equal to 1, then <math>3^{2018} - 2^{2018}</math> is <b>not</b> prime</p>                               | 1     | <b>1 mark</b><br>• Correct explanation                                                                                                                             |
| 17 (e) (i)  |  $x = \frac{1 - k}{k + 1}$ $y = \frac{2 - 4k}{k + 1}$                                                                                                                                     | 1     | <b>1 mark</b><br>• Correct answer                                                                                                                                  |
| 17 (e) (ii) | $3\left(\frac{1 - k}{k + 1}\right) - 4\left(\frac{2 - 4k}{k + 1}\right) - 5 = 0$ $3(1 - k) - 4(2 - 4k) - 5(k + 1) = 0$ $3 - 3k - 8 + 16k - 5k - 5 = 0$ $8k - 10 = 0$ $k = \frac{10}{8}$ $k = \frac{5}{4}$                                                                    | 2     | <b>2 marks</b><br>• Correct solution<br><b>1 mark</b><br>• Substitutes the found coordinates of K into the equation of the line.                                   |

| QUESTION 17...continued.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Marks | Comments                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>17 (f)</b><br>$\frac{AC}{h} = \tan 56^\circ$<br>$AC = h \tan 56^\circ$<br>$\angle ACB = 72^\circ$ (alternate $\angle$ 's = $\parallel$ lines)<br>$\angle ABC + 45^\circ + \angle ACB = 180^\circ$ ( $\angle$ sum of $\Delta$ )<br>$\angle ABC + 45^\circ + 72^\circ = 180^\circ$<br>$\angle ABC = 63^\circ$<br>$\frac{AC}{\sin 63^\circ} = \frac{1}{\sin 72^\circ}$<br>$h \tan 56^\circ = \frac{\sin 63^\circ}{\sin 72^\circ}$<br>$h = \frac{\sin 63^\circ}{\sin 72^\circ \tan 56^\circ}$<br>$h = 0.63199198479...$<br>$h = 632 \text{ m}$ , correct to the nearest metre | 3     | <b>3 marks</b><br><ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <b>2 marks</b><br><ul style="list-style-type: none"> <li>• Finds sufficient sides and angles in <math>\Delta ABC</math> in order to solve the problem</li> </ul> <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Finds an expression for AC in terms of <math>h</math>.</li> <li>• Finds the missing two angles in <math>\Delta ABC</math></li> </ul> <p><i>Note: full marks awarded for the "exact value" for <math>h</math></i></p> |
| QUESTION 18                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>18 (a) (i)</b> Ways = $8!$<br>$= 40320$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 1     | <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Correct answer</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>18(a) (ii)</b> Ways = $2! \times (3 \times 2) \times 6!$<br>$= 8640$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 2     | <b>2 marks</b><br><ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Handles restriction in a logical manner</li> </ul>                                                                                                                                                                                                                                                                                                                                    |
| <b>18 (b) (i)</b> Possibilities = ${}^{22}C_5$<br>$= 26334$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | 1     | <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Correct answer</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>18(b) (ii)</b> Committees = ${}^{22}C_5 - {}^{13}C_5 - {}^9C_5$<br>$= 24921$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | 2     | <b>2 marks</b><br><ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Handles restriction in a logical manner</li> </ul>                                                                                                                                                                                                                                                                                                                                    |
| <b>18 (b) (iii)</b> Ways = $4!$<br>$= 24$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | 1     | <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Correct answer</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>18 (c) (i)</b> Arrangements = $\frac{10!}{3!2!}$<br>$= 302400$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | 1     | <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Correct answer</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>18 (c) (ii)</b> Arrangements = $\frac{7!}{2!} \times {}^8C_3$<br>$= 141120$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | 2     | <b>2 marks</b><br><ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Approaches the restriction in a logical manner</li> </ul>                                                                                                                                                                                                                                                                                                                             |
| <b>18 (d)</b> # triangles = ${}^{10}C_3 - {}^4C_3$<br>$= 116$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | 2     | <b>2 marks</b><br><ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Selects three points from ten, without considering the implication of the collinear points</li> </ul>                                                                                                                                                                                                                                                                                 |
| <b>18 (e)</b> Case 1: 3 red plus 1 other colour = $2 \times \frac{4!}{3!}$<br>$= 8$<br>Case 2: 2 red plus 2 blue = $\frac{4!}{2!2!}$<br>$= 6$<br>Case 3: 2 of one colour plus 2 other colours = $2 \times \frac{4!}{2!}$<br>$= 24$<br>Total # signals = $8 + 6 + 24$<br>$= 38$                                                                                                                                                                                                                                                                                               | 3     | <b>3 marks</b><br><ul style="list-style-type: none"> <li>• Correct solution</li> </ul> <b>2 marks</b><br><ul style="list-style-type: none"> <li>• Finds the correct number of ways in their different cases</li> </ul> <b>1 mark</b><br><ul style="list-style-type: none"> <li>• Considers different cases in an attempt to solve the problem</li> </ul>                                                                                                                                                                               |