



Student Number	
Mark / 24	

Chemistry

**HSC Course
Production of Materials
Theory Test • 2002**

General Instructions

- Reading time – 5 minutes
- Working time – 40 minutes
- Write using black or blue pen
- Draw diagrams using pencil
- Board-approved calculators may be used
- A data sheet and a Periodic Table are provided at the back of this paper
- Write your Student Number at the top of this page

Assessment Weighting – 4%

Total Marks – 24

Part A – 4 marks

- Attempt Questions 1 – 4
- Allow about 5 minutes for this part

Part B – 20 marks

- Attempt Questions 5 – 10
- Allow about 35 minutes for this part

Part A – 4 marks
Attempt Questions 1 – 4
Allow about 5 minutes for this part

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample: $2 + 4 =$ (A) 2 (B) 6 (C) 8 (D) 9
 A ☐ B ☒ C ☐ D ☐

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.


A ☒ B ☒ C ☐ D ☐

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word **correct** and drawing an arrow as follows.

A ☒ B ☒ C ☐ D ☐
 correct ↗

Answer Box for Questions 1 – 4

1	A ☐	B ☐	C ☐	D ☐
2	A ☐	B ☐	C ☐	D ☐
3	A ☐	B ☐	C ☐	D ☐
4	A ☐	B ☐	C ☐	D ☐



Mark your answers for Questions 1 – 4 in the Answer Box on page 1.

- 1 Which of the following lists contains only condensation polymers?
- (A) cellulose, protein, starch
 - (B) cellulose, polyvinyl chloride, polyethylene
 - (C) polystyrene, starch, protein
 - (D) polyvinyl chloride, polyethylene, polystyrene
- 2 Which of the following defines the term *cracking* used in the petrochemical industry?
- (A) addition of hydrogen to a compound
 - (B) preparation of a polymer from a hydrocarbon monomer
 - (C) formation of saturated hydrocarbons from alkanes
 - (D) conversion of long chain hydrocarbons to shorter chain molecules
- 3 A mixture of ethanol and ethylene is heated with concentrated sulfuric acid in a closed container and a reaction occurs. What is the likely outcome?
- (A) more ethylene forms
 - (B) more ethanol forms
 - (C) CO₂ and H₂O form
 - (D) butane forms
- 4 In which of the following equations is the species printed in **bold** type being reduced?
- (A) $3\text{Zn}^{2+} + 2\textbf{Al}_{(\text{s})} \rightarrow 3\text{Zn}_{(\text{s})} + 2\text{Al}^{3+}$
 - (B) $2\text{Br}^{-} + \textbf{Cl}_{2(\text{g})} \rightarrow \text{Br}_{2(\text{l})} + 2\text{Cl}^{-}$
 - (C) $2\text{H}^{+} + \textbf{Mg}_{(\text{s})} \rightarrow \text{Mg}^{2+} + \text{H}_{2(\text{g})}$
 - (D) $2\text{H}_2\text{O}_{(\text{l})} + 3\text{I}_2 + 2\textbf{S}_2\text{O}_3^{2-} \rightarrow \text{S}_4\text{O}_8^{2-} + 4\text{H}^{+} + 6\text{I}^{-}$

Part B – 20 marks**Attempt Questions 5 – 10****Allow about 35 minutes for this part****Show all relevant working in questions involving calculations.**

Question 5 (4 marks)

Three groups of students set out to determine the heat of combustion of the three alkanols... methanol, CH_3OH ; ethanol, $\text{C}_2\text{H}_5\text{OH}$; and 1-propanol, $\text{C}_3\text{H}_7\text{OH}$.

Each group measured out 100 mL of water into a container and heated the water by burning a measured mass of alcohol. Their results are shown below...


Alcohol burned	Mass of H_2O heated (g)	Temperature rise ($^{\circ}\text{C}$)	Mass of alcohol burned (g)	Heat of Combustion (kJ mol^{-1})
methanol	100	10	0.185	725
ethanol	100	10	0.142	
1-propanol	100	10	0.125	2016

- (a) Given that 4.18 J are required to raise the temperature of 1.00 g of water by 1.00°C , use the above data to determine the following values...

- (i) Heat of combustion of ethanol in kJ g^{-1} **(1 mark)**


- (ii) Heat of combustion of ethanol in kJ mol^{-1} **(1 mark)**

Question 5 continues on page 4



Question 5 (continued)

- (b) Plot the heat of combustion (kJ mol^{-1}) against molar mass for all three alkanols. Clearly label the axes. **(1 mark)**




- (c) Use the graph to predict the heat of combustion of 1-butanol, $\text{C}_4\text{H}_9\text{OH}$ in kJ mol^{-1}

Question 6 (3 marks)

- (a) Give a balanced equation for the conversion of ethylene to ethanol. **(1 mark)**

Question 6 continues on page 5




Question 6 (continued)

- (b) Account for ethanol's extensive use as a solvent for polar and non-polar substances. Use a diagram to explain your answer. **(2 marks)**

Question 7 (3 marks)

An electrochemical cell was constructed using two half-cells. One half-cell consisted of tin metal and a tin(II) chloride solution and the other half-cell consisted of zinc metal and zinc chloride solution.


- Draw a diagram of the galvanic cell.
- Label the anode and the cathode.
- Indicate the direction of electron flow.



Question 8 (5 marks)

- (a) Explain the term *biopolymer* and identify an example. **(2 marks)**

- (b) Cellulose is a polymer of β -glucose. A β -glucose molecule is shown below....



Draw a segment of a cellulose molecule by joining three glucose molecules together. **(3 marks)**



Question 9 (2 marks)

A student was asked to perform a first-hand investigation to compare the reactivities of hexane and hexene by observing their reactions with bromine water.

- (a) Describe the reaction(s) observed by the student when the procedures were carried out in a darkened laboratory. **(1 mark)**


- (b) Write an equation to show any addition reaction(s) that occurred. **(1 mark)**

Question 10 (3 marks)

Alkenes and their derivatives are important substances in the production of polymers. Polyvinyl chloride (PVC) is one such polymer.

- (a) Draw the structure of polyvinyl chloride showing three linked monomer units. **(1 mark)**

- (b) Describe **one** use of polyvinyl chloride and a property which makes it useful for this purpose. **(2 marks)**



DATA SHEET

Avogadro's constant, N_A	$6.022 \times 10^{23} \text{ mol}^{-1}$
Volume of 1 mole ideal gas: at 101.3 kPa (1.00 atm) and	
at 273 K (0°C)	22.41 L
at 298 K (25°C)	24.47 L
Ionisation constant for water at 298 K (25°C), K_w	1.0×10^{-14}
Specific heat capacity of water	$4.18 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

Some useful formulae

$$\text{pH} = -\log_{10} [\text{H}^+]$$

$$\Delta H = -m C \Delta T$$

Some standard potentials

$\text{K}^+ + \text{e}^-$	$\rightleftharpoons$	K(s)	-2.94 V
$\text{Ba}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ba(s)	-2.91 V
$\text{Ca}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ca(s)	-2.87 V
$\text{Na}^+ + \text{e}^-$	$\rightleftharpoons$	Na(s)	-2.71 V
$\text{Mg}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Mg(s)	-2.36 V
$\text{Al}^{3+} + 3\text{e}^-$	$\rightleftharpoons$	Al(s)	-1.68 V
$\text{Mn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Mn(s)	-1.18 V
$\text{H}_2\text{O} + \text{e}^-$	$\rightleftharpoons$	$\frac{1}{2}\text{H}_2(\text{g}) + \text{OH}^-$	-0.83 V
$\text{Zn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Zn(s)	-0.76 V
$\text{Fe}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Fe(s)	-0.44 V
$\text{Ni}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ni(s)	-0.24 V
$\text{Sn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Sn(s)	-0.14 V
$\text{Pb}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Pb(s)	-0.13 V
$\text{H}^+ + \text{e}^-$	$\rightleftharpoons$	$\frac{1}{2}\text{H}_2(\text{g})$	0.00 V
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$\text{SO}_2(\text{aq}) + 2\text{H}_2\text{O}$	0.16 V
$\text{Cu}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Cu(s)	0.34 V
$\frac{1}{2}\text{O}_2(\text{g}) + \text{H}_2\text{O} + 2\text{e}^-$	$\rightleftharpoons$	2OH^-	0.40 V
$\text{Cu}^+ + \text{e}^-$	$\rightleftharpoons$	Cu(s)	0.52 V
$\frac{1}{2}\text{I}_2(\text{s}) + \text{e}^-$	$\rightleftharpoons$	I^-	0.54 V
$\frac{1}{2}\text{I}_2(\text{aq}) + \text{e}^-$	$\rightleftharpoons$	I^-	0.62 V
$\text{Fe}^{3+} + \text{e}^-$	$\rightleftharpoons$	Fe^{2+}	0.77 V
$\text{Ag}^+ + \text{e}^-$	$\rightleftharpoons$	Ag(s)	0.80 V
$\frac{1}{2}\text{Br}_2(\text{l}) + \text{e}^-$	$\rightleftharpoons$	Br^-	1.08 V
$\frac{1}{2}\text{Br}_2(\text{aq}) + \text{e}^-$	$\rightleftharpoons$	Br^-	1.10 V
$\frac{1}{2}\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	H_2O	1.23 V
$\frac{1}{2}\text{Cl}_2(\text{g}) + \text{e}^-$	$\rightleftharpoons$	Cl^-	1.36 V
$\frac{1}{2}\text{Cr}_2\text{O}_7^{2-} + 7\text{H}^+ + 3\text{e}^-$	$\rightleftharpoons$	$\text{Cr}^{3+} + \frac{7}{2}\text{H}_2\text{O}$	1.36 V
$\frac{1}{2}\text{Cl}_2(\text{aq}) + \text{e}^-$	$\rightleftharpoons$	Cl^-	1.40 V
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	$\rightleftharpoons$	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	1.51 V
$\frac{1}{2}\text{F}_2(\text{g}) + \text{e}^-$	$\rightleftharpoons$	F^-	2.89 V

Aylward and Findlay, *SI Chemical Data* (4th Edition) is the principal source of data for this examination paper. Some data may have been modified for examination purposes.

PERIODIC TABLE OF THE ELEMENTS

PERIODIC TABLE OF THE ELEMENTS											
KEY											
		Atomic Number		Atomic Weight		Symbol of element					
		79		Au		197.0		Gold			
						Name of element					
1 H 1.008 Hydrogen		4 Be 9.012 Beryllium		5 B 10.81 Boron		6 C 12.01 Carbon		7 N 14.01 Nitrogen		8 O 16.00 Oxygen	
3 Li 6.941 Lithium		12 Mg 24.31 Magnesium		13 Al 26.98 Aluminium		14 Si 28.09 Silicon		15 P 30.97 Phosphorus		16 S 32.07 Sulfur	
11 Na 22.99 Sodium		20 Ca 40.08 Calcium		21 Sc 44.96 Scandium		22 Ti 47.87 Titanium		23 V 50.94 Vanadium		24 Cr 52.00 Chromium	
19 K 39.10 Potassium		38 Sr 87.62 Strontium		39 Y 88.91 Yttrium		40 Zr 91.22 Zirconium		41 Nb 92.91 Niobium		42 Mo 95.94 Molybdenum	
37 Rb 85.47 Rubidium		72 Hf 178.5 Hafnium		73 Ta 180.9 Tantalum		74 W 183.8 Tungsten		75 Re 186.2 Rhenium		76 Os 190.2 Osmium	
55 Cs 132.9 Caesium		88 Ra [226.0] Radium		89-103 Actinides		104 Rf [261.1] Rutherfordium		105 Db [262.1] Dubnium		106 Sg [263.1] Seaborgium	
[223.0] Francium		[226.0] Radium		[223.0] Actinides		[261.1] Rutherfordium		[262.1] Dubnium		[263.1] Seaborgium	



Marking Scheme and Outcomes

Chemistry

HSC Course
Production of Materials
Theory Test • 2002

Assessment Weighting – 4%

Total Marks – 24

Working Time – 40 minutes

		OUTCOMES														
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
QUESTION	1									•						
	2									•						
	3								•	•						
	4							•						•		
	5							•			•			•		
	6									•						
	7								•	•						
	8									•						
	9								•	•				•		
	10							•	•					•		

Part A – Answer Box for Questions 1 – 4

1	A	☒	B	☐	C	☐	D	☐
2	A	☐	B	☐	C	☐	D	☒
3	A	☒	B	☐	C	☐	D	☐
4	A	☐	B	☒	C	☐	D	☐

- 1 Which of the following lists contains only condensation polymers?
- (A) cellulose, protein, starch
(B) cellulose, polyvinyl chloride, polyethylene
(C) polystyrene, starch, protein
(D) polyvinyl chloride, polyethylene, polystyrene
- 2 Which of the following defines the term *cracking* used in the petrochemical industry?
- (A) addition of hydrogen to a compound
(B) preparation of a polymer from a hydrocarbon monomer
(C) formation of saturated hydrocarbons from alkanes
(D) conversion of long chain hydrocarbons to shorter chain molecules
- 3 A mixture of ethanol and ethylene is heated with concentrated sulfuric acid in a closed container and a reaction occurs. What is the likely outcome?
- (A) more ethylene forms
(B) more ethanol forms
(C) CO₂ and H₂O form
(D) butane forms
- 4 In which of the following equations is the species printed in **bold** type being reduced?
- (A) $3\text{Zn}^{2+} + 2\textbf{Al}_{(s)} \rightarrow 3\text{Zn}_{(s)} + 2\text{Al}^{3+}$
(B) $2\text{Br}^{-} + \textbf{Cl}_{2(g)} \rightarrow \text{Br}_{2(l)} + 2\text{Cl}^{-}$
(C) $2\text{H}^{+} + \textbf{Mg}_{(s)} \rightarrow \text{Mg}^{2+} + \text{H}_{2(g)}$
(D) $2\text{H}_2\text{O}_{(l)} + 3\text{I}_2 + 2\textbf{S}_2\text{O}_3^{2-} \rightarrow \text{S}_4\text{O}_8^{2-} + 4\text{H}^{+} + 6\text{I}^{-}$

Question 5 (4 marks)

Three groups of students set out to determine the heat of combustion of the three alkanols... methanol, CH₃OH; ethanol, C₂H₅OH; and 1-propanol, C₃H₇OH.

Each group measured out 100 mL of water into a container and heated the water by burning a measured mass of alcohol. Their results are shown below...

Alcohol burned	Mass of H ₂ O heated (g)	Temperature rise (°C)	Mass of alcohol burned (g)	Heat of Combustion (kJ mol ⁻¹)
methanol	100	10	0.185	725
ethanol	100	10	0.142	
1-propanol	100	10	0.125	2016

- (a) Given that 4.18 J are required to raise the temperature of 1.00 g of water by 1.00 °C, use the above data to determine the following values...

- (i) Heat of combustion of ethanol in kJ g⁻¹ **(1 mark)**

$$\text{Heat of combustion (kJ/g)} = -m C \Delta T / \text{mass ethanol}$$

$$\text{kJ/g} = - (100\text{g} \times 4.18 \text{ J/g/}^\circ\text{C} \times 10^\circ\text{C}) / 0.142 = \underline{\underline{29.4 \text{ kJ/g}}}$$

- (ii) Heat of combustion of ethanol in kJ mol⁻¹ **(1 mark)**

$$\text{Heat of combustion} = -m C \Delta T / n$$

$$n = \# \text{ moles of ethanol} = (0.142\text{g} / 46.068 \text{ g/mol}) = 3.08 \times 10^{-3} \text{ mol}$$


$$\text{kJ/mol} = - (100\text{g} \times 4.18 \text{ J/g/}^\circ\text{C} \times 10^\circ\text{C}) / 3.08 \times 10^{-3} \text{ mol} = \underline{\underline{1360 \text{ kJ/mol}}}$$

☞ Heat of combustion is defined as the heat evolved, ∴ in this linguistic context a negative sign is not required, but will be accepted.

Question 5 continues on page 4

Question 5 (continued)

- (b) Plot the heat of combustion (kJ mol^{-1}) against molar mass for all three alkanols. Clearly label the axes. **(1 mark)**



- (c) Use the graph to predict the heat of combustion of 1-butanol, $\text{C}_4\text{H}_9\text{OH}$ in kJ mol^{-1}

Extrapolated value is about **2600 kJ/mol**

Criteria	Marks
Correct calculation of heat of combustion for the units specified	2
Graph drawn with the proper values on y - axis	1
Correct extrapolated value	1

Question 6 (5 marks)

- (a) Give a balanced equation for the conversion of ethylene to ethanol including reaction conditions. **(1 mark)**



Criterion	Mark
Correct equation including heating with catalyst of conc. H_2SO_4 or H_3PO_4	1


Question 6 continues on page 5

Question 6 (continued)

- (b) Account for ethanol's extensive use as a solvent for polar and non-polar substances. Use a diagram to explain your answer. **(2 marks)**

Criterion	Marks
The answer should include a depiction of the molecular structure of ethanol and as a cause of the resulting charge polarization within the molecule to explain its ability to dissolve various types of substances.	2

Ethanol has both polar (the – OH) and non-polar, (the C₂H₅ –) parts. The polarity of the hydroxy group is due to the much greater electronegativity of oxygen compared with carbon and hydrogen. The non-polar ethyl group enables ethanol to dissolve non-polar substances such as hexane. Hence, ethanol is able to dissolve both polar and non-polar substances. The partial charges on the polar part of the molecule are...



Question 7

An electrochemical cell was constructed using two half-cells. One half-cell consisted of tin metal and a tin(II) chloride solution and the other half-cell consisted of zinc metal and zinc chloride solution.

- Draw a diagram of the galvanic cell.
- Label the anode and the cathode.
- Indicate the direction of electron flow.

Criterion	Marks
Correctly drawn diagram with labels	2
Correct direction of electron flow	1

Question 8

- (a) Explain the term *biopolymer* **(1 mark)** and identify an example **(1 mark)**.

"Biopolymers are polymers that are made totally or in large part by living things." (Smith)

"Naturally occurring polymer generated using renewable resources like microorganisms or plants." (OTEN)

"A natural polymer." (Thickett)


"These are naturally occurring polymers such as cellulose, starch, and gluten..." (Syllabus Notes)

Examples... cellulose, starch, protein, rubber, carboxymethyl cellulose (CMC), PLA, PHA or PHB (poly β-hydroxyalkanoate), etc.


Question 8 continues on page 6

Question 8 (continued)

(b) Cellulose is a polymer of β -glucose. A β -glucose molecule is shown below....



Draw a segment of a cellulose molecule by joining three glucose molecules together. **(3 marks)**



One mark for showing three glucoses joined via the – O – linkage with suitable termination, e.g.

One mark for showing one – O – linkage up and one down.

One mark for showing the three – CH₂OH groups alternately up, down, up.


Question 9 (2 marks)

A student was asked to perform a first-hand investigation to compare the reactivities of hexane and hexene by observing their reactions with bromine water.

- (a) Describe the reaction(s) observed by the student when the procedures were carried out in a darkened laboratory. **(1 mark)**

The hexene immediately decolourised the bromine water.

- (b) Write an equation to show any addition reaction(s) that occurred. **(1 mark)**




☞ Production of bromohydroxyhexane is also acceptable.

Question 10 (3 marks)

Alkenes and their derivatives are important substances in the production of polymers. Polyvinyl chloride (PVC) is one such polymer.

- (a) Draw the structure of polyvinyl chloride showing three linked monomer units. **(1 mark)**



- (b) Describe **one** use of polyvinyl chloride and a property which makes it useful for this purpose. **(2 marks)**

e.g. PVC can be used to make electrical conduit because it is an electrical insulator.
The presence of Cl atoms in the polymer make it flame resistant.

e.g. PVC is used for credit cards because it is a tough, rigid polymer.

Must relate property to usage for **two marks**.

Usage without related property gains **one mark**.

Property without specific use gains **one mark**.