



SAINT IGNATIUS' COLLEGE

Trial Higher School Certificate

2004

MATHEMATICS EXTENSION 1

**1:25pm – 3:30 pm
Thursday 19th August 2004**

Directions to Students

- Reading Time : 5 minutes
- Working Time : 2 hours
- Write using blue or black pen. (sketches in pencil).
- Board approved calculators may be used.
- A table of standard integrals is provided at the back of this paper.
- All necessary working should be shown in every question.
- **Answer each question in the booklets provided and clearly label your name and teacher's name.**
- Total Marks 84
- Attempt Question 1 – 7
- All questions are of equal value

This paper has been prepared independently of the Board of Studies NSW to provide additional exam preparation for students. Although references have been reproduced with permission of the Board of Studies NSW, the publication is in no way connected with or endorsed by the Board of Studies NSW.

Total marks (84)

Attempt Questions 1 – 7

All questions are of equal value

Answer each question in a SEPARATE writing booklet.

QUESTION 1 (12 Marks) Use a SEPARATE writing booklet. Marks

(a) Solve $\frac{5}{2x-1} < 3$. 3

(b) Find the acute angle between the lines $2x - y + 1 = 0$ and $x + 3y - 4 = 0$. 3
Give answer to the nearest degree.

(c) Find the coordinates of the point that divides the interval joining $(-2, 5)$ and $(8, -9)$ internally in the ratio $2 : 3$. 2

(d) If α, β, γ are the roots of the equation $x^3 - 5x^2 - 3x + 2 = 0$, 2
find the value of $\frac{1}{\alpha\beta} + \frac{1}{\beta\gamma} + \frac{1}{\gamma\alpha}$.

(e) Write down the general solution, in terms of π , of the equation 2
$$\cos \theta = -\frac{1}{2}.$$

QUESTION 2 (12 Marks) Use a SEPARATE writing booklet.

Marks

- (a) Use the substitution $x = u^2 + 1$, for $u > 0$, to evaluate

4

$$\int_1^5 (x+1)\sqrt{x-1} \, dx.$$

- (b) Evaluate $\int_0^{\frac{\pi}{4}} \sin^2\left(\frac{1}{2}x\right) dx$.

3

- (c) Prove, using the principle of mathematical induction, that $9^{n+2} - 4^n$ is divisible by 5, for n a positive integer.

5

QUESTION 3 (12 Marks) Use a SEPARATE writing booklet.

Marks


- (a) Find the exact value of $\tan\left(2\sin^{-1}\frac{3}{4}\right)$. **3**
- (b) Consider the function $f(x) = \sin^{-1}(x+1) + \frac{\pi}{2}$.
- (i) What is the domain of $f(x)$? **1**
- (ii) Sketch the graph of $y = f(x)$. **2**
- (c) Consider the function $f(x) = \log_e(2x+1)$.
- (i) Write down the domain of $f(x)$. **1**
- (ii) Find the inverse function of $f(x)$, and write it in the form $f^{-1}(x) = \dots\dots$ **2**
- (iii) Find the gradients of the graphs of $y = f(x)$ and $y = f^{-1}(x)$ at the origin. **1**
- (iv) On the same diagrams, draw the graphs of $y = f(x)$ and $y = f^{-1}(x)$. **2**

QUESTION 4 (12 Marks) Use a SEPARATE writing booklet.

Marks

- (a) Find the coefficient of x^3 in the expansion of $(2 - x)(1 + x)^5$. 3

(b)



A swimming pool is 20 metres long, 8 metres wide, 4 metres deep at one end, and zero depth at the other end. The floor of the pool is a plane rectangular surface.


- (i) When the depth of water at the deeper end is h metres, show that the volume ($V \text{ m}^3$) of water in the pool is given by $V = 20h^2$. 2
- (ii) If water is being poured into the pool at the rate of $2 \text{ m}^3/\text{minute}$, find the rate at which the depth of the water is increasing at the deepest end, when the depth is 1 metre. 2
- (c) The value of a home business, $\$V$, is increasing at a rate proportional to the amount by which the value is less than $\$4000$.
i.e. $\frac{dV}{dt} = k(4000 - V)$
Initially, the value of the business was $\$2000$ and after 5 years it was $\$3000$.
- (i) Show that $V = 4000 - Ae^{-kt}$ satisfies this equation. 1
- (ii) Find the value of A and the value of k to 4 decimal places. 2
- (iii) Find the number of years for the value of the business to grow to $\$3800$. 2

QUESTION 5 (12 Marks) Use a SEPARATE writing booklet.

Marks

- (a) (i) Show that the derivative of $x^2 e^{-x}$ is $xe^{-x}(2-x)$. 1
- (ii) Show that $x^2 e^{-x} = 0.4$ has a root between $x = 1$ and $x = 2$. 1
- (iii) Use Newton's approximation to find an approximation to the root of $x^2 e^{-x} = 0.4$, taking $x = 1$ as a first approximation. 3

(b)



$ABCD$ is a cyclic quadrilateral in which $AB = AC$, and CD is produced to E .
Prove that AD bisects the angle BDE .

3

- (c) In the expansion of $(3+2x)^8$, c_r is the coefficient of x^r .

- (i) Show that $\frac{c_r}{c_{r-1}} = \frac{18-2r}{3r}$. 2
- (ii) Hence or otherwise find the largest coefficient in the expansion of $(3+2x)^8$. 2

QUESTION 6 (12 Marks) Use a SEPARATE writing booklet.

Marks

- (a) The position of a particle at time t is given by:

$$x = 3 \sin 2t - 4 \cos 2t.$$

- | | | |
|-------|---|---|
| (i) | Show that this equation satisfies $\ddot{x} = -n^2 x$. | 2 |
| (ii) | What is the initial velocity of the particle? | 1 |
| (iii) | At what time does the particle first come to rest? | 3 |

- (b) The acceleration of a particle at position x is given by:


$$\ddot{x} = -\frac{1}{4x^3}.$$

Initially the particle is at $x = 1$ moving with a velocity of $\frac{1}{2}$ unit in the positive direction.

- | | | |
|-----|--|---|
| (i) | Prove that the velocity of the particle at position x is given by: | 3 |
|-----|--|---|

$$v = \frac{1}{2x}.$$

- | | | |
|------|---|---|
| (ii) | Hence find the position of the particle at time t . | 3 |
|------|---|---|

QUESTION 7 (12 Marks) Use a SEPARATE writing booklet.**Marks**

An object is thrown from ground level with a speed of 40 m/s at an angle of 60° to the horizontal. Assume acceleration due to gravity is 10 m/s^2 and neglect air resistance.

- (a) Find equations for x and y in terms of time t seconds, starting from the acceleration equations $\ddot{x} = 0$ and $\ddot{y} = -10$, and hence show that: **4**

$$y = \sqrt{3}x - \frac{x^2}{80}.$$

- (b) The object is thrown up a slope with a gradient of $\frac{1}{4}$. **2**
Show that the horizontal distance travelled by the object when it lands on the slope is given by:

$$x = 80\sqrt{3} - 20.$$

- (c) Hence find the distance OA (to the nearest metre) up the slope from the point of projection to the point of landing. **2**


- (d) Show that the maximum height reached by the object above the slope is **4**
 $(61.25 - 10\sqrt{3})$ metres.

End of paper

(2004 Trials)

MATHEMATICS EXTENSION 1 - QUESTION 1

(a) $\frac{5}{2x-1} < 3$
 $5(2x-1) < 3(2x-1)^2$
 $3(2x-1)^2 - 5(2x-1) > 0$
 $(2x-1)[3(2x-1) - 5] > 0$
 $(2x-1)(6x-8) > 0$
 $x < \frac{1}{2} \text{ or } x > \frac{4}{3}$ ③



(b) $2x - y + 1 = 0$ $m_1 = 2$
 $x + 3y - 4 = 0$ $m_2 = -\frac{1}{3}$
 $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$
 $= \left| \frac{2 - (-\frac{1}{3})}{1 + 2(-\frac{1}{3})} \right|$
 $= \frac{2\frac{1}{3}}{\frac{1}{3}}$
 $= 7$
 $\theta = 82^\circ$ ③

(c) $(-2, 5), (8, -9)$ $2:3$
 $\left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right) = \left(\frac{2 \times 8 + 3 \times (-2)}{2+3}, \frac{2 \times (-9) + 3 \times 5}{2+3} \right)$
 $= \left(2, -\frac{3}{2} \right)$ ②

(d) $x^3 - 5x^2 - 3x + 2 = 0$
 $\frac{1}{\alpha\beta} + \frac{1}{\beta\gamma} + \frac{1}{\gamma\alpha} = \frac{\gamma + \alpha + \beta}{\alpha\beta\gamma} = \frac{5}{-2} = -2.5$ ②

(e) $\cos \theta = -\frac{1}{2}$
 $\theta = 2n\pi \pm \frac{2\pi}{3}$
 OR $\theta = (2n+1)\pi \pm \frac{\pi}{3}$
 OR equivalent. ②

Marks Awarded	Marker's Comments
(a). 1 mark 1 mark 1 mark	$5(2x-1) < 3(2x-1)^2$... or critical points. $2(2x-1)(3x-4) > 0$ $x < \frac{1}{2}, x > \frac{4}{3}$ or correctly solving the inequality obtained (unless trivial).
(b). 1 mark 1 mark 1 mark	$m_1 = 2 / m_2 = -\frac{1}{3}$. $\tan \theta = \left \frac{m_1 - m_2}{1 + m_1 m_2} \right $... understanding this is formula to use even if not stated in this form. number crunching $\rightarrow 82^\circ$.
(c). 1 mark. 1 mark	$\frac{(2 \times 8) + 3(-2)}{2+3}$ $\frac{2 \times (-9) + 3 \times 5}{2+3}$ $(2, -\frac{3}{2})$
(d) 1 mark 1 mark	$\frac{\delta + \beta + \alpha}{\alpha\beta\gamma}$ $\alpha + \beta + \alpha = 5$ and $\alpha\beta\gamma = -2$.
(e) 1 mark. 1 mark	$2n\pi$ or equivalent $\pm \frac{2\pi}{3}$ or equivalent (must be $\pm$)

Question 2

$$2) \int_1^5 (x+1)\sqrt{x-1} dx$$

$$x = u^2 + 1$$

$$\frac{dx}{du} = 2u$$

$$dx = 2u du \quad [1]$$

$$x = 1 \quad u = 0$$

$$x = 5 \quad u = 2$$

$$= \int_0^2 (u^2 + 2)\sqrt{u} \cdot 2u du$$

$$= \int_0^2 (2u^4 + 4u^2) du \quad [2]$$

$$= \left[\frac{2u^5}{5} + \frac{4u^3}{3} \right]_0^2$$

$$= \left[\left(\frac{64}{5} + \frac{32}{3} \right) - (0) \right]$$

$$= \frac{352}{15} = 23\frac{7}{15} \quad [1]$$

$$b) \int_0^{\frac{\pi}{4}} \sin^2\left(\frac{1}{2}x\right) dx = \int_0^{\frac{\pi}{4}} \frac{1}{2}(1 - \cos x) dx \quad [1]$$

$$= \frac{1}{2} \left[x - \sin x \right]_0^{\frac{\pi}{4}} \quad [1]$$

$$= \frac{1}{2} \left[\frac{\pi}{4} - \frac{1}{\sqrt{2}} \right] \quad [1]$$

c) Prove $9^{n+2} - 4^n$ is divisible by 5.

[1]

i) let $n=1$ $9^3 - 4 = 725$ $\therefore$ True for $n=1$

ii) Assume true for $n=k$ i.e. $9^{k+2} - 4^k = 5m$ (m is pos integer) [1]

$$\begin{aligned} \text{iii) When } n=k+1 \quad 9^{k+3} - 4^{k+1} &= 9(9^{k+2}) - 4(4^k) \\ &= 9(5m + 4^k) - 4(4^k) \\ &= 45m + 9 \cdot 4^k - 4 \cdot 4^k \\ &= 45m + 5 \cdot 4^k \\ &= 5[9m + 4^k] \end{aligned} \quad [2]$$

This is divisible by 5.

$\therefore$ if true for $n=k$, then true for $n=k+1$

$\therefore$ by induction, it is true for all $n \in \mathbb{N}$ [1]

Marks Awarded	Marker's Comments
<u>1</u>	Many mucked up the conversion.
<u>2</u>	Those with correct step one went on to get the correct integral and find the correct numeric value.
<u>1</u>	Many did not know $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$
<u>1</u>	most integrated correctly.
<u>1</u>	most had correct evaluation.
<u>1</u>	Almost all proved true for $n=1$.
<u>1</u>	$9^{k+2} - 4^k = 5m$ well stated. Most DID NOT state that m was a positive integer.
<u>2</u>	Many could not set out the correct steps for this section. Many used " m " again. Is it the same " m " used earlier?
<u>1</u>	Many were lazy in their final statement. $\therefore$ most did not get this mark.

MATHEMATICS EXTENSION I - QUESTION 3

Mathematics Extension One: Question Number

3

Marker: Mr M'Car

(a) $\tan(2 \sin^{-1} \frac{3}{4})$


Let $\theta = \sin^{-1} \frac{3}{4} \therefore \sin \theta = \frac{3}{4}$

$\tan(2 \sin^{-1} \frac{3}{4}) = \tan 2\theta$

$= \frac{2 \tan \theta}{1 - \tan^2 \theta}$

$= \frac{2 \times \frac{3}{4}}{1 - \frac{9}{16}}$

$= -3\sqrt{7}$




③

(b) $f(x) = \sin^{-1}(x+1) + \frac{\pi}{2}$

(i) Domain: $-1 \leq x+1 \leq 1 \therefore -2 \leq x \leq 0$

①

(ii)



②

(c) $f(x) = \log_e(2x+1)$

(i) Domain: $2x+1 > 0 \therefore x > -\frac{1}{2}$

①

(ii) $y = \log_e(2x+1)$

$2x+1 = e^y$

Inverse is: $2y+1 = e^x$

$y = \frac{1}{2}(e^x - 1)$

$f^{-1}(x) = \frac{1}{2}(e^x - 1)$


②

(iii) $f'(x) = \frac{2}{2x+1}$; $f'(0) = 2$

$\frac{d}{dx} f'(x) = \frac{1}{x^2}$; At $x=0$, $\frac{d}{dx} f'(x) = \frac{1}{x^2}$

①

(v)



②

(a)

Marks Awarded	Marker's Comments
1	$\tan \theta = \frac{3}{\sqrt{7}}$
1	double angle formula substit
1	correct final answer
	Note - 1 mark awarded for calculator answer ≈ 7.94
(b)	
(i) 1	Correct domain $-2 \leq x \leq 0$
(ii) 1	Correct shape
1	Correct position
	Note - Use a stencil !!
(c)	
(i) 1	Correct domain $x > -\frac{1}{2}$
(ii) 1	Interchange $x \leftrightarrow y$
1	Make y the subject
	(Generally well done)
(iii) 1	Either answer $f'(0) = 2$
	or $\frac{d}{dx} f'(0) = \frac{1}{x^2}$
(iv) 1	One correct function
or	
2	Must pass thru' origin } required for full marks
	Show both asymptotes }
	intersect twice }

(poorly answered)


Question 4

$$2) (2-x)(1+x)^5 = (2-x)(1+5x+10x^2+10x^3+\dots)$$

$$= 2(10x^3) + (-x)(10x^2)$$

$$= 20x^3 - 10x^3$$

$$= 10x^3 \quad \therefore \text{Coefficient} = 10$$

i) 

$$\frac{h}{x} = \frac{4}{20} \quad V = \frac{1}{2} h x \times 8$$

$$20h = 4x \quad = \frac{1}{2} h \cdot 5h \cdot 8$$

$$x = 5h \quad = 20h^2$$

ii) $\frac{dV}{dt} = 2$ $h = 1$ $\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt}$

$$\frac{dV}{dh} = ? \quad 2 = 40h \times \frac{dh}{dt}$$

$$\frac{dh}{dt} = 0.05$$

iii) $\frac{dV}{dt} = k(4000 - V)$

i) $V = 4000 - Ae^{-kt}$

$$\frac{dV}{dt} = -k(-Ae^{-kt})$$

$$= +k(4000 - V)$$

ii) $t = 0 \quad V = 2000 \quad 2000 = 4000 - Ae^0$

$$A = 2000$$

$t = 5 \quad V = 3000 \quad 3000 = 4000 - 2000e^{-5k}$

$$e^{-5k} = \frac{1}{2}$$

$$k = -\frac{1}{5} \ln 0.5 = 0.1386$$

iii) $3800 = 4000 - 2000e^{-kt}$

$$e^{-kt} = 0.1$$

Mathematics Extension One: Question Number

4

Marker: NM

Marks Awarded	Marker's Comments
1	Correct expansion
1	Correct collection of coefficients
1	10. (Well done)
1	Correct explanation of why ratio was 5:1.
1	Correct explanation of why $V = 20h^2$. (Poorly done)
1	$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt}$ or equivalent.
1	0.05 or $\frac{1}{20}$ or equivalent. (Well done)
1	Use of $Ae^{-kt} = 4000 - V$.
1	Evaluate A
1	Evaluate k (correct dp's)
1	Correct equation.
1	Correct value of t.

MATHEMATICS EXTENSION I - QUESTION 5.

- (a) $y = x^2 e^{-x}$
- (i) $\frac{dy}{dx} = e^{-x}(2x) + x^2(-e^{-x})$
 $= 2xe^{-x} - x^2 \cdot e^{-x}$
 $= xe^{-x}(2-x)$ ①
- (ii) Let $f(x) = x^2 e^{-x} - 0.4$: $f(1) = e^{-1} - 0.4 = -0.0360$
 $f(2) = 4e^{-2} - 0.4 = 0.19 > 0$
 Since $f(x)$ is continuous and $f(1), f(2)$ have opposite signs,
 a root lies between 1 and 2. ①
- (iii) $x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$ $f'(1) = e^{-1} \times 1$
 $= 1 - \frac{-0.03}{0.3679}$ $= 0.3679$
 $= 1.08$ ③

- b) Let $\angle ADE = \theta$
- $\therefore \angle ABC = \theta$ (exterior angle cyclic quad)
 - $\therefore \angle ACB = \theta$ (base angles isosceles triangle)
 - $\therefore \angle ADB = \theta$ (angles in same segment)
 - $\therefore \angle ADE = \angle ADB$
 - $\therefore AD$ bisects $\angle BDE$.



- (i) $\frac{C_r}{C_{r+1}} = \frac{\binom{8}{r} 3^{8-r} 2^r}{\binom{8}{r+1} 3^{8-r-1} 2^{r+1}}$
 $= \frac{2}{3} \times \frac{8!}{r!(8-r)!} \times \frac{(r-1)!(9-r)!}{8!}$
 $= \frac{2}{3} \times \frac{9-r}{r}$
 $= \frac{18-2r}{3r}$ ②
- (ii) $\frac{18-2r}{3r} > 1$
 $18 > 5r$
 $r < 3\frac{3}{5}$
 Greatest coefficient = coefficient of x^3
 $= \binom{8}{3} 3^5 2^3$
 OR 108 864 ②

Mathematics Extension One: Question Number

5

Marker: Mr M

Marks Awarded	Marker's Comments
(a)	
(i) 1	Correct use of product rule
(ii) 1	Show a change of sign
(iii) 1	$f(1) = -0.03$
1	$f'(1) = 0.368$
1	Correct estimate $x_1 = 1.087$
	Note: must use $f(x) = x^2 e^{-x} - 0.4$
	Poorly answered by many students.
(b)	
1	Exterior $\angle$ of cyclic quad.
1	Base $\angle$ s of isos. Δ
1	Angles in same segment.
	Note - very poor structure
	- drawing a diagram helps.
(c)	
(i) 1	$\frac{\binom{8}{r} 3^{8-r} 2^r}{\binom{8}{r+1} 3^{8-r-1} 2^{r+1}} \approx \frac{n-r+1}{r} \cdot \frac{1}{a}$
1	Use of factorial definition to correctly simplify.
(ii) 1	Solving inequality $r < 3\frac{3}{5}$
1	Finding greatest term 108 864
	Note: part (i) poorly answered.

Question 6

i) $x = 3 \sin 2t - 4 \cos 2t$

$\dot{x} = 6 \cos 2t + 8 \sin 2t$

$\ddot{x} = -12 \sin 2t + 16 \cos 2t$

$= -4[3 \sin 2t - 4 \cos 2t]$

$= -4x \quad (\text{ie } -n^2x)$

ii) $t=0 \quad \dot{x} = 6 \cos 0 + 8 \sin 0$
 $= 6$

iii) $\dot{x} = 0 \quad 6 \cos 2t + 8 \sin 2t = 0$

$8 \sin 2t = -6 \cos 2t$

$\tan 2t = -\frac{3}{4}$

$2t = \pi - 0.6435$

$t = 1.249$

c) i) $\ddot{x} = -\frac{1}{4x^3}$

$\frac{d}{dx}(\frac{1}{2}v^2) = -\frac{1}{4}x^{-3}$

$\frac{1}{2}v^2 = \frac{1}{8}x^{-2} + C$

$v = \frac{1}{2}x = 1 \quad \frac{1}{2}(\frac{1}{4}) = \frac{1}{8} + C$

$\therefore C = 0$

$\frac{1}{2}v^2 = \frac{1}{8x^2}$

$v = \frac{1}{2x}$

ii) $\frac{dx}{dt} = \frac{1}{2x} \quad \frac{dt}{dx} = 2x$

$t = x^2 + C$

Marks Awarded	Marker's Comments
a) i) 1	Correct expression for $\ddot{x}$
1	Correct manipulation to $\ddot{x} = -n^2x$ (well done)
ii) 1	$\dot{x} = 6$ (Very well done)
iii) 1	Let $\dot{x} = 0$ (Generally OK)
1	$\tan 2t = -3/4$ Some poor solutions do
1	$t = 1.249$ $\tan 2t = -3/4$
b) i) 1	Correct version of $\ddot{x} = \frac{d}{dx}(\frac{1}{2}v^2)$
1	Evaluate $C = 0$
1	Correct "tidy up" to $v = \frac{1}{2x}$ (many forgot C)
ii) 1	For $\frac{dt}{dx} = 2x$
1	Evaluate C.
1	Manipulate to $x = \sqrt{t+1}$

MATHEMATICS EXTENSION 1 - QUESTION 7

a) Initially, $\dot{x} = V \cos \theta = 40 \cos 60^\circ = 20$; $\dot{y} = V \sin \theta = 40 \sin 60^\circ = 20\sqrt{3}$

$$\ddot{x} = 0$$

$$\ddot{y} = -10$$

$$\dot{x} = c$$

$$\dot{y} = -10t + k$$

$$\dot{x} = 20$$

$$\text{When } t=0, \dot{y} = 20\sqrt{3} \therefore k = 20\sqrt{3}$$

$$x = 20t + c'$$

$$\therefore \dot{y} = 20\sqrt{3} - 10t$$

$$\text{When } t=0, x=0 \therefore c' = 0$$

$$y = 20\sqrt{3}t - 5t^2 + k'$$

$$\therefore x = 20t \quad (A)$$

$$\text{When } t=0, y=0 \therefore k' = 0$$

$$\therefore y = 20\sqrt{3}t - 5t^2 \quad (B)$$

$$\text{From (A), } t = \frac{x}{20} \therefore y = 20\sqrt{3} \left(\frac{x}{20}\right) - 5 \left(\frac{x}{20}\right)^2$$

$$y = \sqrt{3}x - \frac{x^2}{80}$$

(4)

b) Equation of slope: $y = \frac{1}{4}x$ $\therefore \sqrt{3}x - \frac{x^2}{80} = \frac{1}{4}x$

$$80\sqrt{3}x - x^2 = 20x$$

$$(80\sqrt{3} - 20)x - x^2 = 0$$

$$x[80\sqrt{3} - 20 - x] = 0$$

$$x = 0 \text{ or } x = 80\sqrt{3} - 20$$

$$\therefore \text{Horizontal distance is } 80\sqrt{3} - 20$$

(2)

c) From $y = \frac{1}{4}x$, $y = \frac{1}{4}(80\sqrt{3} - 20) = 29.64$

$$x = 80\sqrt{3} - 20 = 118.56$$

$$\text{Distance OA} = \sqrt{118.56^2 + 29.64^2}$$

$$= 122 \text{ metres (nearest metre)}$$

(2)

d) Height above slope: $H = \sqrt{3}x - \frac{x^2}{80} - \frac{1}{4}x$

$$\frac{dH}{dx} = \sqrt{3} - \frac{1}{4} - \frac{x}{40}$$

$$\frac{dH}{dx} = 0 \therefore x = 40(\sqrt{3} - \frac{1}{4})$$

This is for maximum value of H (concave down parabola)

$$\text{When } x = 40(\sqrt{3} - \frac{1}{4}), H = (\sqrt{3} - \frac{1}{4})40(\sqrt{3} - \frac{1}{4}) - \frac{1}{80} \cdot 40^2(\sqrt{3} - \frac{1}{4})^2$$

$$= 40(\sqrt{3} - \frac{1}{4})^2 - 20(\sqrt{3} - \frac{1}{4})^2$$

$$= 20(\sqrt{3} - \frac{1}{4})^2$$

$$= 20(3 - \frac{\sqrt{3}}{2} + \frac{1}{16})$$

$$= 61.25 - 10\sqrt{3}$$

$$\text{Maximum height is } (61.25 - 10\sqrt{3}) \text{ m.}$$

(4)

Mathematics Extension One: Question Number 7

Marker: C3F

Marks Awarded	Marker's Comments
(a) 1 mark. 1 mark. 1 mark. 1 mark.	$\begin{matrix} \dot{x} = 20 \\ \dot{y} = 20\sqrt{3} \end{matrix} > \text{in whatever form and}$ whenever stated in answer $x = 20t \dots \dots$ deriving equation / finding c' $y = 20\sqrt{3}t - 5t^2 \dots$ deriving equation and finding k and k' (mark not awarded if k and k' ignored) $t = \frac{x}{20} \rightarrow y = 20\sqrt{3} \left(\frac{x}{20}\right) - 5 \left(\frac{x}{20}\right)^2$
(b) 1 mark 1 mark.	$\frac{1}{4}x = \sqrt{3}x - \frac{x^2}{80}$ $x = 80\sqrt{3} - 20$ / derived from above.
(c) 1 mark 1 mark	$y = \frac{1}{4}(80\sqrt{3} - 20)$ and $x = 80\sqrt{3} - 20$ distance OA = 122 m.
(d) 1 mark 1 mark 1 mark 1 mark	$H = \sqrt{3}x - \frac{x^2}{80} - \frac{1}{4}x$ $\frac{dH}{dx} =$ $x = 40(\sqrt{3} - \frac{1}{4})$ $H = 61.25 - 10\sqrt{3}$ Note: $t = 2\sqrt{3} - \frac{1}{2}$ Not $t = 2\sqrt{3}$. This is because lamp is on a slope and maximum height is not in middle