



SAINT IGNATIUS' COLLEGE

Trial Higher School Certificate

2007

MATHEMATICS EXTENSION 1

8:45am – 10:50am
Friday 31st August 2007

Directions to Students

• Reading Time: 5 minutes	• Total Marks: 84
• Working Time: 2 hours	
• Write using blue or black pen (sketches in pencil).	• Attempt Questions 1 – 7
• Board approved calculators may be used	• All questions are of equal value
• A table of standard integrals is provided at the back of this paper.	
• All necessary working should be shown in every question.	
• Answer each question in the booklets provided and clearly label your name and teacher's name.	

This paper has been prepared independently of the Board of Studies NSW to provide additional exam preparation for students. Although references have been reproduced with permission of the Board of Studies NSW, the publication is in no way connected with or endorsed by the Board of Studies NSW.

QUESTION 1 (use a SEPARATE writing booklet)

- (a) Given that if $y = \sin^{-1}\left(\frac{x}{a}\right)$ then $\frac{dy}{dx} = \frac{1}{\sqrt{a^2 - x^2}}$, where $x < |a|$,

write down an expression for $\frac{dy}{dx}$ if y equals:

- (i) $\sin^{-1}(x)$
- (ii) $\sin^{-1}\left(\frac{x}{7}\right)$
- (iii) $\sin^{-1}(7x)$ (3M)

- (b)
- (i) If $f(x) = x^3 - 3x^2 - 4x + 12$, show that $f(3) = 0$ (1M)
- (ii) Hence solve $x^3 - 3x^2 - 4x + 12 = 0$ (2M)

- (c) Prove that
- $$\frac{1}{3}[n(n+1)(n+2)] + (n+1)(n+2) = \frac{1}{3}(n+1)(n+2)(n+3)$$
- (2M)

- (d) Solve for x
- $$\frac{x-2}{x+4} \geq \frac{1}{3}$$
- (4M)

QUESTION 2 (use a SEPARATE writing booklet)

- (a) A particle P moves in a straight line so that its distance x from a central fixed point O at time t is given by

$$x = 2 \sin \left(5t + \frac{\pi}{6} \right)$$

- (i) Write down an expression for the velocity $\dot{x}$
- (ii) Write down an expression for the acceleration $\ddot{x}$
- (iii) The particle P is said to be executing Simple Harmonic Motion. Explain why this is so.
- (3M)

- (b) The parametric equations of a parabola are

$$x = 2t$$

$$y = 2t^2$$

Find the equation of the tangent to this parabola at the point

$$P(2, 2) \quad (2M)$$

- (c) Given that $x^4 + 3x^2 - 100 = 0$ has a root near $x = 3$, use Newton's method once to find a better approximation, giving your answer in exact form.
- (2M)

- (d) Find the numerical value of the co-efficient of x^0 in the expansion of

$$\left(2x^2 - \frac{1}{2x} \right)^{12} \quad (3M)$$

- (e) Find the sum of the six co-efficients (including the co-efficient of x^0) in the expansion of $(3 - x)^5$
- (2M)

QUESTION 3 (use a SEPARATE writing booklet)

- (a) Using the table of standard integrals, evaluate $\int_0^2 \frac{8dx}{x^2 + 4}$ (2M)

- (b) Find in radians all the values of x in the domain $0 \leq x \leq \pi$ which satisfy the equation

$$\sin 2x - \sin x = 0 \quad (3M)$$

- (c) If α , β and γ are the roots of the cubic equation

$$2x^3 - 8x^2 + x + 12 = 0,$$

write down the value of

- (i) $\alpha + \beta + \gamma$
(ii) $\alpha\beta + \beta\gamma + \gamma\alpha$
(iii) $\alpha^2 + \beta^2 + \gamma^2$ (3M)

- (d)

- (i) Show that $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$

- (ii) Deduce from (i) that if $x^2 + 4y^2 = 5$ then $\frac{dy}{dx} = \frac{-x}{4y}$


- (iii) A particle moves at a constant speed of $k \text{ ms}^{-1}$ on the circumference of the curve

$$x^2 + 4y^2 = 5$$

Find $\frac{dy}{dt}$ at the point P where $x = 2$, $y = -\frac{1}{2}$ and $\frac{dx}{dt} = 2 \text{ ms}^{-1}$

- (iv) What is the exact value of k , the constant speed? (4M)

QUESTION 4 (use a SEPARATE writing booklet)



(The above diagrams may help in your presentation)

A particle is projected with a velocity V from a point O at an angle of α to the horizontal, in the x, y plane. In the usual notation using the calculus or otherwise, prove that the horizontal distance x and the vertical distance y , travelled by the particle at time t are given by:

(i) $x = (V \cos \alpha)t$ (2M)

(ii) $y = (V \sin \alpha)t - \frac{1}{2}gt^2$
where g is the acceleration due to gravity. (3M)

(iii) By combining (i) and (ii) deduce that:

$$y = x \tan \alpha - \frac{gx^2}{2V^2}(1 + \tan^2 \alpha) \quad (3M)$$

(iv) Any of the above results may be used in attempting this problem:

A football is kicked at 15ms^{-1} and just passes over a crossbar 5m high and 15m away.


Taking $g = 10\text{ms}^{-2}$ vertically downwards, show that if α is the angle of projection then $\alpha = \frac{\pi}{4}$ or $\alpha = \tan^{-1} 2$

(3M)

(v) If the two values of α (from part iv) are given by A and B (respectively) where $A > B$, write down the exact value of $\tan(A - B)$

(1M)

QUESTION 5 (use a SEPARATE writing booklet)



The above diagram shows two curves

$$y = \cos \frac{x}{4} \quad \text{(i) (thin line)}$$

$$y = \cos^2 \left(\frac{x}{4} \right) \quad \text{(ii) (thick line)}$$

Also, A is the point $(0,1)$, B is the point $(\pi,0)$ and C is the point $(2\pi,0)$

- (i) Prove that the area enclosed by the curve $y = \cos \frac{x}{4}$ and the x and y axes is 4 square units.

(2M)

- (ii) The shaded region $APBA$ is enclosed by $y = \cos^2 \left(\frac{x}{4} \right)$ and the lines AB and $x = \pi$. Prove that the area of this shaded region is 1 square unit.

(3M)

- (iii) Prove that $\cos^4 \left(\frac{x}{4} \right) = \frac{1}{8} \left(3 + 4 \cos \frac{x}{2} + \cos x \right)$


(3M)

- (iv) The shaded region is now rotated through 2π radians about the x -axis. Show, using the result of part (iii) or otherwise that the exact volume of the solid generated is $\pi \left(1 + \frac{\pi}{24} \right)$ cubic units.

(4M)

QUESTION 6 (use a SEPARATE writing booklet)

- (a) If $y = 2007^x$, find an expression for $\frac{dy}{dx}$ in terms of x (2M)



- (b) AB is the diameter of a circle, centre O . P is a point on the circumference. The tangents at B and P meet at T , and AP , BT are produced to meet at Q , and $\hat{ABP} = x^\circ$

- (i) Copy the diagram into your script and explain why $\hat{APB} = 90^\circ$
- (ii) Noting without any explanation, that $\hat{OBQ} = 90^\circ$ or otherwise, explain why $\hat{TQP} = x^\circ$
- (iii) Explain why $\hat{TPQ} = x^\circ$
- (iv) If $x = 30$ show $PQ = 3AP$ (4M)

(c)

- (i) In the usual notation, prove that $\ddot{x} = v \frac{dv}{dx}$ (1M)

- (ii) A particle moves in a straight line and its acceleration at any time t is given by $\ddot{x} = v \frac{dv}{dx} = -e^{-2x}$, where x is the displacement and v the velocity at time t . Also, when $x = 0$, $v = 1$.

By starting with the result of part (i) or otherwise, prove that $v = e^{-x}$. (3M)

- (iii) It is also known that when $t = 0$, $x = 0$. Deduce from (ii), or otherwise prove, that the displacement x at time t is given by $x = \ln(t+1)$ (2M)


QUESTION 7 (use a SEPARATE writing booklet)

- (a) If $y = 3 \cos^{-1} \left(\frac{x}{2} \right)$, write down (i) the domain of x
(ii) the range of y (2M)

- (b) If $y = f(x) = \frac{kx+l}{mx-k}$ (where k, l, m are constants), prove that $x = f(y)$ (2M)

- (c) In attempting the problem below, you may assume without proof that:

- If $y = \cot \theta$, $\frac{dy}{d\theta} = -\operatorname{cosec}^2 \theta$
- If $y = \operatorname{cosec} \theta$, $\frac{dy}{d\theta} = -\operatorname{cosec} \theta \cot \theta$



The diagram shows a house at A , a school at D and a straight canal BC , where $ABCD$ is a rectangle with $AB = 2\text{ km}$ and $BC = 6\text{ km}$.

During the winter, when the canal freezes over, Danny travels from A to D by walking to a point P on the canal, skating along the canal to a point Q and then walking from Q to D . The points P and Q are chosen so that the angles APB and DQC are both equal to θ , $AP = QD$, $BP = QC$.

- (i) Show that $PQ = (6 - 4 \cot \theta)$ (1M)
- (ii) Given that Danny walks at a constant speed of 4 km h^{-1} , and skates at a constant speed of 8 km h^{-1} show that the time, T minutes, taken for Danny to go from A to D along this route is given by
- $$T = 15(3 + 4 \operatorname{cosec} \theta - 2 \cot \theta) \quad (3\text{M})$$
- (iii) Show clearly and carefully that, as θ varies, Danny's minimum time for the journey is approximately 97 minutes. (4M)

END OF PAPER

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

QUESTION 1


Suggested Solutions	Max Mark	Your Mark	Marker's Comments
(a) (i) $\frac{1}{\sqrt{1-x^2}} \quad x < 1$ ✓ (ii) $\frac{1}{\sqrt{49-x^2}} \quad x < 7$ ✓ (iii) $\frac{1}{\sqrt{\left(\frac{1}{7}\right)^2 - x^2}}$ OR $\frac{7}{\sqrt{1-49x^2}} \quad x < \left \frac{1}{7}\right$ ✓ No penalty for omitting domain of x	2 1		
(b) (i) Here $f(3) = 27 - 27 - 12 + 12 = 0$ (ii) From (i) $f(x) = (x-3)(x^2 - 4)$ Here $f(x) = (x-3)(x-2)(x+2) = 0$ Hence the required solutions are $x = 3, 2, -2$ *	1 1 1	✓ ✓ ✓	* No mark awarded if soln. not written.
(c) L.H.S. = $\frac{1}{3}[n(n+1)(n+2) + 3(n+1)(n+2)]$ We take out the factors $[(n+1)(n+2)]$ To get L.H.S. = $\frac{1}{3}[(n+1)(n+2)](n+3)$ (Award 2 marks for L.H.S. = R.H.S by direct multiplication) $\frac{1}{3}(n^3 + 6n^2 + 11n + 6)$	1 1	✓ ✓	Some tried Induction. (+) for 1st two steps; if no solution was shown.
(d) $\frac{x-2}{x+4}$ is undefined if $x+4=0$ ie if $x=-4$ Multiply both sides by $3(x+4)^2$ To get $3(x-2)(x+4) \geq (x+4)^2, (x \neq -4)$ ie $3x^2 + 6x - 24 \geq x^2 + 8x + 16, (x \neq -4)$ Hence $x^2 - x - 20 \geq 0, (x \neq -4)$ ie $(x+4)(x-5) \geq 0$, but $x \neq -4$ $\therefore x < -4$ or $x \geq 5$ To gain the 4th mark, the solution needs to mention $x \neq -4$ just the once.	1 1 1 1	✓ ✓ ✓ ✓	If $x \neq -4$ had to be mentioned, otherwise -1.

QUESTION 2

Suggested Solutions	Max Mark	Your Mark	Marker's Comments
(a) (i) $\dot{x} = 10 \cos\left(5t + \frac{\pi}{6}\right)$ ✓ (ii) $\ddot{x} = -50 \sin\left(5t + \frac{\pi}{6}\right)$ ✓ (iii) $\ddot{x} = -5^2 \left(2 \sin\left(5t + \frac{\pi}{6}\right)\right) = -5^2 x \quad \otimes$ At least 2 explanations are acceptable Eg (α) $x = 2 \sin\left(5t + \frac{\pi}{6}\right) = 2 \cos\left(5t - \frac{\pi}{3}\right)$, defines S.H.M (β) Equation $\otimes \quad \ddot{x} = -n^2 x \quad (n=5)$ defines S.H.M } ✓	1		
(b) Here $\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = 4t \times \frac{1}{2} = 2t$ ✓ OR $\frac{dy}{dx} = 2$ At P, $t=1 \therefore$ Equation is $y = 2x - 2$ OR $2x - y - 2 = 0$ ✓	1		most students used $\ddot{x} = -n^2 x$
(c) Here $f(x) = x^4 + 3x^2 - 100 \therefore f(3) = 8$ $f'(x) = 4x^3 + 6x \quad f'(3) = 126$ } both correct ✓	1		
In the usual notation $x_1 = x_0 - \frac{f(x)}{f'(x)} = 3 - \frac{8}{126}$ $= 2 \frac{59}{63} \quad (= \frac{185}{63})$ ✓	1		
(d) (i) Here $T_{R+1} = \binom{12}{R} \left(\frac{-x^{-1}}{2}\right)^R \cdot (2x^2)^{12-R}$ ✓ Power of x in $T_{R+1} = -R + 24 - 2R$ $= 24 - 3R = 0$ when $R = 8$ ✓ So $T_9 = \binom{12}{8} \left(\frac{1}{2^8}\right) 2^4 = \frac{12 \cdot 11 \cdot 10 \cdot 9}{4 \cdot 3 \cdot 2 \cdot 1} \times \frac{1}{16}$ $= \frac{495}{16} \quad (= 30.9353 = 30 \frac{15}{16})$ ✓	1		
(ii) To sum the 6 co-efficients, set $x = 1$ Required sum $(3 - 1)^5 = 2^5 = 32$ OR ✓ $3^5 + \binom{5}{1} 3^4 (-1) + \binom{5}{2} 3^3 (-1)^2 + \binom{5}{3} 3^2 (-1)^3 + \binom{5}{4} 3 (-1)^4 + (-1)^5$ ✓	1		
	1		

QUESTION 3

Suggested Solutions	Max Mark	Your Mark	Marker's Comments
(a) $I = \left[4 \tan^{-1} \left(\frac{x}{2} \right) \right]_0^2$ $= 4 \left[\tan^{-1} 1 - \tan^{-1} 0 \right] = \frac{4\pi}{4} = \pi$	1	✓	
	1	π ✓	
(b) $2 \sin x \cos x - \sin x = 0$ $\therefore \sin x (2 \cos x - 1) = 0 \rightarrow \sin x = 0$ or $\cos x = \frac{1}{2}$ So $x = 0, \pi$ or 2π or $x = \frac{\pi}{3}$ or $\frac{5\pi}{3}$ But $0 \leq x \leq \pi$ $x = 0, \frac{\pi}{3}, \pi$ (Penalty of 1 if degrees are used instead of radians)	1	✓	
	1	✓	
	1	0, $\frac{\pi}{3}$, π ✓	Any answers other than these three did not get a mark here.
(c) (i) 4 (ii) $\frac{1}{2}$	1	4 ✓	
	1	$\frac{1}{2}$ ✓	
(iii) $\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \alpha\gamma) = 15$	1	15 ✓	
(d) (i) Use product rule $\frac{d}{dx}(y \cdot y) = yy' + y' y = 2yy'$ OR note $\frac{d}{dx}(y^2) = \frac{d}{dy}(y^2) \cdot \frac{dy}{dx} = 2yy'$ (ii) Differentiate both sides with respect to x To get $2x + 8y \cdot \frac{dy}{dx} = 0 \therefore \frac{dy}{dx} = \frac{-x}{4y}$ (iii) $\frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt} = \frac{-x}{4y} (2) = 2ms^{-1}$ (iv) Note: we focus on <u>speed</u> (and not velocity) At P, speed along x axis = speed along the y axis $= 2ms^{-1} \therefore$ Resulting speed $= \sqrt{2^2 + 2^2}$ $k = 2\sqrt{2}$	1	$2y \frac{dy}{dx}$ ✓	Some tried to reproduce the Question - No marks. All other methods were accepted.
	1	$\frac{-x}{4y}$ ✓	
	1	2 ✓	
	1	$2\sqrt{2}$ ✓	



QUESTION 4

Suggested Solutions	Max Mark	Your Mark	Marker's Comments
(i) HORIZONTAL MOTION The initial velocity in a horizontal direction is $V \cos \alpha$ and since no forces are acting in the horizontal direction, this velocity will remain uniform. $\ddot{x} = 0$ That is $\frac{dx}{dt} = V \cos \alpha$ ✓ Thus $x = (V \cos \alpha)t + c$ But $x = 0$ when $t = 0$, thus $c = 0$ Hence $x = (V \cos \alpha)t \dots (i)$ ✓	1		
(ii) VERTICAL MOTION The initial velocity in a vertical direction is $V \sin \alpha$. If we take the upwards direction as positive the particle experiences a uniform acceleration of $-g$ due to gravity. Thus $\frac{d^2y}{dt^2} = -g$ ✓ Hence $\frac{dy}{dt} = -gt + c$ But $\frac{dy}{dt} = V \sin \alpha$ when $t = 0$, thus $c = V \sin \alpha$ Thus $\frac{dy}{dt} = V \sin \alpha - gt$ ✓ Hence $y = (V \sin \alpha)t - \frac{1}{2}gt^2 + k$ But $y = 0$ when $t = 0$, thus $k = 0$ Giving $y = (V \sin \alpha)t - \frac{1}{2}gt^2 \dots (ii)$ ✓	1		<u>Note:</u> You must show the initial conditions to work out the constants.
(iii) Eliminating t From (i) $t = \frac{x}{V \cos \alpha}$ ✓ Substituting in (ii) $y = \frac{(V \sin \alpha)x}{V \cos \alpha} - \frac{1}{2}g\left(\frac{x}{V \cos \alpha}\right)^2$ ✓ i.e. $y = x \tan \alpha - \frac{gx^2}{2V^2} \sec^2 \alpha = x \tan \alpha - \frac{gx^2}{2V^2}(1 + \tan^2 \alpha) \dots (iii)$ ✓	1		Remember $\sec^2 \alpha = 1 + \tan^2 \alpha$
(iv) Using equation (iii) $5 = 15 \tan \alpha - 5 - 5 \tan^2 \alpha$, ✓ $5 = 15t - 5 - 5t^2$ where $t = \tan \alpha$ $\therefore 5t^2 - 15t + 10 = 0$ $t^2 - 3t + 2 = 0$ ✓ $(t-2)(t-1) = 0$ Hence $t = 1$ and $t = 2$ $\therefore \alpha = \frac{\pi}{4}$ & $\alpha = \tan^{-1} 2$ ✓	2		1st mark: correct substitution
	1		2nd mark: correct equation
(v) $\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B} = \frac{2-1}{1+2} = \frac{1}{3}$	1		N.B question $A > B \therefore \text{answer} = \frac{1}{3}$

MATHEMATICS EXTENSION 1
Solutions, Marking Scheme & Comments

QUESTION 5

Suggested Solutions	Max Mark	Your Mark	Marker's Comments
(i) From diagram, Area = $\int_0^{2\pi} \cos \frac{x}{4} dx$	1	✓	
i.e. $A = \left[4 \sin \frac{x}{4} \right]_0^{2\pi} = 4 \sin \frac{\pi}{2} - 0 = 4$	1	✓	
(ii) Shaded Area = $\int_0^{\pi} \cos^2 \left(\frac{x}{4} \right) dx - \Delta BOA$	1		
$= \frac{1}{2} \int_0^{\pi} \left(\cos \frac{x}{2} + 1 \right) dx - \frac{1}{2} (\pi \times 1)$			$\cos^2 A = \cos 2A + 1$
$= \frac{1}{2} \left[2 \sin \left(\frac{x}{2} \right) + x \right]_0^{\pi} - \frac{\pi}{2}$	1		$\frac{1}{2} \left[2 \sin \left(\frac{\pi}{2} \right) + \pi \right]_0^{\pi} - \frac{\pi}{2}$
$= \left[\sin \frac{\pi}{2} + \frac{\pi}{2} - 0 - 0 \right] - \frac{\pi}{2} = 1$	1		$= \frac{1}{2} \left[2 \times \sin \frac{\pi}{2} + \pi - 0 \right] - \frac{\pi}{2}$ $= \sin \frac{\pi}{2} + \frac{\pi}{2} - \frac{\pi}{2} = 1$
(iii) Since $\cos^2 \left(\frac{x}{4} \right) = \frac{1}{2} \left[1 + \cos \left(\frac{x}{2} \right) \right]$	1	✓	
Therefore $\cos^4 \frac{x}{4} = \frac{1}{4} \left[1 + \cos \left(\frac{x}{2} \right) \right]^2$			
$\therefore \cos^4 \frac{x}{4} = \frac{1}{4} \left[\left(1 + 2 \cos \frac{x}{2} + \cos^2 \left(\frac{x}{2} \right) \right) \right]$	1	✓	
$= \frac{1}{4} \left[1 + 2 \cos \left(\frac{x}{2} \right) + \frac{1}{2} (1 + \cos x) \right]$			
$= \frac{1}{8} \left(3 + 4 \cos \frac{x}{2} + \cos x \right)$	1	✓	
(iv) Required Volume = $\pi \int_0^{\pi} \cos^4 \left(\frac{x}{4} \right) dx - v_c$	1	✓	
where v_c = volume of a cone with radius unity and height π .			
ie $v_c = \frac{1}{3} \cdot \pi \cdot 1^2 \cdot \pi = \frac{\pi^2}{3}$	1	✓	
$\therefore$ Required Volume = $\frac{\pi}{8} \int_0^{\pi} \left(3 + 4 \cos \frac{x}{2} + \cos x \right) dx - \frac{\pi^2}{3}$ from			
(iii)			
$= \frac{\pi}{8} \left[3x + 8 \sin \frac{x}{2} + \sin x \right]_0^{\pi} - \frac{\pi^2}{3}$	1	✓	
$= \frac{\pi}{8} [3\pi + 8 + 0] - \frac{\pi}{8} [0 + 0 + 0] - \frac{\pi^2}{3}$	1		$\rightarrow \frac{3\pi^2}{8} + 1 - \frac{\pi^2}{3}$
$= \frac{8\pi}{8} + \pi^2 \left[\frac{3}{8} - \frac{1}{3} \right] = \pi + \frac{\pi^2}{24} = \pi \left[1 + \frac{\pi}{24} \right]$	1	✓	Few got to the end yet if 4 right steps full mark was awarded.

QUESTION 6

Suggested Solutions	Max Mark	Your Mark	Marker's Comments
(a) Taking logs to base e of each side $(\ln y) = (x) \ln 2007$ Now differentiate both sides with respect to x: $\frac{1}{y} \cdot \frac{dy}{dx} = \ln 2007$ now multiply both sides by $y = (2007)^x$ to get $\frac{dy}{dx} = (2007)^x (\ln 2007)$	1 1		poorly answered some Extension 2 students O.K.
(b) (i) $\hat{APB} = 90^\circ$ since it is the angle in the semi circle diameter AB (ii) $\hat{OBQ} = 90^\circ$ (given) $\therefore \hat{PBQ} = 90^\circ - x^\circ$ But $\hat{APB} = \hat{PBQ} + \hat{PQB}$ (Ext $\angle$ of Δ = sum of interior opp $\angle$'s) $90^\circ = 90^\circ - x^\circ + \hat{PQB}$ $\therefore \hat{PQB} = x^\circ = \hat{TPQ}$ (same angle)	1 1 1		Note: You cannot assume $PT \parallel AB$ I have written in the correct values for each angle in terms of the pronumerals.
(iii) $TP = TB$ (tangents from ext point to a circle are equal) $\hat{TPB} = \hat{BTP} = 90^\circ - x^\circ$ So $\hat{TPQ} = [90^\circ - (90^\circ - x^\circ)] = x^\circ$ (iv) $\frac{AP}{AQ} = \frac{AB \cos 60^\circ}{\left(\frac{AB}{\cos 60^\circ}\right)} = \left(\frac{1}{2} \times \frac{1}{2}\right) = \frac{1}{4} \therefore PQ = 3AP$	1		
(c) (i) $\ddot{x} = \frac{dv}{dt} = \frac{dx}{dt} \cdot \frac{dv}{dx} = v \frac{dv}{dx}$	1		
(ii) Using part (i) $v \frac{dv}{dx} = -e^{-2x}$ and rearranging, $\int v dv = -\int e^{-2x} dx$ Thus $\frac{v^2}{2} = \frac{1}{2} e^{-2x} + K$ Set $x = 0$ and $v = 1$ To get $\frac{1}{2} = \frac{1}{2}(1) + K \Rightarrow K = 0$ thus $v^2 = e^{-2x}$ so $v = e^{-x}$	1 1 1		* Some used $\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \ddot{x}$ this was O.K.
(iii) From part (ii) $\frac{dx}{dt} = e^{-x}$ So again rearranging, $\frac{dx}{e^{-x}} = dt$ and hence $\int e^x dx = \int dt \Rightarrow e^x = t + c$ set $t = 0$ and $x = 0$, (data), $e^0 = 0 + c \therefore c = 1$ ie $e^x = t + 1$ Finally take logs of each side to base e, to get $x = \ln t + 1$	1 1 1		Note: $\frac{dx}{dt} = e^{-x}$ to assist in correct integration

QUESTION 7

$-2 \leq x \leq 2$

Suggested Solutions

Max
Mark

Your
Mark

Marker's Comments

(a) (i) $x \leq |2|$ (ii) $0 \leq y \leq 3\pi$

1+1

so many

disappointing that
students didn't get
these correct.

(b) Note that $mxy - ky = kx + l \therefore mxy - kx = ky + l$ and so

$x(my - k) = ky + l \Rightarrow x = \frac{ky + l}{my - k} = f(y)$

1

→ attempting to make
 x the subject

1

→ correct result

(neither x , or $y = \frac{k}{m}$)

(c) (i) $PQ = BC - 2BP$ and in right angled $\triangle ABP$

$\tan \theta = \frac{2}{BP} \therefore \cot \theta = \frac{BP}{2} \therefore BP = 2 \cot \theta$

Hence $PQ = BC - 2(2 \cot \theta) = 6 - 4 \cot \theta$

1

(ii) By similar reasoning as in (i)

$AP = 2 \operatorname{cosec} \theta$ so $AP + QD = 4 \operatorname{cosec} \theta$

Now focus on time in MINUTES (t_m)

t_m for D to travel $AP + QD = 60 \left(\frac{4 \operatorname{cosec} \theta}{4} \right) = 60 \operatorname{cosec} \theta$

t_m for D to travel $PQ = 60 \left(\frac{6 - 4 \cot \theta}{8} \right) = (45 - 30 \cot \theta)$

Let $T = \text{Total } t_m$, so $T = 60 \operatorname{cosec} \theta + 45 - 30 \cot \theta$
 $= 15(3 + 4 \operatorname{cosec} \theta - 2 \cot \theta)$

(iii) $\therefore \frac{dT}{d\theta} = -60 \operatorname{cosec} \theta \cot \theta + 30 \operatorname{cosec}^2 \theta$

$= 30 \operatorname{cosec}^2 \theta (-2 \cos \theta + 1)$ Note $30 \operatorname{cosec}^2 \theta > 0$

So (in discussing) the sign of $\frac{dT}{d\theta}$ we need focus on sign of $(-2 \cos \theta + 1)$

Now for a Max or a Min $\frac{dT}{d\theta} = 0$ so $-2 \cos \theta + 1 = 0$ or $\cos \theta = \frac{1}{2}$ and

$\theta = \frac{\pi}{3}$ (Note θ is acute)

If $\theta = \frac{\pi}{3}$, $T = 15 \left(\frac{8}{\sqrt{3}} + 3 - \frac{2}{\sqrt{3}} \right) = 45 + 30\sqrt{3}$

$\approx 45 + 51.963 \approx 96.963 \dots$

Hence for $\theta = \frac{\pi}{3}$, T is just slightly below 97 minutes. We have now to show this is

a minimum. Note $\frac{dT}{d\theta}$ takes the same sign as $(1 - 2 \cos \theta)$

Now if $0 \leq \theta < \frac{\pi}{3}$, $\cos \theta > \frac{1}{2} \therefore 1 - 2 \cos \theta < 0$

Now if $\frac{\pi}{3} < \theta \leq \frac{\pi}{2}$, $\therefore \cos \theta < \frac{1}{2}$, $\therefore 1 - 2 \cos \theta > 0$

Hence as θ passes through $\frac{\pi}{3}$, $\frac{dT}{d\theta}$ goes from negative through 0 to positive.

Hence at $\theta = \frac{\pi}{3}$ we get the least value of T .

when $\theta = \frac{\pi}{3}$, $T \doteq 97$ minutes

1

→ correct expression
for length of AP or QD .

1

→ correct expression for
time taken (even if in
hours)

1

→ correct conversion from
time in hours to time
in minutes.

1

→ correct expression
for $\frac{dT}{d\theta}$

1

* many students didn't
use the given results at
beginning of question.

1

→ correct solution
for $\theta \left(\frac{\pi}{3} \right)$

1

→ testing that
 $\theta = \frac{\pi}{3}$ gives a
minimum time

1

→ substituting $\theta = \frac{\pi}{3}$
into expression for T
and obtaining $T \doteq 97$ min.

QUESTION 7 (c) (iii)

Suggested Solutions	Max Mark	Your Mark	Marker's Comments
2nd way of showing that $\theta = \frac{\pi}{3}$ gives a minimum time for Danny's journey. Note that $\frac{1}{30} \frac{dT}{d\theta} = \operatorname{cosec}^2 \theta (1 - 2 \cos \theta)$ $\therefore \frac{1}{30} \frac{d^2T}{d\theta^2} = (1 - 2 \cos \theta) [-2 \operatorname{cosec}^2 \theta \cot \theta] + \operatorname{cosec}^2 \theta (2 \sin \theta) \dots (L)$ RHS of L $= [-2 \operatorname{cosec}^2 \theta \cot \theta + 4 \operatorname{cosec}^2 \theta \cot \theta \cos \theta] + 2 \operatorname{cosec} \theta$ $= 2 \operatorname{cosec}^2 \theta \cot \theta (2 \cos \theta - 1) + 2 \operatorname{cosec} \theta$ $= 2 \operatorname{cosec}^2 \frac{\pi}{3} \cot \frac{\pi}{3} (2 \cos \frac{\pi}{3} - 1) + 2 \operatorname{cosec} \frac{\pi}{3}, \text{ if } \theta = \frac{\pi}{3}.$ $= 0 + \frac{4}{\sqrt{3}}, \text{ as } 2 \cos \frac{\pi}{3} = 1$ $= \frac{4}{\sqrt{3}}$ $\therefore \frac{1}{30} \cdot \frac{d^2T}{d\theta^2} = \frac{4}{\sqrt{3}}$ $\therefore \frac{d^2T}{d\theta^2} = 30 \times \frac{4}{\sqrt{3}} = 40\sqrt{3} > 0$ $\therefore \theta = \frac{\pi}{3}$ gives a minimum time for Danny's journey. Source of question 7(c) University of London GCE Pure Mathematics 405, January 1988. 371/405/420 Question 14	1 1 1 1		