

**SYDNEY GIRLS HIGH SCHOOL**  
**TRIAL HIGHER SCHOOL CERTIFICATE**



**1999**

**MATHEMATICS**

**3 UNIT (Additional)**  
**and**  
**3/4 UNIT (Common)**

Time allowed - 2 hours  
(Plus 5 minutes' reading time)

**DIRECTIONS TO CANDIDATES**

NAME \_\_\_\_\_

- Attempt ALL questions.
  - ALL questions are of equal value.
  - All necessary working should be shown in every question. Marks may be deducted for careless or badly arranged work.
  - Board-approved calculators may be used.
  - Each question attempted should be started on a new sheet. Write on one side of the paper only.
-

## **QUESTION ONE**

a) If  $3 \cot x = 4$ , find the value of

$$\frac{6 \sin x - 4 \cos x}{\operatorname{cosec} x + \sec x} \quad (x \text{ is acute}) \quad [2]$$

b) Evaluate  $\int_0^2 x e^{x^2} dx$  [2]

c) Differentiate  $x^3 \sin^{-1} 4x$  [2]

d) Given  $\log_a b = 0.3$  and  $\log_a c = 0.4$ , find  $\log_a \left( \frac{b}{c} \right) + \log_a ac$  [2]

e) Find the exact value of  $\cos 2x$  if  $\sin x = \sqrt{3} - 1$  [2]

f) A cosine curve has an amplitude of 5 and a period of  $3\pi$ . It has a minimum turning point at  $(0, 5)$ . Find its equation. [2]

## **QUESTION TWO**

a) Write down the domain of the function

$$y = \frac{1}{x^2 + 5x + 6} \quad [1]$$

b) The roots of  $x^3 + 5x^2 + 8x + 2 = 0$  are  $\alpha, \beta$ , and  $\gamma$  [4]

i) Find  $(\alpha + 1) + (\beta + 1) + (\gamma + 1)$

ii) Find  $(\alpha + 1)(\beta + 1)(\gamma + 1)$

c) The half life of a radioactive substance is 24 hrs. How long will it take for only 15% of the substance to remain. (Assume  $M = M_0 e^{-kt}$  and give your answer to the nearest hour) [2]

d) Find the equation of the tangent to the curve  $y = e^{\tan^{-1}x}$  at the point where it cuts the y-axis. [2]

e) The area of the region below the curve  $y = e^{-x}$  and above the x-axis, between  $x = 0.5$  and  $x = 1.5$  is rotated about the x-axis. Find the volume of the solid generated. (Answer correct to 2 decimal places) [3]

### **QUESTION THREE**

a) If  $\frac{dx}{dt} = 5(x-3)$  [3]

i) Show that  $x = 3 + A e^{5t}$  is a solution, where A is a constant.

ii) Find A if  $x = 20$  when  $t = 0$ .

b) [5]  
The points  $P(2ap, ap^2)$ ,  $Q(2aq, aq^2)$  lie on the parabola  $x^2 = 4ay$ .

i) If PQ passes through  $(4a, 0)$  show that  $pq = 2(p+q)$


ii) Hence find the locus of M, the mid point of PQ.

c) Find the size of the acute angle between the lines [2]  
 $y = -x$  and  $\sqrt{3}y = 2x$  (Answer to the nearest minute)


d) Differentiate  $\log_e \left( \frac{3+x}{3-x} \right)$  [2]

### QUESTION FOUR

- a) A right circular cone of vertical angle  $60^\circ$  is being filled with liquid. The depth of liquid in the cone is increasing at a rate of  $4\text{ cm/s}$ . Find the rate of increase of the volume of the liquid in the cone when the depth is  $9\text{ cm}$ . [3]



- b) A projectile is fired at an angle of  $\tan^{-1}(\frac{5}{12})$  to the horizontal with initial velocity  $130\text{ m/s}$ . Using  $g=10\text{ m/s}^2$  [6]
- Derive equations for the horizontal and vertical position of the projectile at time  $t$ .
  - What is the horizontal range of this projectile?
- c)  $AB$  is the diameter of the circle centre  $O$ .  $AD$  is parallel to  $OC$ , and angle  $BAD = 40^\circ$ . Find the size of angle  $DCO$ , giving reasons. [3]



(figure not to scale)

## **QUESTION FIVE**

a)

[9]

i) Find the remainder when  $P(x) = x^3 - (k+1)x^2 + kx + 12$  is divided by  $A(x) = x - 3$

ii) Find  $k$  if  $P(x)$  is divisible by  $A(x)$

iii) Find the zeros of  $P(x)$ , for this value of  $k$

iv) Solve  $P(x) > 0$

b) It is known that  $\log_e x + \sin x = 0$  has one root close to  $x = 0.5$ .  
Use one application of Newton's method to obtain a better approximation of the root correct to 3 decimal places.

[3]

## **QUESTION SIX**

a) Show that  $7^n + 2$  is divisible by 3, for all positive integral  $n$ .

[3]

b) Find the general solution of  $\cos 2x = \sin x$

[3]

c) Find the area bounded by the curve  $y = \frac{1}{\sqrt{25-x^2}}$ , the  $x$  axis and the ordinates at  $x = -2$  and  $x = 2$ .

(Answer correct to 2 decimal places)

[2]

d) Differentiate  $\log_e (\sec x + \tan x)$  and hence find  $\int_0^{\frac{\pi}{4}} \sec x \, dx$ , in simplest exact form.

[4]

## **QUESTION SEVEN**

a)

A Particle moving on a horizontal line has a velocity of  $v$  m / s  
given by  $v^2 = 64 - 4x^2 + 24x$  [5]

i) Prove that the motion is simple harmonic

ii) Find the centre of the motion

iii) Write down the period and amplitude of the motion

iv) Initially the particle is at the centre of the motion and moving to the left. Write down an expression for the displacement as a function of time.

b)

i) Write the expression for  $\sqrt{2} \cos \theta + \sin \theta$  in terms of  $t$ . [4]

(where  $t = \tan \frac{\theta}{2}$ )

ii) Hence or otherwise solve  $\sqrt{2} \cos \theta + \sin \theta = 1$  for  $0^\circ < \theta < 360^\circ$

c) Find  $\int \frac{x dx}{(25 + x^2)^{\frac{3}{2}}}$

using the substitution  $x = 5 \tan \theta$

[3]


1) a)  $3 \cot x = 4$   
 $\cot x = \frac{4}{3}$   
 $\frac{6 \sin x - 4 \cos x}{\sec x + \csc x}$   
 $= \left[ 6\left(\frac{3}{5}\right) - 4\left(\frac{4}{5}\right) \right] \div \left[ \frac{5}{3} + \frac{5}{4} \right]$   
 $= \frac{24}{175}$

b)  $\frac{d}{dx} e^{x^2} = 2x e^{x^2}$   
 $\therefore \int 2x e^{x^2} dx = e^{x^2} + C$   
 $\therefore \int_0^2 x e^{x^2} dx = \frac{1}{2} [e^{x^2}]_0^2$   
 $= \frac{1}{2} (e^4 - 1)$

c)  $y = x^3 \sin^{-1} 4x$   
 put  $u = x^3$ ,  $\frac{du}{dx} = 3x^2$   
 $v = \sin^{-1} 4x$ ,  $\frac{dv}{dx} = \frac{4}{\sqrt{1-16x^2}}$   
 $\frac{dy}{dx} = 3x^2 \sin^{-1} 4x + \frac{4x^3}{\sqrt{1-16x^2}}$

or  $\frac{x^3}{\sqrt{1-16x^2}} + 3x^2 \sin^{-1} 4x$   
 d)  $\log_a \left( \frac{b}{c} \right) + \log_a ac$   
 $= (\log_a b - \log_a c) + (\log_a a + \log_a c)$   
 $= (0.3 - 0.4) + (1 + 0.4)$   
 $= 1.3$

e)  $\sin x = \sqrt{3} - 1$   
 $= \frac{\sqrt{3}-1}{1}$   
 $\cos 2x = \cos^2 x - \sin^2 x$   
 $= 2\sqrt{3} - 3 - (\sqrt{3}-1)^2$   
 $= 2\sqrt{3} - 3 - (4 - 2\sqrt{3})$   
 $= 2\sqrt{3} - 3 - 4 + 2\sqrt{3}$   
 $= 4\sqrt{3} - 7$



$y = 10 - 5 \cos \frac{2x}{3}$   
 or  $y = 10 + 5 \cos \left( \frac{2x}{3} - \pi \right)$   
 $y = 5 \sin \left( \frac{2x}{3} - \frac{\pi}{2} \right) + 10$   
 all real except  $x^2 + 5x + 6 = 0$   
 i.e. all real except  $x = -2, -6$

b)  $P(x) = x^3 + 5x^2 + 8x + 2$   
 $\alpha + \beta + \gamma = -\frac{b}{a} = -5$   
 $\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a} = 8$   
 $\alpha\beta\gamma = -\frac{d}{a} = -2$   
 1)  $(\alpha+1) + (\beta+1) + (\gamma+1)$   
 $= \alpha + \beta + \gamma + 3$   
 $= -5 + 3$   
 $= -2$


ii)  $(\alpha+1)(\beta+1)(\gamma+1)$   
 $= (\alpha+1)(\beta\gamma + \beta + \gamma + 1)$   
 $= (\alpha\beta\gamma) + (\alpha\beta + \alpha\gamma + \beta\gamma) + (\alpha + \beta + \gamma) + 1$   
 $= -2 + 8 - 5 + 1$   
 $= 2$

c)  $M = M_0 e^{-kt}$   
 when  $t = 24$ ,  $m = \frac{M_0}{2}$   
 i.e.  $\frac{M_0}{2} = M_0 e^{-24k}$   
 $0.5 = e^{-24k}$   
 $-24k = \log_e 0.5$   
 $k = \frac{\log_e 0.5}{-24} [\approx 0.0297]$

If 15% remains  
 $m = 0.15 M_0$   
 i.e.  $0.15 = e^{-kt}$   
 $\log_e (0.15) = \log_e e^{-kt}$   
 $-kt = \log (0.15)$   
 $t = \frac{\log (0.15)}{-k}$   
 $= 65 \text{ hrs } 41' 14''$

d)  $y = e^{\tan^{-1} x}$   
 on the y axis  $x = 0$   
 $y = e^{\tan^{-1}(0)}$   
 $= e^0 = 1$  i.e. (0,1)  
 $\frac{dy}{dx} = \frac{1}{1+x^2} \times e^{\tan^{-1} x}$   
 $= \frac{e^{\tan^{-1} x}}{1+x^2}$   
 when  $x = 0$   
 $m = \frac{e^{\tan^{-1}(0)}}{1+0}$   
 $= 1$


$y - y_1 = m(x - x_1)$   
 $y - 1 = 1(x - 0)$   
 $y = x + 1$



$V = \pi \int_{0.5}^{1.5} (e^{-x})^2 dx$   
 $= \pi \int_{0.5}^{1.5} e^{-2x} dx$

$\pi \left[ -\frac{1}{2} e^{-2x} \right]_{0.5}^{1.5}$   
 $= -\frac{\pi}{2} [e^{-3} - e^{-1}]$   
 $= \frac{\pi}{2} \left[ \frac{1}{e} - \frac{1}{e^3} \right] \text{ units}^3$   
 $= 0.50 \text{ (2 d.p.)}$   
 a) i)  $x = 3 + Ae^{5t}$   
 $\frac{dx}{dt} = 5Ae^{5t}$   
 $= 5(x-3) \text{ [since } Ae^{5t} = x-3]$

ii)  $x = 20$  when  $t = 0$   
 $20 = 3 + A$   
 $A = 17$



i) Gradient  $PQ = \frac{ap^2 - aq^2}{2ap - 2aq}$   
 $= \frac{p+q}{2}$

Eqn PQ  
 $y - y_1 = m(x - x_1)$   
 $y - ap^2 = \frac{p+q}{2} (x - 2ap)$

$2y - 2ap^2 = (p+q)(x - 2ap)$   
 subst  $x = 4a$ ,  $y = 0$   
 $-2ap^2 = (p+q)(4a - 2ap)$   
 $-2ap^2 = 4ap - 2ap^2 + 4aq - 2apq$   
 $2apq = 4ap + 4aq$   
 $pq = 2p + 2q$   
 $= 2(p+q)$

ii) Coordinates of M

$$x = \frac{2ap + 2aq}{2}$$

$$y = \frac{ap^2 + aq^2}{2}$$

$$x = a(p+q)$$

$$y = \frac{a}{2}(p^2 + q^2) \quad \textcircled{B}$$

$$\therefore p+q = \frac{x}{a} \quad \textcircled{A}$$

$$\text{Now } y = \frac{a}{2}[p^2 + q^2]$$

$$= \frac{a}{2}[(p+q)^2 - 2pq]$$

$$= \frac{a}{2}\left[\left(\frac{x}{a}\right)^2 - 2pq\right] \quad \text{from A}$$

$$= \frac{a}{2}\left[\left(\frac{x}{a}\right)^2 - 2(2)(2)(p+q)\right] \quad \text{from part i)}$$

$$= \frac{a}{2}\left[\left(\frac{x}{a}\right)^2 - 4(p+q)\right]$$

$$= \frac{a}{2}\left[\frac{x^2}{a^2} - 4\frac{x}{a}\right] \quad \text{from A}$$

$$\text{or } 2ay = x^2 - 4ax$$

$$\text{or } (x-2a)^2 = 2a(y+2a) \quad \textcircled{3}$$

c)  $y = -x, y = \frac{2}{\sqrt{3}}x$   
 $\therefore m_1 = -1, m_2 = \frac{2}{\sqrt{3}}$

$$\tan \theta = \frac{m_1 - m_2}{1 + m_1 m_2}$$

$$= \frac{(-1 - \frac{2}{\sqrt{3}})}{(1 - \frac{2}{\sqrt{3}})}$$

$$\theta = 85^\circ 54' \quad \textcircled{A}$$

d)  $y = \log_e \left(\frac{3+x}{3-x}\right)$   
 $= \log_e(3+x) - \log_e(3-x)$

$$\frac{dy}{dx} = \frac{1}{3+x} - \frac{-1}{3-x}$$

$$= \frac{1}{3+x} + \frac{1}{3-x}$$

$$= \frac{6}{(3-x)(3+x)}$$


Q4 Let depth be  $h$   
 then  $\frac{dh}{dt} = 4 \text{ cm s}^{-1}$

Find  $\frac{dV}{dt}$  when  $h = 9$

$$V = \frac{1}{3}\pi r^2 h$$

$$\frac{dV}{dt} = \frac{1}{3}\pi r^2$$

$$\frac{r}{h} = \tan 30^\circ$$



$$V = \frac{1}{3}\pi r^2 h$$

$$= \frac{1}{3}\pi \left(\frac{h}{3}\right)^2 h$$

$$= \frac{\pi h^3}{9}$$

$$\frac{dV}{dh} = \frac{\pi h^2}{3}$$


$$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt}$$

$$= \frac{\pi h^2}{3} \times 4$$

$$\text{when } h = 9$$

$$\frac{dV}{dt} = \frac{\pi (81)(4)}{3}$$


$$= 108\pi \text{ cm}^3 \text{ s}^{-1}$$



$$\tan \theta = \frac{5}{12}$$

$$\sin \theta = \frac{5}{13}$$

$$\cos \theta = \frac{12}{13}$$



Initially

$$x = 0, y = 0$$

$$\dot{x} = 130 \cos \theta$$

$$= 130 \times \frac{12}{13}$$

$$= 120$$

$$y = 130 \sin \theta$$

$$= 130 \times \frac{5}{13}$$

$$= 50$$

Horizontal Motion

$$\ddot{x} = 0$$

$$\dot{x} = C_1$$

$$\text{when } t = 0, \dot{x} = 120$$

$$\dot{x} = 120$$

$$x = 120t + C_2$$

$$\text{when } t = 0, x = 0$$

$$x = 120t$$

Vertical Motion

$$\ddot{y} = -10$$

$$\dot{y} = -10t + C_1$$

$$\text{when } t = 0, \dot{y} = 50$$

$$\dot{y} = -10t + 50$$

$$y = -5t^2 + 50t + C_2$$

$$\text{when } t = 0, y = 0$$

$$y = -5t^2 + 50t$$

$$\text{at max range } y = 0$$


$$0 = -5t^2 + 50t$$

$$5t = 50 \quad t = 10$$

$$x_{\text{max}} = 120(10)$$

$$= 1200 \text{ metres}$$

c)



$$\angle AOC + 40^\circ = 180^\circ \quad (\text{coint. } \angle \text{s})$$

$$\angle AOC = 140^\circ$$

$$\text{Major } \angle AOC = 360^\circ - 140^\circ$$

$$= 220^\circ \quad (\angle \text{ at } \text{circ})$$

$$\angle ADC = 220^\circ \div 2 \quad (\angle \text{ at cen twice } \angle \text{ at circ})$$

$$= 110^\circ$$

$$\angle OCO = 180^\circ - 110^\circ \quad (\text{coint. } \angle \text{s})$$

$$= 70^\circ$$

$$AD \parallel OC$$

Q5

a) i)  $R = P(3)$

$$= 3^3 - (k+1)9 + 3k + 1$$

$$= 30 - 6k$$

ii) If divisible  $P(3) = 0$

$$0 = 30 - 6k$$

$$k = 5$$

iii)  $P(x) = x^3 - 6x^2 + 5x + 12$

$$x-3 \overline{) x^3 - 6x^2 + 5x + 12}$$

$$\underline{x^3 - 3x^2}$$

$$-3x^2 + 5x$$

$$\underline{-3x^2 + 9x}$$

$$-4x + 12$$


$$\underline{-4x + 12}$$

$$0$$

$$\therefore P(x) = (x-3)(x^2 - 3x - 4)$$

$$= (x-3)(x-4)(x+1)$$

$$\text{Zeros at } x = 3, x = 4$$



$P(x) > 0$  for  $-1 < x < 3$  and  $x > 4$

b)  $y = \ln x + \sin x$

$a_2 = a_1 - \frac{f(a_1)}{f'(a_1)}$

$y = \frac{1}{x} + \cos x$

$a_2 = 0.5 - \frac{(\ln 0.5 + \sin 0.5)}{(-\frac{1}{0.5} + \cos 0.5)}$   
 $= 0.574$  (to 3 d.p.)

### Question 6.

i) Step 1. Verify for  $n=1$   
 i.e.  $7^1 + 2 = 9$  which is  
 divisible by 3

Step 2. a) Assume true for  $n=k$   
 i.e.  $7^k + 2 = 3P$  ( $P$  integer)

b) Prove true for  $n=k+1$

$7^{k+1} + 2 = 7^k \cdot 7 + 2$   
 $= 7(3P-2) + 2$  (from assp)  
 $= 21P - 14 + 2$   
 $= 3(7P-4)$

Since  $P$  is an integer,  $(7P-4)$   
 is an integer and

$7^{k+1} + 2$  is divisible by 3  
 if the assumption is true.

i.e. true for  $n=k+1$  if true  
 for  $n=k$ .

### Missing Solns

Step 3 Since statement is true  
 for  $n=1$ , it is true for  $n=2$ .  
 Since true for  $n=2$ , then  
 true for  $n=3$ , and so  
 on for all positive integers.

b)  $\cos 2x = \sin x$   
 $1 - 2\sin^2 x = \sin x$

$\therefore 2\sin^2 x + \sin x - 1 = 0$

$(2\sin x - 1)(\sin x + 1) = 0$

$\sin x = \frac{1}{2}$  or  $\sin x = -1$

$\therefore x = n\pi + (-1)^n \sin^{-1} \frac{1}{2}$  or

$n\pi + (-1)^n \sin^{-1}(-1)$

i.e.  $x = n\pi + (-1)^n (\frac{\pi}{6})$  or

$n\pi + (-1)^n (-\frac{\pi}{2})$  (3)

c)  $y > 0$  for all  $x$   
 (i.e. does not cut  $x$  axis)

$\therefore A = \int_{-2}^2 \frac{dx}{\sqrt{25-x^2}}$

$= 2 \int_0^2 \frac{dx}{\sqrt{25-x^2}}$  since fn.  
 is even

$= 2 \left[ \sin^{-1} \frac{x}{5} \right]_0^2 = 2 \left( \sin^{-1} \frac{2}{5} \right)$

$\text{Area} \approx 0.82 \text{ u}^2$  (2)

d)  $y = \log(\sec x + \tan x)$

let  $u = \sec x + \tan x$

$= (\cos x)^{-1} + \tan x$

$\frac{du}{dx} = -(\cos x)^{-2} \cdot -\sin x + \sec^2 x$

$= \frac{\sin x}{\cos^2 x} + \sec^2 x$

$= \tan x \cdot \sec x + \sec^2 x$

(see bottom of next page)

Q6 (i)  $V^2 = 64 - 4x^2 + 24x$

Q7 For SHM  $\ddot{x} = -n^2 x$  or  $\ddot{x} = -n^2 x$

Now  $\frac{d}{dx} \left( \frac{1}{2} V^2 \right) = a = \ddot{x}$

$\therefore \frac{1}{2} V^2 = 32 - 2x^2 + 12x$

$\frac{d}{dx} \left( \frac{1}{2} V^2 \right) = -4x + 12$

i.e.  $\ddot{x} = -4(x-3)$  is of form  $\ddot{x} = -n^2 x$

$\therefore$  motion is SHM.

(ii) Centre of motion  $x=0$  i.e.  $x-3=0$  or when  $v=0$   
 $x=3$

(iii) Period  $T = \frac{2\pi}{n}$  where  $n=2$

$\therefore T = \frac{2\pi}{2} = \pi$  Period =  $\pi$  secs

Amplitude: is from centre to end

i.e. from 3 to 8

$\therefore$  amplitude = 5 m.

or complete  $V^2 = n^2(a^2 - x^2)$

(iv)

$x = -5 \sin 2t + 3$

$x = 5 \sin(-2t) + 3$

$x = 5 \cos(2t - \frac{3\pi}{2}) + 3$

Q6 (cont'd)

$\frac{dy}{dx} = \frac{\sec x (\tan x + \sec x)}{\sec x + \tan x}$

$= \sec x$

$\therefore \int_0^{\pi/4} \sec x \, dx = \left[ \ln(\sec x + \tan x) \right]_0^{\pi/4}$

$= \ln \left( \sec \frac{\pi}{4} + \tan \frac{\pi}{4} \right) - \ln(\sec 0 + \tan 0)$

$= \ln(\sqrt{2} + 1) - \ln(1 + 0)$

$= \ln(\sqrt{2} + 1)$  (4)

$$1) \sqrt{2} \cos \theta + \sin \theta = 1$$

$$\therefore \Rightarrow \sqrt{2} \left( \frac{1-t^2}{1+t^2} \right) + \frac{2t}{1+t^2}$$

$$\Rightarrow \frac{\sqrt{2}(1-t^2) + 2t}{1+t^2}$$

$$(ii) \text{ Now } \sqrt{2} \cos \theta + \sin \theta = 1.$$

$$\therefore \frac{\sqrt{2}(1-t^2) + 2t}{1+t^2} = 1$$

$$\sqrt{2}(1-t^2) + 2t = 1+t^2$$

$$\sqrt{2} - \sqrt{2}t^2 + 2t = 1+t^2$$

$$t^2(1+\sqrt{2}) - 2t + (1-\sqrt{2}) = 0.$$

$$\therefore t = \frac{2 \pm \sqrt{4 - 4(1+\sqrt{2})(1-\sqrt{2})}}{2(1+\sqrt{2})}$$

$$t = \frac{2 \pm \sqrt{4 - 4(1-2)}}{2(1+\sqrt{2})}$$

$$\therefore t = \frac{2(1+\sqrt{2})}{2(1+\sqrt{2})} \text{ or } t = \frac{1-\sqrt{2}}{1+\sqrt{2}} \times \frac{1-\sqrt{2}}{1-\sqrt{2}}$$

$$t = 1 \quad t = \frac{2\sqrt{2}-3}{1}$$

$$7) \text{ Soln's } \cos \theta = \frac{1-t^2}{1+t^2} \text{ where } t = \tan \frac{\theta}{2}$$

$$\sin \theta = \frac{2t}{1+t^2}$$

$$0^\circ < \theta < 360^\circ$$

$$\text{when } t=1 \quad \text{when } t = \frac{2\sqrt{2}-3}{1}$$

$$\text{Then } \tan \frac{\theta}{2} = 1 \quad \tan \frac{\theta}{2} = -0.1715$$

$$\therefore \frac{\theta}{2} = \frac{\pi}{4} \quad \theta = -19^\circ 28'$$

$$* \theta = \frac{\pi}{2} \quad \text{But } 0 < \theta < 360^\circ$$

$$\therefore \theta = 360^\circ - 19^\circ 28'$$

$$\begin{cases} \theta = 340^\circ 32' \\ \theta = \frac{\pi}{2} \end{cases}$$

$$C) \int \frac{x \cdot dx}{(25+x^2)^{\frac{3}{2}}}$$

$$I = \frac{\int 25 \tan \theta \sec^3 \theta \cdot d\theta}{125 \sec^3 \theta}$$

$$I = \frac{1}{5} \int \frac{\tan \theta}{\sec \theta} \cdot d\theta$$

$$I = \frac{1}{5} \int \frac{\sin \theta}{\cos \theta} \cdot d\theta$$

$$I = \frac{1}{5} \int \sin \theta \cdot d\theta$$

$$I = -\frac{1}{5} \cos \theta + C$$

$$\therefore I = \frac{-1}{\sqrt{25+x^2}} + C$$

$$x = 5 \tan \theta, \\ dx = 5 \sec^2 \theta \cdot d\theta$$


$$\therefore x \cdot dx = 25 \tan \theta \sec^2 \theta \cdot d\theta$$

$$25+x^2 = 25+25 \tan^2 \theta$$

$$= 25 \sec^2 \theta$$

$$(25+x^2)^{\frac{3}{2}} = 125 \sec^3 \theta$$

But



$$\cos \theta = \frac{5}{\sqrt{25+x^2}}$$