



2010 Trial Examination

FORM VI

MATHEMATICS EXTENSION 1

Wednesday 11th August 2010

General Instructions

- Reading time — 5 minutes
- Writing time — 2 hours
- Write using black or blue pen.
- Board-approved calculators may be used.
- A list of standard integrals is provided at the end of the examination paper.
- All necessary working should be shown in every question.
- Start each question in a new booklet.

Structure of the paper

- Total marks — 84
- All seven questions may be attempted.
- All seven questions are of equal value.

Collection

- Write your candidate number clearly on each booklet.
- Hand in the seven questions in a single well-ordered pile.
- Hand in a booklet for each question, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Place the question paper inside your answer booklet for Question One.

Checklist

- SGS booklets — 7 per boy
- Candidature — 132 boys

Examiner
SJE

QUESTION ONE (12 marks) Use a separate writing booklet.**Marks**

- (a) The polynomial $P(x) = x^4 - x^3 + kx - 4$ has a factor $(x + 1)$. Find the value of k . **1**
- (b) Differentiate $y = \sin(\log_e x)$. **1**
- (c) Find, correct to the nearest degree, the acute angle between the lines **2**
 $x - 3y + 4 = 0$ and $2x + y - 1 = 0$.
- (d) Find the coordinates of the point that divides the interval from $(-3, 4)$ to $(5, -2)$ in the ratio $1 : 3$. **2**
- (e) Find the exact value of $\int_0^2 \frac{4}{4 + x^2} dx$. **2**
- (f) Find $\lim_{x \rightarrow 0} \left(\frac{\sin x \cos x}{x} \right)$. **1**
- (g) Find the term independent of x in the expansion of $\left(x^2 + \frac{2}{x} \right)^{15}$. **3**

QUESTION TWO (12 marks) Use a separate writing booklet.**Marks**

- (a) The equation $x^3 + bx^2 + cx + d = 0$ has roots $2 + \sqrt{3}$, $2 - \sqrt{3}$ and -3 . Use the sum and the product of the roots to find b , c and d . **3**
- (b) Consider the curve $f(x) = \sin^{-1}(2x)$. **2**
- (i) Sketch the curve. **2**
- (ii) Find the gradient of the tangent to the curve at the point where $x = \frac{1}{4}$. **2**
- (c) A particle is undergoing simple harmonic motion subject to the equation $\frac{d^2x}{dt^2} = -6x$. Initially it is at rest at $x = 2$. **1**
- (i) In which direction will it start to move? **1**
- (ii) Show that $v^2 = 6(4 - x^2)$. **2**
- (iii) State the period and the amplitude of the motion. **2**

QUESTION THREE (12 marks) Use a separate writing booklet.**Marks**

(a) Use the substitution $u = x - 1$ to find $\int x(x - 1)^4 dx$. **2**

(b) (i) Express $\cos \theta - \sqrt{3} \sin \theta$ in the form $R \cos(\theta + \alpha)$, where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$. **3**

(ii) Hence, or otherwise, solve $\cos \theta - \sqrt{3} \sin \theta = 1$, for $0 \leq \theta \leq 2\pi$. **2**

(c) Consider the function $f(x) = \frac{x - 1}{x - 2}$.

(i) Show that $f^{-1}(x) = \frac{2x - 1}{x - 1}$. **1**


(ii) Find the vertical and horizontal asymptotes of $f^{-1}(x)$. **2**

(iii) Sketch $f^{-1}(x)$ showing the asymptotes and any x or y intercepts. **1**

(iv) Hence, or otherwise, solve $\frac{2x - 1}{x - 1} \geq 1$. **1**

QUESTION FOUR (12 marks) Use a separate writing booklet.**Marks**

(a)



A square pyramid has altitude PQ of length h and base $ABCD$ of side length x , as shown above. Each face makes an angle α with PQ and each edge makes an angle β with PQ . Assume that it is a right pyramid, so that Q lies in the centre of the base.

(i) Show that $AQ = \frac{x}{\sqrt{2}}$. **1**

(ii) Hence express x in terms of h and β . **1**

(iii) Show that $\sqrt{2} \tan \alpha = \tan \beta$. **1**

Exam continues overleaf ...

QUESTION FOUR (Continued)

(b) Solve for x and y :

3

$$\log_3 x + \log_3 y = 6$$

$$\log_2 x - \log_2 y = 4$$

(c) (i) Write $\cos^2 x$ in terms of $\cos 2x$.


1

(ii) Hence evaluate $\int_0^{\frac{\pi}{3}} \sin 2x \cos^2 x \, dx$.

2

(d)

3



In the diagram above, two circles intersect and P is one of the points of intersection. A straight line is drawn through P cutting the two circles at X and Y . An isosceles triangle XYZ is constructed with $XZ = YZ$. Suppose that XZ cuts the smaller circle at A and YZ cuts the larger circle at B . Suppose also that the line AB cuts the circles at C and D . Let $\angle XYZ$ be θ .

Prove that $\triangle CPD$ is isosceles.

NOTE: You do not have to copy the diagram above. It has been reproduced for you on a tear-off sheet at the end of this paper. Insert this sheet into your answer booklet.

QUESTION FIVE (12 marks) Use a separate writing booklet.**Marks**

- (a) Consider the two points $P(4t, 2t^2)$ and $Q(8t, 8t^2)$ on the parabola $x^2 = 8y$. The tangents at P and Q intersect at R .

(i) Find the equations of the tangents at P and Q .

2

(ii) Find the coordinates of R .

2

(iii) Hence find the locus of R .

1

- (b) By substituting a suitable value for x in the expansion of $(1+x)^n$, show that

2

$$1 + 2 \binom{n}{1} + 4 \binom{n}{2} + \dots + 2^{n-1} \binom{n}{n-1} = 3^n - 2^n.$$

- (c) A forensic scientist is called upon to determine the time of death of a corpse found in a room which is maintained at a constant temperature of 20°C . The temperature T of the corpse was initially measured at midnight to be 29°C . The scientist measured the temperature of the corpse one hour later and it had fallen to 26°C . Assume that the temperature of the body at the time of death was 36.8°C and that the rate of temperature decrease obeys Newton's law of cooling. Let t be the number of hours after midnight.

(i) Show that $T = 20 + 9e^{-kt}$ satisfies the cooling equation $\frac{dT}{dt} = -k(T - 20)$.

1

(ii) Show that $k = \ln \frac{3}{2}$.

2

(iii) Hence estimate the time of death, to the nearest minute.

2

QUESTION SIX (12 marks) Use a separate writing booklet.**Marks**

- (a) Prove by mathematical induction that for all positive integer values of
- n
- ,

3

$$\frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} + \dots + \frac{n}{(n+1)!} = 1 - \frac{1}{(n+1)!}.$$

- (b) A film director must decide whether a stuntman is able to perform a dangerous stunt. The stuntman must leap from a building onto the centre of some erected scaffolding. The centre of the scaffolding is 5 m below his initial position and at a horizontal distance of 14 m. The stuntman jumps at an angle of 30° above the horizontal. Let the stuntman's initial velocity be V , and let x and y be his horizontal and vertical displacements respectively from his initial position. You may assume that the velocity and displacement equations are:

$$\begin{aligned} \dot{x} &= V \cos 30^\circ & \dot{y} &= -10t + V \sin 30^\circ \\ x &= Vt \cos 30^\circ & y &= -5t^2 + Vt \sin 30^\circ \end{aligned}$$

- (i) Show that the Cartesian equation of the stuntman's path is

2


$$y = -\frac{20x^2}{3V^2} + \frac{x}{\sqrt{3}}.$$

- (ii) Hence determine the required initial velocity V so that he lands in the centre of the scaffolding. Write your answer to the nearest m/s.
- (iii) Safety requirements are such that if the impact velocity is greater than 15 m/s, then padding must be placed on the scaffolding. Assuming that the stuntman leaps at the required speed, determine whether or not padding is needed.

2**2**

QUESTION SIX (Continued)

(c)



The diagram above shows two straight lines OX and OY . The points $P_0, P_2, P_4, \dots$ lie on OX , while the points $P_1, P_3, P_5, \dots$ lie on OY .

P_1 is the foot of the perpendicular from P_0 to OY ,
 P_2 is the foot of the perpendicular from P_1 to OX ,
 P_3 is the foot of the perpendicular from P_2 to OY , and so on.

Let $\angle XOY = \theta$, where $0^\circ < \theta < 90^\circ$, let $OP_0 = x$, and let the length of the line joining P_{r-1} to P_r be denoted by d_r , for $r = 1, 2, 3, \dots$

(i) Show that the lengths $d_1, d_2, d_3, \dots$ form a geometric series.


2

(ii) Hence prove that $\sum_{r=1}^{\infty} d_r = x \cot \frac{\theta}{2}$.

1

QUESTION SEVEN (12 marks) Use a separate writing booklet.**Marks**


(a)



The diagram above shows a semi-circle with centre O , radius r and diameter AB . Let P be a point on arc AB . The arc PQ has centre A and Q lies on AB . Let $\angle PAQ = \theta$.

- (i) Show that $AP = 2r \cos \theta$. 1
- (ii) Prove that as θ varies, the arc PQ will have maximum length when $\theta \sin \theta = \cos \theta$. 3
- (iii) Taking $\theta = 1$ as a first approximation to the value of θ that maximises the arc PQ , use one application of Newton's method to find a better approximation. Round your answer to two decimal places. 2

(b)



The diagram above shows a rotating wheel with radius 40 cm and a connecting rod AP with length 120 cm. The pin P slides back and forth along the x -axis as the wheel rotates anticlockwise at a rate of 6 revolutions per second. In each part below you need only address the case where A is in the first quadrant.

- (i) Show that $\alpha = \sin^{-1} \left(\frac{\sin \theta}{3} \right)$. 1
- (ii) Use the chain rule to show that $\frac{d\alpha}{dt} = \frac{12\pi \cos \theta}{\sqrt{9 - \sin^2 \theta}}$ radians per second. 1
- (iii) Show that $x = 40 \left(\cos \theta + \sqrt{9 - \sin^2 \theta} \right)$. 2
- (iv) Find an expression for the velocity of the pin P in terms of θ . 2

END OF EXAMINATION

B L A N K P A G E

The following list of standard integrals may be used:

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$

Question 1

$$\begin{aligned} (a) \quad P(-1) &= (-1)^4 - (-1)^3 + kx - 4 = 0 \\ &= 1 + 1 - k - 4 = 0 \\ &= -k - 2 = 0 \\ &= k = -2 \quad \checkmark \quad (1) \end{aligned}$$

$$\begin{aligned} (b) \quad y &= \sin(\log_e x) \\ \frac{dy}{dx} &= \frac{\cos(\log_e x)}{x} \quad \checkmark \quad (1) \end{aligned}$$

$$\begin{aligned} (c) \quad \tan \theta &= \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \\ &= \left| \frac{\frac{1}{3} - -2}{1 + \frac{1}{3}(-2)} \right| \quad \checkmark \\ &= \left| \frac{\frac{7}{3}}{\frac{1}{3}} \right| \\ &= 7 \\ \therefore \theta &\hat{=} 82^\circ \text{ (nearest degree)} \quad \checkmark \quad (2) \end{aligned}$$

$$\begin{aligned} (d) \quad x &= \frac{1(5) + 3(-3)}{1+3} & y &= \frac{1(-2) + 3(4)}{1+3} \quad \checkmark \\ &= -1 & &= \frac{5}{2} \\ \therefore \text{Coordinates are} & & & (-1, \frac{5}{2}) \quad \checkmark \quad (2) \end{aligned}$$

$$\begin{aligned} (e) \quad \int_0^2 \frac{4}{4+x^2} dx &= 4 \left[\frac{1}{2} \tan^{-1}\left(\frac{x}{2}\right) \right]_0^2 \quad \checkmark \\ &= 2 \left[\tan^{-1}(1) - \tan^{-1} 0 \right] \\ &= 2 \cdot \frac{\pi}{4} \\ &= \frac{\pi}{2} \quad \checkmark \quad (2) \end{aligned}$$

$$\begin{aligned} (f) \quad \lim_{x \rightarrow 0} \left(\frac{\sin x \cos x}{x} \right) &= \lim_{x \rightarrow 0} \left(\frac{\sin x}{x} \right) \times \lim_{x \rightarrow 0} (\cos x) \\ &= 1 \times 1 \\ &= 1 \quad \checkmark \quad (1) \end{aligned}$$

$$\begin{aligned} (g) \quad \left(x^2 + \frac{2}{x} \right)^{15} \quad \text{General Term: } & {}^{15}C_r (x^2)^r \left(\frac{2}{x} \right)^{15-r} \quad \checkmark \\ &= {}^{15}C_r x^{2r} x^{-15+r} 2^{15-r} \\ &= {}^{15}C_r x^{3r-15} 2^{15-r} \end{aligned}$$

So Term independent of x has

$$\begin{aligned} 3r - 15 &= 0 \quad \checkmark \\ r &= 5 \end{aligned}$$

$$\therefore \text{Term is } {}^{15}C_5 2^{10} = 3075072 \quad \checkmark \quad (3)$$

Question 2

(a) Sum of roots: $2 + \sqrt{3} + 2 - \sqrt{3} - 3 = -\frac{b}{a}$

$$1 = -b$$

$$b = -1 \quad \checkmark$$

Product of roots: $(2 + \sqrt{3})(2 - \sqrt{3})(-3) = -\frac{d}{a}$

$$(4 - 3)(-3) = -d$$


$$d = 3 \quad \checkmark$$

Also, $(2 + \sqrt{3})(2 - \sqrt{3}) + (2 - \sqrt{3})(-3) + (-3)(2 + \sqrt{3}) = c$

$$4 - 3 + -6 + 3\sqrt{3} - 6 - 3\sqrt{3} = c$$

$$-11 = c \quad \checkmark \quad \textcircled{3}$$

(b) (i)



(ii) $f'(x) = \frac{2}{\sqrt{1 - (2x)^2}}$ $\checkmark$

$$f'\left(\frac{1}{4}\right) = \frac{2}{\sqrt{1 - \frac{1}{4}}}$$

$$= \frac{2}{\frac{\sqrt{3}}{2}}$$

$$= \frac{4}{\sqrt{3}} \quad \checkmark$$

④

(c) $\frac{d^2x}{dt^2} = -6x$

$t = 0$
 $x = 2$
 $v = 0$

(i) Negative direction towards the origin $\checkmark$

(ii) $\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -6x$

$$\frac{1}{2}v^2 = -3x^2 + c_1$$

$$v^2 = -6x^2 + c_2$$

Now $v = 0, x = 2$

$$0 = -24 + c_2$$

$$\therefore c_2 = 24$$

So $v^2 = -6x^2 + 24$

$$= 6(4 - x^2) \quad \text{as required} \quad \checkmark$$

(iii) $\frac{d^2x}{dt^2} = -n^2x$

$$\therefore n^2 = 6$$

$$n = \sqrt{6}$$

Period = $\frac{2\pi}{n}$

$$= \frac{2\pi}{\sqrt{6}}$$

$$= \frac{\pi\sqrt{6}}{3}$$

Amplitude = 2 $\checkmark$

⑤

12

Question 3

(a) $\int x(x-1)^4 dx$

$u = x-1, x = u+1$
 $du = dx$

$= \int (u+1)u^4 du$ ✓

$= \int u^5 du + \int u^4 du$

$= \frac{u^6}{6} + \frac{u^5}{5} + c$

$= \frac{(x-1)^6}{6} + \frac{(x-1)^5}{5} + c$ ✓ (2)

(b) (i) $\cos \theta - \sqrt{3} \sin \theta = R \cos(\theta + \alpha)$

$= R [\cos \theta \cos \alpha - \sin \theta \sin \alpha]$ ✓

Equating coefficients:

$R \cos \alpha = 1$

$R \sin \alpha = \sqrt{3}$

Squaring and adding

$R^2 = 4$

$\therefore R = 2$ ✓

$\cos \alpha = \frac{1}{2}$ (α acute)

$\therefore \alpha = \frac{\pi}{3}$ ✓

So $\cos \theta - \sqrt{3} \sin \theta = 2 \cos(\theta + \frac{\pi}{3})$

(ii) $2 \cos(\theta + \frac{\pi}{3}) = 1$ $\frac{\pi}{3} \leq \theta + \frac{\pi}{3} \leq \frac{7\pi}{3}$

$\cos(\theta + \frac{\pi}{3}) = \frac{1}{2}$

$\theta + \frac{\pi}{3} = \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}$ ✓

$\therefore \theta = 0, \frac{4\pi}{3}, 2\pi$ ✓ (5)

(c) (i) $f(x) = y = \frac{x-1}{x-2}$

Interchange x and y

$x = \frac{y-1}{y-2}$

$x(y-2) = y-1$

$xy - y = -1 + 2x$ ✓

$y(x-1) = 2x-1$

$y = \frac{2x-1}{x-1}$

$\therefore f^{-1}(x) = \frac{2x-1}{x-1}$ as required.

(ii) Vertical asymptote: $x = 1$ ✓

$f^{-1}(x) = \frac{2 - \frac{1}{x}}{1 - \frac{1}{x}}$

as $x \rightarrow \pm \infty$


$f^{-1}(x) \rightarrow 2$

$\therefore$ Horizontal asymptote: $y = 2$ ✓

(iii) y -intercept ($x=0$)

x -intercept ($y=0$)

$\left. \begin{matrix} y = 1 \\ x = \frac{1}{2} \end{matrix} \right\}$



(iv) $\frac{2x-1}{x-1} \geq 1$ when $x \leq 0$ or $x > 1$ ✓

Question 4

$$(a) (i) AC^2 = x^2 + x^2 \\ = 2x^2$$

$$AC = \sqrt{2}x$$

$$AQ = \frac{\sqrt{2}x}{2}$$

$$= \frac{x}{\sqrt{2}} \text{ as required}$$

$$(ii) \tan \beta = \frac{AQ}{h}$$

$$= \frac{\frac{x}{\sqrt{2}}}{h}$$

$$\therefore x = \sqrt{2}h \tan \beta$$

$$(iii) \tan \alpha = \frac{x}{\frac{x}{2}}$$

$$\therefore 2 \tan \alpha = \frac{x}{h} \text{ and } \sqrt{2} \tan \beta = \frac{x}{h}$$

$$\therefore 2 \tan \alpha = \sqrt{2} \tan \beta$$

$$\sqrt{2} \tan \alpha = \tan \beta$$

$$(b) \log_3 x + \log_3 y = 6 \\ \log_2 x - \log_2 y = 4$$

$$xy = 3^6$$

$$\frac{x}{y} = 2^4$$

Solving simultaneously

$$x^2 = 11664$$

$$x = 108 \quad (x > 0) \checkmark$$

$$y = \frac{3^6}{108}$$

$$= 6^{3/4}$$

$$(c) (i) \cos^2 x = \frac{1}{2} \cos 2x + \frac{1}{2} \checkmark$$

$$(ii) \int_0^{\pi/3} \sin 2x \cos^2 x \, dx = \frac{1}{2} \int_0^{\pi/3} (\sin 2x \cos 2x + \sin 2x) \, dx$$

$$= \frac{1}{2} \left[\frac{\sin^2 2x}{4} \right]_0^{\pi/3} + \frac{1}{2} \left[-\frac{\cos 2x}{2} \right]_0^{\pi/3} \checkmark$$

$$= \frac{1}{8} \left[\left(\frac{\sqrt{3}}{2} \right)^2 - 0 \right] + \frac{1}{4} \left[-\frac{1}{2} - -1 \right]$$

$$= \frac{3}{32} + \frac{3}{8}$$

$$= \frac{15}{32}$$

(c) Alternate Solution

$$\int_0^{\pi/3} \sin 2x \cos^2 x \, dx = \frac{1}{2} \int_0^{\pi/3} \sin 2x (1 + \cos 2x) \, dx$$

$$= \frac{1}{2} \int_0^{\pi/3} \sin 2x \, dx + \frac{1}{2} \int_0^{\pi/3} \sin 2x \cos 2x \, dx$$

$$= -\frac{1}{4} \left[\cos 2x \right]_0^{\pi/3} + \frac{1}{4} \int_0^{\pi/3} \sin 4x \, dx$$

$$= \left[-\frac{\cos 2x}{4} \right]_0^{\pi/3} + \left[-\frac{\cos 4x}{16} \right]_0^{\pi/3}$$

$$= \frac{1}{8} + \frac{1}{4} + \frac{1}{32} + \frac{1}{16}$$

$$= \frac{15}{32}$$

(c) Alternate Solution.

$$\begin{aligned} \int_0^{\pi/3} \sin 2x \cos^2 x \, dx &= 2 \int_0^{\pi/3} \cos^3 x \sin x \, dx \\ &= -2 \left[\frac{\cos^4 x}{4} \right]_0^{\pi/3} \\ &= -2 \left(\frac{1}{64} - \frac{1}{4} \right) = \frac{15}{32} \end{aligned}$$

(d) $\angle YXZ = \theta$ (base angles of isosceles triangle XYZ)
 $\angle PXA = \angle PCA$ (angles subtended at the circumference by arc PA) ✓
 $= \theta$
 $\angle PDC = \theta$ (exterior angle of a cyclic quadrilateral PYBD) ✓

$$\therefore \angle PCO = \angle PDC = \theta$$

$\therefore \triangle CPO$ is isosceles

(3)

12

Question 5

(a) $x^2 = 8y$, $\therefore a = 2$

(i) gradient of tangent at $P(4t, 2t^2)$ is t
 Equation of tangent at P: $y - 2t^2 = t(x - 4t)$
 $y = tx - 2t^2$ ✓

gradient of tangent at $Q(8t, 8t^2)$ is $2t$
 Equation of tangent at Q: $y - 8t^2 = 2t(x - 8t)$
 $y = 2tx - 8t^2$ ✓

(ii) Solving the equations of the tangents simultaneously

$$tx - 2t^2 = 2tx - 8t^2$$

$$6t^2 = tx$$

$$x = 6t$$
 ✓

$$\begin{aligned} y &= t(6t) - 2t^2 \\ &= 4t^2 \end{aligned}$$
 ✓

$\therefore$ Coordinates of R are $(6t, 4t^2)$

(iii) Locus of R: $t = \frac{x}{6}$

$$y = 4\left(\frac{x}{6}\right)^2$$

$$= \frac{4x^2}{36}$$

$$\therefore x^2 = 9y \quad \checkmark \quad (5)$$

$$b) (1+x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots + \binom{n}{n}x^n$$

Substitute $x=2$

$$3^n = 1 + \binom{n}{1}2 + \binom{n}{2}4 + \dots + \binom{n}{n-1}2^{n-1} + \binom{n}{n}2^n$$

$$\text{Now } \binom{n}{n} = 1$$

$$\therefore 3^n = 1 + 2\binom{n}{1} + 4\binom{n}{2} + \dots + 2^{n-1}\binom{n}{n-1} + 2^n$$

$$3^n - 2^n = 1 + 2\binom{n}{1} + 4\binom{n}{2} + \dots + 2^{n-1}\binom{n}{n-1}$$

as required (2)

(c) At midnight: $t=0$, $T=29^\circ\text{C}$

$$t=1, T=26^\circ\text{C}$$

$$(i) T = 20 + 9e^{-kt} \Rightarrow T - 20 = 9e^{-kt}$$

$$\frac{dT}{dt} = -k \cdot 9e^{-kt}$$

$$= -k(T - 20) \quad \text{as required.}$$

$$(ii) T = 20 + 9e^{-kt}$$

$$t=1, T=26$$

$$26 = 20 + 9e^{-k}$$

$$6 = 9e^{-k}$$

$$\frac{2}{3} = e^{-k}$$

$$\ln\left(\frac{2}{3}\right) = -k$$

$$\therefore k = \ln\left(\frac{3}{2}\right)$$

Question 5. (cont.)

(c) (iii) For time of death solve for t when $T=36.8$.

$$36.8 = 20 + 9e^{-kt} \quad (k = \ln \frac{3}{2})$$

$$\frac{16.8}{9} = e^{-kt}$$

$$\ln\left(\frac{16.8}{9}\right) = -kt$$

$$t = -\frac{\ln\left(\frac{16.8}{9}\right)}{\ln\left(\frac{3}{2}\right)}$$

$$\doteq -1.54 \text{ hours}$$

$$\doteq -1 \text{ hour } 32 \text{ mins.}$$

$\therefore$ Time of death is approximately 10:28 p.m.
(1 hour 32 minutes before midnight)

(5)

12

Question 6

a) Step 1: $n=1$

$$\text{LHS} = \frac{1}{2!}$$

$$= \frac{1}{2}$$

$$\text{RHS} = 1 - \frac{1}{2}$$

$$= \frac{1}{2}$$

$$= \text{LHS}$$

Hence, the result is true for $n=1$. ✓

Step 2: Suppose the result is true for $n=k$.

$$\text{i.e. } \frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} + \dots + \frac{k}{(k+1)!} = 1 - \frac{1}{(k+1)!} \quad (*)$$

We need to show that the result is true for $n=k+1$

$$\text{i.e. } \frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} + \dots + \frac{k}{(k+1)!} + \frac{k+1}{(k+2)!} = 1 - \frac{1}{(k+2)!}$$

$$\text{LHS} = 1 - \frac{1}{(k+1)!} + \frac{k+1}{(k+2)!} \quad \text{by } (*)$$

$$= 1 - \frac{(k+2) - (k+1)}{(k+2)!}$$

$$= 1 - \frac{k+2-k-1}{(k+2)!}$$

$$= 1 - \frac{1}{(k+2)!}$$

$$= \text{RHS}$$

Step 3 It follows from Step 1 and Step 2 by mathematical induction that it is true for all positive integers.


Question 6 (cont.)

(b) (i)

$$x = Vt \cos 30^\circ$$

$$= \frac{V\sqrt{3}}{2} t$$

$$\therefore t = \frac{2x}{\sqrt{3}V}$$



$$\begin{aligned} \text{So } y &= -5\left(\frac{2x}{\sqrt{3}V}\right)^2 + \frac{2x}{\sqrt{3}} \cdot \frac{1}{2} \\ &= -\frac{20x^2}{3V^2} + \frac{x}{\sqrt{3}} \quad \text{as required.} \end{aligned}$$

(ii) Sub. $x=14$, $y=-5$ and solve for V .

$$-5 = -\frac{20(14)^2}{3V^2} + \frac{14}{\sqrt{3}}$$

$$-15V^2 = -20 \times 196 + 14\sqrt{3}V^2$$

$$V^2(14\sqrt{3} + 15) = 20 \times 196$$

$$V^2 = \frac{3920}{14\sqrt{3} + 15}$$

$$\div 99.87589\dots$$

$$\therefore V \div 10 \text{ m/s}$$

iii) Impact Velocity = $\sqrt{\dot{x}^2 + \dot{y}^2}$

First, calculate time of flight, $x=14$, $v=10$

$$t = \frac{2(14)}{\sqrt{3}(10)}$$

$$= \frac{14}{5\sqrt{3}} \text{ s}$$

$$\dot{x} = \frac{10\sqrt{3}}{2}$$

$$= 5\sqrt{3}$$

$$\dot{y} = -10 \frac{14}{5\sqrt{3}} + 10 \cdot \frac{1}{2}$$

$$= -\frac{28}{\sqrt{3}} + 5$$

$$= \frac{-28\sqrt{3} + 15}{3}$$

$$v = \sqrt{(5\sqrt{3})^2 + \left(\frac{-28\sqrt{3} + 15}{3}\right)^2}$$

$$= \sqrt{199.99 \dots}$$

$$\approx 14.14 \text{ m/s}$$

$\therefore$ Padding is not required

⑥

Question 6. (cont.)

(c) (i) $\angle OP_0P_1 = 90 - \theta$ (angle sum of $\triangle OP_0P_1$)
 $\angle P_2P_1P_0 = \theta$ (angle sum of $\triangle P_2P_1P_0$)

Now $\cos \theta = \frac{d_2}{d_1}$

Similarly $\angle P_3P_2P_1 = \theta$

and $\cos \theta = \frac{d_3}{d_2}$

Geometric series as $\frac{d_3}{d_2} = \frac{d_2}{d_1} = \cos \theta$

$a = d_1 = r \sin \theta$ (from $\triangle P_1OP_0$)

$r = \cos \theta$

(ii) $\sum_{n=1}^{\infty} dr = \frac{r \sin \theta}{1 - \cos \theta}$

since $|\cos \theta| < 1$

$$= r \frac{2 \sin \theta/2 \cos \theta/2}{2 \sin^2 \theta/2}$$

$$= r \frac{\cos \theta/2}{\sin \theta/2}$$

$$= r \cot \theta/2 \text{ as required}$$

③

Question 7

a) (i) Join PB $\therefore \angle APB = 90^\circ$ (angle in a semi-circle)

$$\text{So } \frac{AP}{2r} = \cos \theta$$

$$AP = 2r \cos \theta$$

(ii) Let l be arc length PQ

$$l = AP \times \theta$$

$$= 2r \cos \theta \times \theta \quad (\theta \text{ in radians})$$

$$\frac{dl}{d\theta} = 2r \cos \theta + 2r\theta(-\sin \theta)$$

$$= 2r (\cos \theta - \theta \sin \theta)$$

Stationary point when $\cos \theta - \theta \sin \theta = 0$
 $\theta \sin \theta = \cos \theta$

Now $\frac{d^2l}{d\theta^2} < 0$ for a maximum

$$\frac{d^2l}{d\theta^2} = 2r(-\sin \theta) - 2r[\sin \theta + \theta \cos \theta]$$

$$= 2r(-2\sin \theta - \cos \theta)$$

$$= -2r(2\sin \theta + \cos \theta) < 0$$

since $0 < \theta < \frac{\pi}{2}$ (θ in a semi-circle)

(iii) let $f(\theta) = \theta \sin \theta - \cos \theta$

$$f'(\theta) = \theta \cos \theta + 2\sin \theta$$

$$\theta_0 = 1$$

Question 7 (cont.)

(a) (iii) cont. $\theta_1 = 1 - \frac{f(\theta_0)}{f'(\theta_0)}$

$$= 1 - \frac{1 \times \sin 1 - \cos 1}{1 \times \cos 1 + 2 \sin 1}$$

$$= \frac{\cos 1 + 2 \sin 1 - \sin 1 + \cos 1}{\cos 1 + 2 \sin 1}$$

$$= \frac{2 \cos 1 + \sin 1}{\cos 1 + 2 \sin 1}$$

$$\doteq 0.86 \quad (2 \text{ dec. pl.})$$

6

(b) (i) Using sine rule:

$$\frac{\sin \alpha}{40} = \frac{\sin \theta}{120}$$

$$\sin \alpha = \frac{\sin \theta}{3}$$

$$\alpha = \sin^{-1}\left(\frac{\sin \theta}{3}\right) \text{ as required}$$

$$0 \leq \alpha < \frac{\pi}{2}$$

(ii) $\frac{d\alpha}{dt} = \frac{d\alpha}{d\theta} \cdot \frac{d\theta}{dt}$

Now $\frac{d\theta}{dt} = 6 \text{ rev/s}$

$$= 12\pi \text{ radians/sec}$$

$$\frac{d\alpha}{d\theta} = \frac{1}{3} \cos \theta \cdot \frac{1}{\sqrt{1 - \frac{\sin^2 \theta}{9}}}$$

$$\begin{aligned}\frac{dx}{d\theta} &= \frac{1}{3} \cos \theta \cdot \frac{3}{\sqrt{9 - \sin^2 \theta}} \\ &= \frac{\cos \theta}{\sqrt{9 - \sin^2 \theta}} \quad \checkmark\end{aligned}$$

$$\begin{aligned}\text{So } \frac{dx}{dt} &= \frac{\cos \theta}{\sqrt{9 - \sin^2 \theta}} \cdot 12\pi \\ &= \frac{12\pi \cos \theta}{\sqrt{9 - \sin^2 \theta}} \quad \text{as required}\end{aligned}$$

(iii) Drop a perpendicular AX from A intersecting OP at X

$$x = OX + XP$$

$$\text{Now } OX = 40 \cos \theta \quad \checkmark$$

$$XP = 120 \cos \alpha$$

$$\begin{aligned}\therefore x &= 40 \cos \theta + 120 \sqrt{1 - \sin^2 \alpha} \\ &= 40 \cos \theta + 120 \sqrt{1 - \frac{\sin^2 \theta}{9}} \quad \text{from (i)} \\ &= 40 \left(\cos \theta + 3 \frac{\sqrt{9 - \sin^2 \theta}}{3} \right) \quad \checkmark \\ &= 40 (\cos \theta + \sqrt{9 - \sin^2 \theta})\end{aligned}$$

Alternate solution to (iii) using cosine rule

$$\cos \theta = \frac{40^2 + x^2 - 120^2}{2(40)x}$$

$$x^2 - 80x \cos \theta + (40^2 - 120^2) = 0$$

Using the quadratic formula

$$x = \frac{80 \cos \theta \pm \sqrt{80^2 \cos^2 \theta - 4(40^2 - 120^2)}}{2}$$

$$= 40 \cos \theta \pm 40 \sqrt{\cos^2 \theta + 8}$$

$$= 40 (\cos \theta + \sqrt{\cos^2 \theta + 8}), \quad x > 0$$

$$= 40 (\cos \theta + \sqrt{9 - \sin^2 \theta}) \quad \text{as required.}$$

$$(iv) \quad \frac{dx}{dt} = \frac{dx}{d\theta} \cdot \frac{d\theta}{dt} \quad \checkmark$$

$$= 40 \left(-\sin \theta + \frac{1}{2} \frac{1}{\sqrt{9 - \sin^2 \theta}} \cdot 2 \sin \theta \cos \theta \right) \times 12\pi$$

$$\begin{aligned}&= 480\pi \left[-\sin \theta - \frac{\cos \theta \sin \theta}{\sqrt{9 - \sin^2 \theta}} \right] \quad \checkmark\end{aligned}$$

(6)