

Name: _____

Class: _____

SYDNEY TECHNICAL HIGH SCHOOL

YEAR 12

HSC ASSESSMENT TASK 3

JUNE 2009

MATHEMATICS Extension 1

Time Allowed: 70 minutes

Instructions:

- Write your name and class at the top of each page.
- All necessary working must be shown. Marks may be deducted for careless or badly arranged work.
- Marks indicated are a guide only and may be varied if necessary.
- Start **each** question on a **new page**.
- Standard integrals can be found on the last page.

Question 1	Question 2	Question 3	Question 4	Question 5	Question 6	Total
/10	/11	/10	/10	/11	/11	/63

Question 1**Marks**

- a) Solve $\log_e x + \log_e(x - 3) = \log_e 4$ 3
- b) Evaluate and leave in exact form
- (i) $\cos^{-1}\left(\frac{-\sqrt{3}}{2}\right)$ 1
- (ii) $\sin\left(2 \cos^{-1}\frac{4}{5}\right)$ 2
- c) Find (i) $\int \frac{2}{9+x^2} dx$ 2
- (ii) $\int \frac{dx}{\sqrt{1-9x^2}}$ 2

Question 2

- a) Consider the function $f(x) = 8x - x^2$
- (i) Sketch $y = f(x)$ clearly showing the x and y intercepts and the vertex. 2
- (ii) State the largest domain containing $x = 8$ for which $f(x)$ has an inverse function, $f^{-1}(x)$ 1
- (iii) State the domain of $f^{-1}(x)$ 1
- (iv) Find the equation of $f^{-1}(x)$ 2
- (v) For what value(s) of x does $f(x) = f^{-1}(x)$? 2
- (vi) Evaluate $f^{-1}(f(-1))$ 1
- b) Sketch the function $y = \sin^{-1}x$ showing clearly the domain and range. 2


Question 3**Marks**

Differentiate

- | | | |
|-------|--|---|
| (i) | $y = e^{3x} \log_e 3x$ | 2 |
| (ii) | $y = \log_e \left(\frac{x-2}{x} \right)$ | 2 |
| (iii) | $y = \log_3 x$ | 2 |
| (iv) | $y = \log_e (\cos x)$ (give answer in simplified form) | 2 |
| (v) | $y = \tan (\log_e x)$ | 2 |

Question 4

- | | | |
|----|--|---|
| a) | (i) Express $\sin^2 x$ in terms of $\cos 2x$ | 1 |
| | (ii) Use this result or otherwise to evaluate $\int_0^{\frac{\pi}{4}} \sin^2 2x dx$ | 3 |
| b) | Differentiate $\sin^2 3x$ and hence evaluate $\int_0^{\frac{\pi}{6}} \sin 3x \cos 3x dx$ | 3 |
| c) | | 3 |



Find the shaded area and leave it in exact form.


Question 5

Marks

- a) Find the general solution for $\sin\theta = \frac{1}{\sqrt{2}}$ 2
- b) Differentiate (i) $y = \tan^{-1}3x$ 2
- (ii) $y = \sin^{-1}(\cos x)$ 2
- c) Sketch $y = 1 - 3\sin 2x$ between $0 \leq x \leq 2\pi$ 3
- d) Sketch $y = \cos^{-1}(\cos x)$ 2

Question 6

- a) A spherical balloon is being inflated so that the surface area is increasing at the rate of $0.3\text{cm}^2\text{s}^{-1}$. When the balloon's radius is 4cm, find the rate of increase. $\left[v = \frac{4}{3}\pi r^3, SA = 4\pi r^2\right]$
- i) In the radius (correct to 3 decimal places) 2
- ii) In the volume (correct to one decimal place) 2
- b) In the diagram, a vertical pole AB, 3 metres high, is placed on top of a 1 metre high support. The pole subtends an angle of θ radians at the point P, which is x metres from the base O of the support.



- (i) Show that $\theta = \tan^{-1} \frac{4}{x} - \tan^{-1} \frac{1}{x}$ 2
- (ii) Show that θ has a stationary point when $x = 2$. 3
- Assume it is a maximum.
- (iii) Deduce that the maximum angle subtended at P is $\theta = \tan^{-1} \frac{3}{4}$. 2

Teacher's Name:

Student's Name/N^o:

2009 Ext. 1 Task 3 Solutions

Question 1

$$a) \log_e x + \log_e (x-3) = \log_e 4$$

$$\log_e x(x-3) = \log_e 4$$

$$x^2 - 3x = 4$$


$$x^2 - 3x - 4 = 0$$

$$(x+1)(x-4) = 0$$

$$x = -1 \text{ or } 4$$

but x must be > 0

$$\therefore x = 4$$

$$ii) \sin(2 \cos^{-1} \frac{4}{5})$$


Let $\cos^{-1} \frac{4}{5} = \theta$

$$\therefore \sin 2\theta = 2 \sin \theta \cos \theta$$

$$= 2 \times \frac{3}{5} \times \frac{4}{5}$$

$$= \frac{24}{25}$$

$$cii) \int \frac{dx}{\sqrt{1-9x^2}}$$

$$\frac{1}{3} \int \frac{dx}{\sqrt{(\frac{1}{3})^2 - x^2}}$$

$$\frac{1}{3} \sin^{-1} \frac{x}{\frac{1}{3}}$$

$$\frac{1}{3} \sin^{-1} 3x + c$$


Student's Name/N^o:

$$b) i) \cos^{-1}(-\frac{\sqrt{3}}{2})$$

Range of $\cos^{-1}$ is

$$0 \leq \theta \leq \pi \quad \therefore \theta$$

is in 2nd quadrant



$$\theta = \pi - \frac{\pi}{6}$$

$$\theta = \frac{5\pi}{6}$$


$$c) i) \int \frac{2}{9+x^2} dx$$

$$\frac{2}{3} \int \frac{1}{9+x^2} dx$$

$$= \frac{2}{3} \tan^{-1} \frac{x}{3} + c$$

Question 2

$$a) i) f(x) = 8x - x^2$$



$$cii) x \geq 4$$

$$civ) y = 8x - x^2$$

$$x = 8y - y^2$$

$$y^2 - 8y + 16 = -x + 16$$

$$(y-4)^2 = -x + 16$$

$$\rightarrow y-4 = \pm \sqrt{16-x}$$

$$y = 4 \pm \sqrt{16-x}$$

$$ciii) x \leq 16$$

Teacher's Name:

Student's Name/N^o:

v) They intersect on

$$y = x$$

$$\therefore \text{Solve } y = x \text{ and } y = 8x - x^2$$

$$x = 8x - x^2$$

$$x^2 = 7x$$

$$x(x-7) = 0$$

$$x = 7 \Rightarrow (7, 7)$$

$$cvi) f^{-1}(f(-1))$$


$$= f^{-1}(-9)$$

$$= 4 + \sqrt{16 - (-9)}$$

$$= 4 + 5$$

$$= 9$$

$$b) y = \sin^{-1} x$$



Question 3

$$ci) y = e^{3x} \log_e 3x$$

$$y' = e^{3x} \times \frac{3}{3x} + 3e^{3x} \log_e 3x$$

$$= \frac{e^{3x}}{x} + 3e^{3x} \log_e 3x$$

$$cii) y = \log_e \left(\frac{x-2}{x} \right)$$

$$= \log_e (x-2) - \log_e x$$

$$y' = \frac{1}{x-2} - \frac{1}{x}$$

$$ciii) y = \log_3 x$$

$$y = \frac{\log_e x}{\log_e 3}$$

$$y' = \frac{1}{x \log_e 3}$$

$$civ) y = \log_e (\cos x)$$

$$y' = \frac{-\sin x}{\cos x}$$

$$y' = -\tan x$$

$$v) y = \tan (\log_e x)$$

$$y' = \sec^2 (\log_e x) \times \frac{1}{x}$$

$$y' = \frac{\sec^2 (\log_e x)}{x}$$

Teacher's Name:

Student's Name/N°:

Question 4

$$a) (i) \sin^2 x = \frac{1 - \cos 2x}{2}$$

$$(ii) \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin^2 2x \, dx$$

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1 - \cos 4x}{2} \, dx$$

$$\frac{1}{2} \left[x - \frac{\sin 4x}{4} \right]_0^{\frac{\pi}{4}}$$

$$\frac{1}{2} \left[\frac{\pi}{4} - 0 - (0) \right]$$

$$= \frac{\pi}{8}$$

$$b) \frac{d}{dx} (\sin^2 3x) = 2 \sin 3x \cos 3x \times 3$$

$$= 6 \sin 3x \cos 3x$$

$$\therefore \frac{1}{6} \frac{d}{dx} (\sin^2 3x) = \sin 3x \cos 3x$$

$$\therefore \frac{1}{6} \int_0^{\frac{\pi}{6}} \frac{d}{dx} (\sin^2 3x) \, dx$$

$$= \int_0^{\frac{\pi}{6}} \sin 3x \cos 3x \, dx$$

$$LHS = \frac{1}{6} \left[\sin^2 3x \right]_0^{\frac{\pi}{6}}$$

$$= \frac{1}{6} \left(\sin^2 \frac{\pi}{2} \right)$$

$$= \frac{1}{6}$$

c) Area shaded = Rectangle - Area to y axis

$$= e^2 \times 2 - \int_0^2 x \, dy$$

$$= 2e^2 - \int_0^2 e^y \, dy$$

$$= 2e^2 - [e^y]_0^2$$

$$= 2e^2 - (e^2 - e^0)$$

$$= e^2 + 1$$

Question 5

$$a) \sin \theta = \frac{1}{\sqrt{2}}$$

$$\theta = n\pi + (-1)^n \sin^{-1} \frac{1}{\sqrt{2}}$$

$$\theta = n\pi + (-1)^n \times \frac{\pi}{4}$$

$$b) (i) y = \tan^{-1} 3x$$

$$y = \tan^{-1} \frac{x}{3}$$

$$y' = \frac{\frac{1}{3}}{x^2 + (\frac{1}{3})^2}$$

$$= \frac{3}{9x^2 + 1}$$

$$(ii) y = \sin^{-1}(\cos x)$$

$$y' = \frac{-\sin x}{\sqrt{1 - \cos^2 x}}$$


$$y' = -1$$

Teacher's Name:

Student's Name/N°:

$$c) y = 1 - 3 \sin 2x$$

Period π , Amplitude 3




$$d) y = \cos^{-1}(\cos x)$$

Domain: All real x

Range: $0 \leq y \leq \pi$

Since $\cos x$ is even, so is y .



Question 6

$$a) \frac{dA}{dt} = 0.3 \quad A = 4\pi r^2 \quad (ii) \frac{dV}{dt} = \frac{dV}{dr} \times \frac{dr}{dt}$$

$$\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$$

$$0.3 = 8\pi r \times \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{0.3}{8\pi \times 4}$$

$$\frac{dr}{dt} = 0.003 \text{ cm/s}$$

$$= 4\pi r^2 \times 0.003$$

$$= 4\pi \times 4^2 \times 0.003$$

$$= 0.6 \text{ cm}^3/\text{s}$$

$$b) (i) \tan \angle BPO = \frac{4}{x} \Rightarrow \angle BPO = \tan^{-1} \frac{4}{x} \quad \theta = \angle BPO - \angle APO$$

$$\tan \angle APO = \frac{1}{x} \Rightarrow \angle APO = \tan^{-1} \frac{1}{x} \therefore \theta = \tan^{-1} \frac{4}{x} - \tan^{-1} \frac{1}{x}$$

$$(ii) \frac{d\theta}{dx} = \frac{-\frac{4}{x^2}}{1 + (\frac{4}{x})^2} - \frac{-\frac{1}{x^2}}{1 + (\frac{1}{x})^2}$$

$$= \frac{-4}{x^2 + 16} + \frac{1}{x^2 + 1}$$

$$= \frac{-4x^2 - 4 + x^2 + 16}{(x^2 + 16)(x^2 + 1)}$$

$$\frac{d\theta}{dx} = 0 \text{ if } -3x^2 + 12 = 0$$

$$x^2 = 4$$

$$x = 2, (x > 0)$$

$$(iii) \text{Sub. } x = 2 \text{ into}$$

$$\theta = \tan^{-1} \frac{4}{2} - \tan^{-1} \frac{1}{2}$$

$$\tan \theta = \tan [\tan^{-1} 2 - \tan^{-1} \frac{1}{2}]$$

$$= \frac{\tan(\tan^{-1} 2) - \tan(\tan^{-1} \frac{1}{2})}{1 + \tan(\tan^{-1} 2) \tan(\tan^{-1} \frac{1}{2})}$$

$$= \frac{2 - \frac{1}{2}}{1 + 2 \times \frac{1}{2}}$$

$$= \frac{1\frac{1}{2}}{2}$$

$$= \frac{1\frac{1}{2}}{2}$$

$$\therefore \theta = \tan^{-1} \frac{3}{4}$$