



SAINT IGNATIUS' COLLEGE

Trial Higher School Certificate

2007

MATHEMATICS EXTENSION 2

12:30pm – 3:35pm

Tuesday 28th August 2007

Directions to Students

• Reading Time : 5 minutes	• Total Marks 120
• Working Time : 3 hours	
• Write using blue or black pen. (sketches in pencil).	• Attempt Question 1 – 8
• Board approved calculators may be used	• All questions are of equal value
• A table of standard integrals is provided at the back of this paper.	
• All necessary working should be shown in every question.	
• Answer each question in the booklets provided and clearly label your name and teacher's name.	


This paper has been prepared independently of the Board of Studies NSW to provide additional exam preparation for students. Although references have been reproduced with permission of the Board of Studies NSW, the publication is in no way connected with or endorsed by the Board of Studies NSW.

Question 1 (15 marks) Use a SEPARATE writing booklet	Marks
(a) (i) Show that $\sin(A+B) + \sin(A-B) = 2 \sin A \cos B$	1
(ii) Hence find the indefinite integral $\int \sin 5x \cos 3x \, dx$	2
(b) Evaluate $\int_0^5 \frac{t \, dt}{\sqrt{t+4}}$	2
(c) Evaluate $\int_{-1}^1 3^x \, dx$ correct to three significant figures.	2
(d) Evaluate $\int_0^{\frac{1}{3}} \frac{dx}{\sqrt{1-9x^2}}$	2
(e) Find $\int \frac{1}{1+\sin x} \, dx$, using the substitution $t = \tan \frac{x}{2}$	3
(f) Use integration by parts to find $\int \frac{\cos^{-1} x}{\sqrt{1+x}} \, dx$	3

Question 2 (15 marks) Use a SEPARATE writing booklet**Marks**

- (a) Prove that $f(x) = \frac{x^3}{\sin x}$ is an even function. 2

- (b) A sketch of $f(x) = -4(x+1)(x-2)$ is shown below.



With the aid of the above diagram, and without the use of calculus, draw a separate half page sketch for each of the following.

- (i) $y = |f(x)|$ 1
- (ii) $y = f(2x)$ 1
- (iii) $y = f(-x)$ 1
- (iv) $y = \frac{1}{f(x)}$ 2
- (v) $y = \sqrt{f(x)}$ 2
- (vi) $y = \log_e f(x)$ 2
- (c) Using calculus, show that $e^{-x} + x - 1 \geq 0$ for real x . 4


Question 3 (15 marks) Use a SEPARATE writing booklet

Marks

- (a) Express the following in the form $(x+iy)$, where x and y are real: 2

$$\frac{i^2-1}{i} + \frac{1}{1+i}$$

- (b)



The Argand diagram above, shows a regular hexagon with vertex A at the point $(0, 3i)$. O is the centre of the hexagon.

- (i) Copy the diagram into your writing booklet.
- (ii) On your diagram show the region within the hexagon in which both the inequalities $|z| \leq 2$ and $-\frac{\pi}{6} \leq \arg z \leq \frac{\pi}{6}$ are satisfied. 2
- (iii) Find in the form $|z - z_1| = R$, the equation of the circle through the points O, B and F . 1
- (iv) Find the complex numbers, in modulus argument form, represented by the points B and C . 2
- (v) The hexagon is rotated anticlockwise about the origin through an angle of $\frac{\pi}{4}$. Express in the form $r(\cos \theta + i \sin \theta)$, where θ is the principal argument, the complex numbers represented by the new positions of B and C . 3

Question 3 continues on page 5

Question 3 (continued)

- (c) If $1 - 2i$ is a root of the equation $z^2 - (3 + i)z + k = 0$,
- (i) explain why the conjugate $1 + 2i$ cannot be a root to the equation 1
 - (ii) show that the other root is $2 + 3i$ 1
 - (iii) find the value of k 1
 - (iv) hence, or otherwise, find the two square roots of $-24 + 10i$. 2

Question 4 (15 marks) Use a SEPARATE writing booklet

Marks

- (a) A polynomial $p(x) = x^n + ax^2 - 2$ has a factor of $(x-1)$ and leaves a remainder of -6 on division by $(x+2)$.
Find:
- (i) the value of a 1
 - (ii) the value of n 1
 - (iii) the zeros of $p(x)$. 2
- (b) Find the values of a and b so that $p(x) = 2x^3 - (2a+1)x^2 + (2+b)x - 1$ has a double root at $x = 1$. 4
- (c) If l, m, n are the roots of the equation $x^3 - 2x + 5 = 0$,
- (i) find the cubic equation whose roots are $2l, 2m, 2n$. 2
 - (ii) find the value of $l^3 + m^3 + n^3$. 2
- (d) Find all the values of k for which the polynomial equation $3x^4 - 4x^3 + k = 0$ has no real roots. 3

Question 5 (15 marks) Use a SEPARATE writing booklet**Marks**

- (a) An ellipse has the equation $\frac{x^2}{25} + \frac{y^2}{16} = 1$. O is the centre of the ellipse and S and S' are the foci.
- (i) Find
- (α) the eccentricity 1
 - (β) the co-ordinates of the foci 1
 - (γ) the equations of the directrices. 1
- (ii) Make a third of a page sketch of the ellipse showing the features found in part (i) 2
- (iii) If $P(x_0, y_0)$ is a point on the ellipse show that $(PS + PS')$ is constant. You may mark point P in quadrant one of the above mentioned diagram. 2
- (iv) Show that the equation of the tangent at $P(x_0, y_0)$ is 2
- $$\frac{xx_0}{25} + \frac{yy_0}{16} = 1.$$
- (v) The tangent at P meets the nearer focus at R . If S is the nearer focus to P ,
- (α) write down the co-ordinates of R . 1
 - (β) find expressions for the gradients of PR and SR in terms of x_0 and y_0 . 2
 - (γ) show that the angle PSR is a right angle. 1
- (b) A hyperbola has its centre at the origin and asymptotes $y = \pm \frac{2}{3}x$. Find its equation. 2

Question 6 (15 marks) Use a SEPARATE writing booklet

Marks

- (a) The region bounded by the curve $y = x(2 - x)$ and the x -axis is rotated about the y -axis. Find the volume of the solid of revolution by taking slices perpendicular to the y -axis. 4
- (b) The region bounded by the curve $y = \ln x$, the line $y = 1$ and the co-ordinate axes is rotated about the x -axis.
- (i) By dividing the resulting solid into cylindrical shells, show that each shell has an approximate volume : $\delta v = 2\pi y e^y \delta y$ where δy is the thickness of the shell. 2
- (ii) Hence calculate the volume of the solid. 2
- (c) The base of a particular solid is the circle $x^2 + y^2 = 8$. Find the volume of the solid if every cross section to the x -axis is an isosceles – right angled triangle with the hypotenuse in the base of the solid. 4
- (d) Show that the straight line $lx + my + n = 0$ is a tangent to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ if $a^2 l^2 - b^2 m^2 = n^2$. 3

Question 7 (15 marks) Use a SEPARATE writing booklet**Marks**


- (a) Mr Kirkpatrick's mathematics class brought him a ride in a Gondola at Queenstown in New Zealand, during a recent trip. When Mr Kirkpatrick's hands were H metres above the Earth's surface, he dropped overboard his packet of beer nuts of mass m kg. The packet of beer nuts encounters air resistance proportional to its velocity v (which is in metres per second), that is the resistive force is equal to mkv .
Taking Mr Kirkpatrick's hands as the origin and downwards displacement as positive:
- (i) Write down an equation of motion representing the passage of the packet of beer nuts. 2
- (ii) Find the terminal velocity, w , of the packet of beer nuts. 1
- (iii) Show that the equation of motion in part (i) can be written as $\ddot{x} = k(w - v)$. 1
- (iv) Show that the displacement, x metres, of the packet of beer nuts from Mr Kirkpatrick's hands is given by: $x = -\frac{v}{k} - \frac{w}{k} \ln\left(\frac{w-v}{w}\right)$. 4
- (v) If the packet reaches the Earth's surface with a velocity of u metres per second, show that $\ln\left(1 - \frac{u}{w}\right) + \frac{u}{w} + \frac{kH}{w} = 0$. 1
- (vi) Consider the moment when the packet of beer nuts has reached 75% of its terminal velocity.
Find:
- (α) the time, t seconds, for this moment to be reached. 3
- (β) the distance fallen at this moment. 3

Question 8 (15 marks) Use a SEPARATE writing booklet

Marks

(a) Draw a neat half page sketch of the graph for $y^2 = x^2(4 - x^2)$. 3

(b)



In the above diagram AB and BC are chords of a circle, and F is on the arc ABC such that arc AF is equal to arc FC . E is the foot of the perpendicular from F to the chord BC . CB is extended to P so that $PE = EC$. (Note that B is inside the triangle APF)


(i) Show that the triangle APF is isosceles. 3

(ii) Show that $AB + BE = EC$. 4

Question 8 continues on page 11

Question 8 (continued)

(c)



A triangle ABC has sides of varying length a , b and c with a fixed interior angle of $BAC = \frac{\pi}{3}$ as shown in the above diagram.

Use the cosine rule to show that:

- | | | |
|------|--|---|
| (i) | $a^2 \geq bc$, and hence, | 3 |
| (ii) | the area of triangle $ABC \leq \frac{a^2\sqrt{3}}{4}$ $ABC \leq \frac{a^2\sqrt{3}}{4}$ | 2 |

End of examination

Solutions



SAINT IGNATIUS' COLLEGE

Trial Higher School Certificate

2007

MATHEMATICS EXTENSION 2

- This is a study resource for HSC preparation
- This document is designed to help students understand the questions
- The solutions should be treated as aids only
- There may be better solutions to some questions.

Question 1

$$(a)(i) LHS = \sin(A+B) + \sin(A-B)$$

$$= \sin A \cos B + \cos A \sin B + \sin A \cos B - \cos A \sin B$$

$$= 2 \sin A \cos B = RHS$$

$$(ii) \int \sin 5x \cos 3x \, dx = \frac{1}{2} \int (\sin 8x + \sin 2x) \, dx$$

$$= \frac{1}{2} \left(-\frac{1}{8} \cos 8x - \frac{1}{2} \cos 2x \right) + C$$

$$= -\frac{1}{16} \cos 8x - \frac{1}{4} \cos 2x + C$$

$$(b) I = \int_0^5 \frac{t \, dt}{\sqrt{t+4}}$$

$$\text{Let } u = t+4$$

$$du = dt$$

$$\text{When } t=5, u=9$$

$$t=0, u=4$$

$$I = \int_4^9 \frac{u-4}{u^{1/2}} \, du$$

$$I = \int_4^9 \left(u^{\frac{1}{2}} - 4u^{-\frac{1}{2}} \right) \, du$$

$$I = \left[\frac{2}{3} u^{\frac{3}{2}} - 2 \times 4 u^{\frac{1}{2}} \right]_4^9$$

$$I = \left[\frac{2}{3} u^{\frac{3}{2}} - 8u^{\frac{1}{2}} \right]_4^9$$

$$I = \left[\left(\frac{2}{3} \times 27 - 24 \right) - \left(\frac{2}{3} \times 8 - 8 \times 2 \right) \right]$$

$$I = 18 - 24 - \frac{16}{3} + 16$$

$$I = 34 - 29\frac{1}{3}$$

$$I = 4\frac{2}{3}$$

$$\begin{aligned} (c) \quad \int_{-1}^1 3^x dx &= \frac{1}{\ln 3} [3^x]_{-1}^1 \\ &= \frac{1}{\ln 3} \left(3 - \frac{1}{3} \right) \\ &= \frac{8}{3 \ln 3} \end{aligned}$$

$$= 2.43 \text{ c.s.f.} \quad \text{Calculator Face } 2.427304604$$

$$(d) \quad \int_0^{\frac{1}{3}} \frac{dx}{\sqrt{1-9x^2}}$$

Note $1-9x^2 = 9\left(\frac{1}{9}-x^2\right)$

$$\sqrt{1-9x^2} = 3\sqrt{\frac{1}{9}-x^2}$$

$$= \frac{1}{3} \int_0^{\frac{1}{3}} \frac{1}{\sqrt{\left(\frac{1}{3}\right)^2 - x^2}} dx$$

$$= \frac{1}{3} \left[\sin^{-1} \frac{x}{\frac{1}{3}} \right]_0^{\frac{1}{3}}$$

$$= \frac{1}{3} \left[\sin^{-1} 3x \right]_0^{\frac{1}{3}}$$

$$= \frac{1}{3} [\sin^{-1} 1 - \sin^{-1} 0]$$

$$= \frac{1}{3} \times \frac{\pi}{2}$$

$$= \frac{\pi}{6}$$

$$(e) \int \frac{1}{1+\sin x} dx$$

$$\text{when } t = \tan \frac{x}{2}$$

$$\frac{dt}{dx} = \frac{1}{2} \sec^2 \frac{x}{2}$$

$$2 dt = (1 + \tan^2 \frac{x}{2}) dx$$

$$\frac{2 dt}{1+t^2} = dx$$

$$I = \int \frac{1}{1 + \frac{2t}{1+t^2}} \cdot \frac{2 dt}{1+t^2}$$

$$I = 2 \int \frac{1+t^2}{(1+t^2+2t)(1+t^2)} dt$$

$$I = 2 \int \frac{1}{(1+t)^2} dt$$

$$I = 2 \int (1+t)^{-2} dt$$

$$I = 2 \frac{(1+t)^{-1}}{-1} + C$$

$$I = -\frac{2}{1+t} + C$$

$$I = -\frac{2}{1+\tan \frac{x}{2}} + C$$

$$(f) \int \frac{\cos^{-1} x}{\sqrt{1+x}} dx$$

$$\text{Let } u = \cos^{-1} x \Rightarrow u' = -\frac{1}{\sqrt{1-x^2}}$$

$$v' = (1+x)^{-\frac{1}{2}} \Rightarrow v = \frac{(x+1)^{\frac{1}{2}}}{\frac{1}{2} \times 1} = 2\sqrt{x+1}$$

$$I = 2\sqrt{x+1} \cos^{-1} x - \int -\frac{1}{\sqrt{1-x^2}} \times 2\sqrt{x+1} dx$$

$$I = 2\sqrt{x+1} \cos^{-1} x + 2 \int \frac{\sqrt{x+1}}{\sqrt{1-x^2}} dx$$

$$I = 2\sqrt{x+1} \cos^{-1} x + 2 \int \frac{1}{\sqrt{1-x}} dx$$

$$I = 2\sqrt{x+1} \cos^{-1} x + 2 \int (1-x)^{-\frac{1}{2}} dx$$

$$I = 2\sqrt{x+1} \cos^{-1} x + 2 \times \frac{(1-x)^{\frac{1}{2}}}{\frac{1}{2} \times -1} + C$$

$$I = 2\sqrt{x+1} \cos^{-1} x - 4\sqrt{1-x} + C$$

Question 2

(a) $f(x) = \frac{x^3}{\sin x}$

$$f(-x) = \frac{(-x)^3}{\sin(-x)}$$

$$= \frac{-x^3}{-\sin x}$$


$$= \frac{x^3}{\sin x}$$

$$= f(x)$$

$\therefore f(x)$ is an even function


(b) $f(x) = -4(x+1)(x-2)$

(i)




$y = |f(x)|$

(ii)




$y = f(2x)$

(iii)




$y = f(-x)$


(iv)



(v)



(vi)



$$(c) \quad e^{-x} + x - 1 \geq 0$$

$$\text{Let } y = e^{-x} + x - 1$$

$$\text{Now } \frac{dy}{dx} = -e^{-x} + 1$$

$$\frac{d^2y}{dx^2} = e^{-x} = \frac{1}{e^x} > 0 \text{ for all } x \text{ in the domain of the reals.}$$

Note: The meaning of a negative index
 $a^{-m} = \frac{1}{a^m}$

Now since $\frac{d^2y}{dx^2} > 0$ we can say that the curve.

$y = e^{-x} + x - 1$ is always concave up.

$$\text{When } \frac{dy}{dx} = 0$$

$$-e^{-x} + 1 = 0$$

$$e^x = 1$$

$$e^x = e^0$$

$$\therefore x = 0$$

$$\text{and when } x = 0, y = e^0 + 0 - 1$$

$y = 0$
 $\therefore$ minimum turning point at $(0, 0)$

Hence $y \geq 0$ for all x hence

$$e^{-x} + x - 1 \geq 0 \text{ for real } x.$$

(C) $(1-2i)$ a root of $z^2 - (3+i)z + k = 0$

(I) Theorem: If polynomial equation co-efficients are real then complex roots exist in conjugate pairs. Hence $(1+2i)$ is not a root.

(II) Sum of roots: $\alpha + \beta = -\frac{[-(3+i)]}{1}$

$$\alpha + \beta = 3+i$$

If $\alpha = 1-2i$

Then $1-2i + \beta = 3+i$

$$\therefore \beta = 2+3i$$

(III) Product of roots: $\alpha\beta = \frac{k}{1}$

$$\alpha\beta = k$$

substitution: $(1-2i)(2+3i) = k$

$$k = 2 + 3i - 4i - 6i^2$$

$$k = 2 + 6 - i$$

$$k = 8 - i$$

(IV) Now considering $z^2 - (3+i)z + (8-i) = 0$

$$z = \frac{(3+i) \pm \sqrt{(3+i)^2 - 4(1)(8-i)}}{2(1)}$$

$$z = \frac{(3+i) \pm \sqrt{9+6i+i^2-32+4i}}{2}$$

$$z = \frac{(3+i) \pm \sqrt{-24+10i}}{2}$$

Note here that
There is an
alternative
method for
finding square
roots

Let one of the square roots be $(a+bi)$

$$\therefore 1-2i = \frac{(3+i) + a+bi}{2}$$

$$2-4i = 3+i+a+bi$$

$$-1-5i = a+bi$$

Equating real and imaginary parts.

$$a = -1, b = -5$$

Similarly it can also be shown that $a=1, b=5$

$\therefore$ The two square roots are $\pm(1+5i)$

Question 4

(a) $p(x) = x^n + ax^2 - 2$

(i) When $x=1$, $p(x) = 0$

$\therefore 0 = (1)^n + a(1)^2 - 2$

$\therefore a = 1$

(ii) When $p(x) = -6$, $x = -2$

$\therefore -6 = (-2)^n + 1 \times (-2)^2 - 2$

$-6 = (-2)^n + 4 - 2$

$-6 = (-2)^n + 2$

$(-2)^n = -8$

$(-2)^n = (-2)^3$

$n = 3$

(iii) $\therefore p(x) = x^3 + x^2 - 2$

Now $p(x) \div (x-1)$

$$\begin{array}{r} x-1 \overline{) \begin{array}{r} x^3 + x^2 + 0x - 2 \\ \underline{x^3 - x^2} \\ 2x^2 + 0x \\ \underline{2x^2 - 2x} \\ 2x - 2 \\ \underline{2x - 2} \\ 0 + 0 \end{array}} \end{array}$$

$\therefore p(x) = (x-1)(x^2 + 2x + 2)$

When $x^2 + 2x + 2 = 0$

$x = \frac{-2 \pm \sqrt{4-8}}{2}$

$x = \frac{-2 \pm \sqrt{4-8}}{2}$

$x = \frac{-2 \pm 2i}{2} = -1 \pm i$

Hence solutions for $p(x) = 0$ are $x = 1$ or $x = -1 \pm i$

$$(b) \quad p(x) = 2x^3 - (2a+1)x^2 + (2+b)x - 1$$

$$p'(x) = 6x^2 - 2(2a+1)x + (2+b)$$

Now There is a double root at $x=1$

$$\therefore p(1) = p'(1) = 0$$

$$\text{So } 2 - (2a+1) \times 1 + (2+b) \times 1 - 1 = 0$$

$$2 - 2a - 1 + 2 + b - 1 = 0$$

$$-2a + b + 2 = 0$$

$$2a - b = 2 \quad \text{--- (1)}$$

$$\text{And } 6 - 2(2a+1) \times 1 + 2 + b = 0$$

$$6 - 4a - 2 + 2 + b = 0$$

$$-4a + b + 6 = 0$$

$$4a - b = 6 \quad \text{--- (2)}$$

$$\text{Solving: } (1) - (2) \quad -2a = -4$$

$$a = 2$$

$$\text{Sub in (1)} \quad 4 - b = 2$$

$$b = 2$$

$$\text{Answer } a=2, b=2$$

$$(b) \quad l, m, n \text{ roots of } x^3 - 2x + 5 = 0$$

$$(1) \quad \text{Let } y = 2x \Rightarrow x = \frac{y}{2}$$

Substitute:

$$\left(\frac{y}{2}\right)^3 - 2\left(\frac{y}{2}\right) + 5 = 0$$

$$\frac{y^3}{8} - y + 5 = 0$$

$$y^3 - 8y + 40 = 0$$

reverting to the variable x :

$$\text{The required equation is } x^3 - 8x + 40 = 0$$

C (ii) If l, m, n are the roots of the equation $x^3 = 2x - 5$ Then:

$$l^3 = 2l - 5 \quad \text{--- ①}$$

$$m^3 = 2m - 5 \quad \text{--- ②}$$

$$n^3 = 2n - 5 \quad \text{--- ③}$$

Add these equations:

$$l^3 + m^3 + n^3 = 2(l + m + n) - (5 \times 3)$$

Now the sum of the roots of the given equation is zero, i.e. $(l + m + n) = 0$

$$\therefore l^3 + m^3 + n^3 = -15$$

(d) considering $3x^4 - 4x^3 + k = 0$.

$$\text{Let } p(x) = 3x^4 - 4x^3 + k.$$

$$p'(x) = 12x^3 - 12x^2 = 12x^2(x - 1)$$

$$p''(x) = 36x^2 - 24x = 12x(3x - 2)$$

For stationary points $p'(x) = 0 \therefore x = 0$ or 1

When $x = 0$ $p(x) = k$


and $p''(x) = 0$

It appears that $(0, k)$ is a horizontal point of inflection

When $x = 1$ $p(x) = k - 1$

and $p''(x) = 12 > 0$

$\therefore (1, k - 1)$ is a minimum turning point.



If $k > 1$ The point $(1, k - 1)$ is above the x-axis and hence $(3x^4 - 4x^3 + k) = 0$ has no real roots.

(a) $\frac{x^2}{25} - \frac{y^2}{16} = 1$ Question 5

$a = 5, b = 4$

(i) $b^2 = a^2(e^2 - 1)$

$16 = 25(e^2 - 1)$

$\frac{16}{25} = e^2 - 1$

$e^2 = \frac{25}{25} + \frac{16}{25}$

$e^2 = \frac{41}{25}$

$e = \frac{\sqrt{41}}{5}, e > 0$

(ii) $S \equiv (ae, 0); S' \equiv (-ae, 0)$

$ae = \frac{5 \times \sqrt{41}}{5} = \sqrt{41}$

$\therefore S = (\sqrt{41}, 0) \text{ and } S' = (-\sqrt{41}, 0)$


(iii) $y = \pm \frac{b}{a}x$

$y = \pm \frac{4}{5}x$

(iv) $x = \pm \frac{a}{e}$

$x = \pm \frac{25}{\sqrt{41}}$

(b)



(c) $\frac{PS}{PM} = e$ and $\frac{PS'}{PM'} = e$ This is from the locus definition of a Hyperbola.

$$PS - PS' = ePM - ePM'$$

$$= e(PM - PM')$$

$$|PS - PS'| = \left| \frac{\sqrt{41}}{5} \left[x_0 - \frac{25}{\sqrt{41}} - \left(x_0 + \frac{25}{\sqrt{41}} \right) \right] \right|$$

$$= \left| \frac{\sqrt{41}}{5} \left(x_0 - \frac{25}{\sqrt{41}} - x_0 - \frac{25}{\sqrt{41}} \right) \right|$$

$$= \left| \frac{\sqrt{41}}{5} \left(-\frac{50}{\sqrt{41}} \right) \right|$$

$$= 10 \text{ i.e. a constant (This is the length of the major axis).}$$

(d) Equation of tangent:

$$\frac{2x}{25} - \frac{4}{8} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{16x}{25y}$$

Gradient of tangent: $m = \frac{16x_0}{25y_0}$

tangent: $y - y_1 = m(x - x_1)$

$$y - y_0 = \frac{16x_0}{25y_0} (x - x_0)$$

$$25y_0y - 25y_0^2 = 16x_0x - 16x_0^2$$

$$16x_0x - 25y_0y = 16x_0^2 - 25y_0^2$$

$$\frac{x_0x}{25} - \frac{y_0y}{16} = \frac{x_0^2}{25} - \frac{y_0^2}{16} \quad \neq$$

$$\text{i.e. } \frac{x_0x}{25} - \frac{y_0y}{16} = 1 \quad \text{--- --- --- } \textcircled{1}$$

Note: (x_0, y_0) satisfies the equation of the hyperbola

Hence $\frac{x_0^2}{25} - \frac{y_0^2}{16} = 1$

(e) (i) Solving ① with $x_R = \frac{25}{\sqrt{41}}$ --- ② for the co-ordinates of R.

$$\text{Sub ② in ①} \quad \frac{x_0}{25} \times \frac{25}{\sqrt{41}} - \frac{y_0}{16} = 1$$

$$\frac{x_0}{\sqrt{41}} - 1 = \frac{y_0}{16}$$

$$y_R = \frac{16}{y_0} \left(\frac{x_0}{\sqrt{41}} - 1 \right)$$

The co-ordinates of R are given by (x_R, y_R)

(ii) Gradients : $m = \frac{y_2 - y_1}{x_2 - x_1}$

$$m_{PS} = \frac{y_0 - 0}{x_0 - \sqrt{41}} = \frac{y_0}{x_0 - \sqrt{41}}$$

$$m_{SR} = \frac{\frac{16}{y_0} \left(\frac{x_0}{\sqrt{41}} - 1 \right) - 0}{\frac{25}{\sqrt{41}} - \sqrt{41}} = \frac{\frac{16}{y_0 \sqrt{41}} (x_0 - \sqrt{41})}{-\frac{16}{\sqrt{41}}}$$

(iii) For perpendicular lines $m_1 m_2 = -1$

$$m_{PS} \times m_{SR} = \frac{y_0}{x_0 - \sqrt{41}} \times \frac{\sqrt{41}}{-16} \times \frac{16}{y_0 \sqrt{41}} (x_0 - \sqrt{41})$$

$$= -1$$



Therefore $PS \perp SR$

Hence $\widehat{PSR} = 90^\circ$ i.e. a right angle.

Question 6

(a)

$$y = 2x - x^2 = x(2-x)$$



$$A(y) = \pi x_2^2 - \pi x_1^2$$

$$\Delta V \doteq \pi(x_2^2 - x_1^2) \Delta y.$$

$$V = \lim_{\Delta y \rightarrow 0} \int_{y=0}^{y=1} \pi x 4 \sqrt{1-y} \Delta y$$

$$V = 4\pi \int_0^1 (1-y)^{\frac{1}{2}} dy$$

$$V = 4\pi \left[\frac{(1-y)^{\frac{3}{2}}}{\frac{3}{2} \times -1} \right]_0^1$$

$$V = 4\pi \times -\frac{2}{3} [0 - 1]$$

$$V = \frac{8\pi}{3} \text{ units}^3$$

Note To find x_1 and x_2

Solve the quadratic equation in x where y is a fixed value

$$x^2 - 2x + y = 0$$

$$x = \frac{2 \pm \sqrt{4-4y}}{2}$$

$$x = 1 \pm \sqrt{1-y}$$

$$x_2 = 1 + \sqrt{1-y}$$


$$x_1 = 1 - \sqrt{1-y}$$

And $x_2^2 - x_1^2 = (x_2 + x_1)(x_2 - x_1)$

$$= 2(2\sqrt{1-y})$$

$$= 4\sqrt{1-y}$$

(b)



$$A(y) = 2\pi y \times x = 2\pi xy$$

$$V \doteq 2\pi xy \Delta y$$

$$\delta V \doteq 2\pi x e^y xy \delta y$$

$$\text{i.e. } \delta V \doteq 2\pi y e^y \delta y$$

$$V = \lim_{\delta x \rightarrow 0} \sum_{y=0}^{y=1} 2\pi y e^y \delta y$$

$$V = 2\pi \int_0^1 y e^y dy$$

Now integration by parts with $u=y \Rightarrow u'=1$
 $v'=e^y \Rightarrow v=e^y$



$$V = 2\pi \left\{ \left[y e^y \right]_0^1 - \int_0^1 e^y dy \right\}$$

$$V = 2\pi \left\{ (e - 0) - \left[e^y \right]_0^1 \right\}$$

$$V = 2\pi (e - e + 1)$$

$$V = 2\pi \text{ units}^3$$

(c)



$$\text{Now } \frac{h}{2y} = \cos 45^\circ = \frac{1}{\sqrt{2}} \Rightarrow h = \frac{2y}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = y\sqrt{2}$$

$$A(x) = \frac{b \times h}{2} = \frac{y\sqrt{2} \times y\sqrt{2}}{2} = y^2$$

$$\Delta V \doteq y^2 \Delta x$$

$$\Delta V \doteq (8 - x^2) \Delta x$$

$$V = \lim_{\Delta x \rightarrow 0} \sum_{x=-2\sqrt{2}}^{x=2\sqrt{2}} (8 - x^2) \Delta x$$

$$V = 2 \int_0^{2\sqrt{2}} (8 - x^2) dx$$

note even function property.

$$V = 2 \left[8x - \frac{x^3}{3} \right]_0^{2\sqrt{2}}$$

$$V = 2 \left[16\sqrt{2} - \frac{16\sqrt{2}}{3} - 0 \right]$$

$$V = \frac{64\sqrt{2}}{3} \text{ units}^3$$

$$(d) \quad lx + my + n = 0 \quad \text{--- --- --- (1)}$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad \text{--- --- --- (2)}$$

$$\text{From (1)} \quad y = -\frac{lx+n}{m}$$

Sub in (2)

$$\frac{x^2}{a^2} + \frac{(lx+n)^2}{b^2m^2} = 1$$

$$b^2m^2x^2 + a^2(l^2x^2 + 2lnx + n^2) = a^2b^2m^2$$

$$b^2m^2x^2 + a^2l^2x^2 + 2a^2lnx + a^2n^2 - a^2b^2m^2 = 0$$

$$(b^2m^2 + a^2l^2)x^2 + 2a^2lnx + a^2(n^2 - b^2m^2) = 0$$

In this quadratic equation $\Delta = 0$ for tangency

$$(2a^2ln)^2 - 4(b^2m^2 + a^2l^2) \times a^2(n^2 - b^2m^2) = 0$$

$$\div 4a^2 : a^2l^2n^2 - (b^2m^2 + a^2l^2)(n^2 - b^2m^2) = 0$$

$$a^2l^2n^2 - (b^2m^2n^2 - b^4m^4 + a^2l^2n^2 - a^2l^2b^2m^2) = 0$$

$$a^2l^2n^2 - b^2m^2n^2 + b^4m^4 - a^2l^2n^2 + a^2l^2b^2m^2 = 0$$

$$-b^2m^2n^2 + b^4m^4 + a^2l^2b^2m^2 = 0$$


$$-m^2n^2 + b^2m^4 + a^2l^2m^2 = 0$$

$$-n^2 + b^2m^2 + a^2l^2 = 0$$

$$\text{ie } n^2 = a^2l^2 + b^2m^2$$

Question 7

(a)



$$F = ma$$

$$m\ddot{x} = mg - mkv$$

$$\ddot{x} = g - kv \quad \text{--- (1)}$$

(b) For terminal velocity w let $\ddot{x} = 0$

$$\therefore g - kw = 0$$

$$w = \frac{g}{k}$$

(c)

Now from (1)

$$\ddot{x} = k \left(\frac{g}{k} - v \right)$$

$$\ddot{x} = k(w - v) \quad \text{--- (2)}$$

(d) In (2) let $\ddot{x} = v \frac{dv}{dx}$

$$v \frac{dv}{dx} = k(w - v)$$

$$\frac{dv}{dx} = k \left(\frac{w - v}{v} \right)$$

$$\frac{dx}{dv} = \frac{1}{k} \times \left(\frac{v}{w - v} \right)$$

$$\frac{dx}{dv} = \frac{1}{k} \left[-1 - \frac{w}{v - w} \right]$$

Note

$$\begin{aligned} \frac{v}{w - v} &= - \left(\frac{v}{v - w} \right) \\ &= - \left(\frac{v - w + w}{v - w} \right) \\ &= - \left(1 + \frac{w}{v - w} \right) \\ &= -1 - \frac{w}{v - w} \end{aligned}$$

(f) Now replacing $\ddot{x} = \frac{dv}{dt}$ in ②

$$(1) \quad \frac{dv}{dt} = k(w-v)$$

$$\frac{dt}{dv} = \frac{1}{k} \times \left(\frac{1}{w-v} \right) \quad \text{Note Several students used substitution here: let } u = w-v \text{ etc}$$

$$t = \frac{1}{k} \ln |w-v| + c$$

$$t = -\frac{1}{k} \ln |w-v| + c$$

Now when $t = 0$, $v = 0$

$$0 = -\frac{1}{k} \ln |w| + c$$

$$c = \frac{1}{k} \ln |w|$$

$$\therefore t = \frac{1}{k} \ln |w| - \frac{1}{k} \ln |w-v|$$

$$t = \frac{1}{k} \ln \left| \frac{w}{w-v} \right| \quad \rightarrow \quad \text{--- (4)}$$

Now 75% of terminal velocity means $v = 0.75w$.

$$t = \frac{1}{k} \ln \left| \frac{w}{w-0.75w} \right|$$

$$t = \frac{1}{k} \ln \frac{1}{0.25}$$

$$t = \frac{1}{k} \ln \frac{1}{\frac{1}{4}}$$

$$t = \frac{1}{k} \ln 4 \text{ seconds.}$$

ie the time to reach 75% of terminal velocity

Note
 $0.25 = \frac{1}{4}$

$$x = \frac{1}{k} \left[-v - w \ln|v-w| \right] + C = -\frac{v}{k} - \frac{w}{k} \ln|v-w| + C$$

Now $x=0$ when $v=0$

$$0 = -\frac{w}{k} \ln|-w| + C.$$

$$C = \frac{w}{k} \ln|-w|$$

$$\therefore x = -\frac{v}{k} - \frac{w}{k} \ln|v-w| + \frac{w}{k} \ln|-w|$$

$$x = -\frac{v}{k} - \frac{w}{k} \left[\ln|v-w| - \ln|-w| \right]$$

$$x = -\frac{v}{k} - \frac{w}{k} \ln \left| \frac{v-w}{-w} \right|$$

$$x = -\frac{v}{k} - \frac{w}{k} \ln \left| \frac{w-v}{w} \right| \quad \text{--- (3)}$$

(e) Now when $x=H$, $v=w$ sub in

$$H = -\frac{u}{k} - \frac{w}{k} \ln \left| \frac{w-u}{w} \right|$$

$$kH = -u - w \ln \left| 1 - \frac{u}{w} \right|$$

$$w \ln \left| 1 - \frac{u}{w} \right| + u + kH = 0$$

$$\ln \left| 1 - \frac{u}{w} \right| + \frac{u}{w} + \frac{kH}{w} = 0$$

(ii) Back to equation (4)

$$t = \frac{1}{k} \ln \left| \frac{w}{w-v} \right|$$

$$kt = -\ln \left| \frac{w-v}{w} \right|$$

$$\ln \left| \frac{w-v}{w} \right| = -kt$$

$$\frac{w-v}{w} = e^{-kt}$$

$$w-v = we^{-kt}$$

$$v = w - we^{-kt}$$

$$\frac{dx}{dt} = w - we^{-kt}$$

$$x = wt + \frac{w}{k} e^{-kt} + c$$

Now when $x = 0$, $t = 0$

$$0 = 0 + \frac{w}{k} e^0 + c$$

$$c = -\frac{w}{k}$$

$$x = wt + \frac{w}{k} (e^{-kt} - 1)$$

When $t = \frac{1}{k} \ln 4$

$$x = \frac{w}{k} \ln 4 + \frac{w}{k} \left(e^{-k \times \frac{1}{k} \ln 4} - 1 \right)$$

$$x = \frac{w}{k} \left[\ln 4 + e^{-\ln 4} - 1 \right]$$

$$x = \frac{w}{k} \left(\ln 4 + e^{\ln 4^{-1}} - 1 \right)$$

$$x = \frac{w}{k} \left(\ln 4 + 4^{-1} - 1 \right)$$

$$x = \frac{w}{k} \left(\ln 4 - \frac{3}{4} \right) \text{ metres.}$$

(ii) Alternative approach:

Substitute $v = 0.75w$
in equation (3) part (i)

$$x = -\frac{0.75w}{k} - \frac{w}{k} \ln \left(\frac{w-0.75w}{w} \right)$$

$$x = -\frac{w}{k} \left[0.75 + \ln \left(\frac{w(1-0.75)}{w} \right) \right]$$

$$x = -\frac{w}{k} \left[\frac{3}{4} + \ln 0.25 \right]$$

$$x = -\frac{w}{k} \left(\frac{3}{4} + \ln \frac{1}{4} \right)$$

$$x = -\frac{w}{k} \left(\frac{3}{4} + \ln 4^{-1} \right)$$

$$x = -\frac{w}{k} \left(\frac{3}{4} - \ln 4 \right)$$


$$x = \frac{w}{k} \left(\ln 4 - \frac{3}{4} \right) \text{ metres.}$$

This is the distance fallen at 75% of the terminal velocity


Question 8

(a) $y^2 = x^2(x^2 - 4)$

Dotted Graph: $g(x) = x^2(x^2 - 4)$



(b)



(i) arc AF = arc FC {given} $\therefore AF = FC$ --- ①

But in $\triangle PFC$, EF perpendicularly bisects base PC.

$\therefore \triangle PFC$ is isosceles {converse of the Theorem for an isosceles $\triangle$ which states that the altitude is the perpendicular bisector of the base}

$\therefore PF = FC$ --- ②

From ① and ② $AF = PF$ $\therefore \triangle APF$ is isosceles.

(ii) Now $\alpha_1 = \alpha_2$ {Angles in the same segment Theorem - BF subtends equal angles at A and C}

And $\alpha_2 = \alpha_3$ {base angles of isosceles $\triangle PFC$ }

Now since $\triangle APF$ is isosceles


$\angle FPA = \angle FAP = \beta$, say

$\therefore \angle BPA = \angle BAP = \beta - \alpha$

$\therefore \triangle PBA$ is isosceles and $\therefore AB = BP$ --- ③

Hence, $EC = EP = EB + BP = EB + AB$ (from ③) $= AB + BE$.

(c)



(i)

using The cosine rule :

$$a^2 = b^2 + c^2 - 2bc \cos \frac{\pi}{3}$$

$$\therefore a^2 = b^2 + c^2 - 2bc \left(\frac{1}{2}\right)$$

$$\therefore a^2 = b^2 + c^2 - bc$$

Now $a^2 - bc = b^2 + c^2 - 2bc$ (subtracting bc from both sides)

$$\therefore a^2 - bc = (b - c)^2$$

Now $(b - c)^2$ is a perfect square

$$\therefore a^2 - bc \geq 0 \quad (\text{note equality if } b = c)$$

$$\therefore a^2 \geq bc \quad \text{--- ①}$$

(iii) The area of $\triangle ABC = \frac{1}{2} bc \sin \frac{\pi}{3}$

$$\text{Area} = \frac{1}{2} bc \times \frac{\sqrt{3}}{2}$$

$$\text{Area} = \frac{\sqrt{3}}{4} bc$$

$$\text{Area} \leq \frac{\sqrt{3}}{4} a^2 \quad (\text{using part (i) ①})$$