

Question 1 (15 Marks)**Marks**

(a) Sketch the curve $(x + 1)(y - 2) = 1$

2

(b) Write the function in part (a) in the form $y = f(x)$. Hence, or otherwise, sketch the curve

(i) $y = f(x - 2)$

1

(ii) $y = \sqrt{f(x)}$

2

(c) Evaluate

(i) $\int_2^3 \frac{4}{(x-1)(4-x)} dx$

3

(ii) $\int_0^1 \frac{dx}{9-4x^2}$

2

(d) (i) The point P moves so that the sum of its distances from the points $(-2, 0)$ and $(2, 0)$ is 6 units. Prove that the equation of the locus of P is $\frac{x^2}{9} + \frac{y^2}{5} = 1$.

4

(ii) Find the eccentricity of this locus.

1**Question 2 (15 Marks) START A NEW PAGE****Marks**

(a) The hyperbola H is given by the equation $\frac{x^2}{16} - \frac{y^2}{25} = 1$.

(i) Write down the equations of the asymptotes.

2

(ii) P is an arbitrary point with coordinates $(4\sec\theta, 5\tan\theta)$. Show that P lies on H .

2

(iii) Prove that the tangent to H at P has equation $\frac{x\sec\theta}{4} - \frac{y\tan\theta}{5} = 1$.

3

(iv) This tangent cuts the asymptotes in L and M . Prove that $LP = PM$.

3

Question 2 Continued**Marks**

(b) Let n be a positive integer and $I_n = \int_1^2 (\ln x)^n dx$.

(i) Prove that $I_n = 2(\ln 2)^n - nI_{n-1}$.

2

(ii) Hence, evaluate $\int_1^2 (\ln x)^3 dx$.

3**Question 3 (15 Marks) START A NEW PAGE****Marks**

(a) Find $\int \frac{x+1}{x^2+1} dx$.

2

(b) On the attached sheet, you are given the curve of $y = f(x)$.
Sketch neatly on separate diagrams

(i) $y = |f(x)|$

1

(ii) $y = \frac{1}{f(x)}$

2

(c) (i) Evaluate $\int_{-\pi}^{\pi} x \cos x dx$.

2

(ii) Evaluate $\int_0^{\frac{\pi}{2}} \cos^5 \theta d\theta$.

3

(d) F is the point $(4, 0)$ and d is the line $x = 1$. M is the foot of the perpendicular from a variable point P to d , and P moves so that $FP = 2PM$.

(i) Derive the equation of the locus of P .

3

(ii) Find the acute angle between the asymptotes to the nearest degree.

2

Question 4 (15 Marks) START A NEW PAGE**Marks**

(a) Use the substitution $t = \tan \frac{x}{2}$ to prove that $\int_0^{\frac{\pi}{2}} \frac{dx}{2 + \sin x} = \frac{\pi}{3\sqrt{3}}$.

4

(b) (i) Show that $\int_0^{2a} f(x) dx = \int_0^a \{f(x) + f(2a - x)\} dx$.

3

(ii) Hence, or otherwise evaluate $\int_0^{\pi} \frac{x dx}{2 + \sin x}$.

3

(c) (i) Evaluate $\int_0^{\frac{\pi}{4}} \sec \theta d\theta$.

2

(ii) Hence, show that $\int_0^{\frac{\pi}{4}} \sec^3 \theta d\theta = \frac{1}{2} [\sqrt{2} + \ln(\sqrt{2} + 1)]$

3


~ END OF ASSESSMENT~

Question 3 Part (b)
HAND IN WITH YOUR ANSWERS

Sketch neatly on separate diagrams


(i) $y = |f(x)|$

1



(ii) $y = \frac{1}{f(x)}$


2



~SOLUTIONS Year 12 Extension 2 Term 1 Assessment 2007~


Question 1

(a)




Asymptotes $x = -1$ & $y = 2 \rightarrow$ 1 mark
Shape $\rightarrow$ 1 Mark

(b) (i)



$y = f(x - 2)$ is a horizontal shift of $y = f(x)$ 2 units to the right. $\rightarrow$ 1 Mark
(Asymptotes at $y = 2$ & $x = 1$)

(ii)



$$y = \sqrt{f(x)}$$

Asymptote at $x = -1$ & $y = \sqrt{2}$, $\rightarrow$ ½ Mark

y - intercept = $\sqrt{3}$ &

x - intercept = $-1.5 \rightarrow$ 1 Mark

No graph below x - axis $\rightarrow$ ½ Mark

Shape is relatively similar to the original function $\therefore$ no marks awarded

Q1 (c)

$$(i) \int_2^3 \frac{4}{(x-1)(4-x)} dx$$

$$= \frac{4}{3} \int_2^3 \left(\frac{1}{x-1} + \frac{1}{4-x} \right) dx$$

By Partial Fractions

$$= \frac{4}{3} [\ln(x-1) - \ln(4-x)]_2^3$$

$$= \frac{4}{3} [(\ln 2 - \ln 1) - (\ln 1 - \ln 2)]$$

$$= \frac{4}{3} [2 \ln 2] = \frac{8 \ln 2}{3}$$

Correct values of A and B → 1 Mark

Correct integration → 1 Mark

Correct answer → 1 Mark

$$(ii) \int_0^1 \frac{dx}{9-4x^2}$$

$$= \frac{1}{6} \int_0^1 \left(\frac{1}{3+2x} + \frac{1}{3-2x} \right) dx$$

$$= \frac{1}{6} \left[\frac{\ln(3+2x)}{2} - \frac{\ln(3-2x)}{2} \right]_0^1$$

$$= \frac{1}{12} [(\ln 5 - \ln 3) - (\ln 1 - \ln 3)]$$

$$= \frac{\ln 5}{12}$$

This is not a inverse Trigonometry Question!
No marks awarded if used incorrect method.

Correct use of partial fractions & integration → 1 Mark

Correct evaluation → 1 Mark

(d) (i) P is (x, y) then by data

$$\sqrt{(x+2)^2 + y^2} + \sqrt{(x-2)^2 + y^2} = 6$$

$$\text{i.e. } 12\sqrt{x^2 + 4x + 4 + y^2} = 36 - 8x$$

$$\therefore 9(x^2 - 4x + 4 + y^2) = 81 - 36x + 4x^2$$

$$\therefore 5x^2 + 9y^2 = 45 \Rightarrow \frac{x^2}{9} + \frac{y^2}{5} = 1$$

→ 1 Mark

→ 1 Mark

→ 1 Mark

→ 1 Mark

(ii) Using $b^2 = a^2(1 - e^2)$ since locus is an ellipse where $a = 3$ and $b = \sqrt{5}$, we get

$$e = \frac{2}{3}.$$

→ 1 Mark

Question 2

(a) (i) Eqn. of asymptotes $\frac{x^2}{16} - \frac{y^2}{25} = 0$

$$\therefore y = \pm \frac{5x}{4}$$

(ii) If P is $(4\sec\theta, 5\tan\theta)$ then

$$\begin{aligned} \text{LHS} &= \frac{x^2}{16} - \frac{y^2}{25} \\ &= \frac{16\sec^2\theta}{16} - \frac{25\tan^2\theta}{25} \\ &= \tan^2\theta + 1 - \tan^2\theta \\ &= 1 = \text{RHS} \end{aligned}$$

$\therefore P$ lies on hyperbola H .

(iii) $\frac{dy}{dx} = \frac{5\sec\theta}{4\tan\theta}$ at P .

$\therefore$ eqn. of tangent at P is

$$y - 5\tan\theta = \frac{5\sec\theta}{4\tan\theta}(x - 4\sec\theta)$$

$$4\tan\theta y - 5x\sec\theta = 20(\tan^2\theta - \sec^2\theta)$$

$$\therefore \frac{x\sec\theta}{4} - \frac{y\tan\theta}{5} = 1$$

(iv) tangent cuts $y = \frac{5x}{4}$ when

$$\frac{x\sec\theta}{4} - \frac{5x}{4} \cdot \frac{\tan\theta}{5} = 1$$

$$\therefore x = \frac{4}{\sec\theta - \tan\theta} \text{ and } y = \frac{5}{\sec\theta - \tan\theta}.$$

$$\therefore L \left[\frac{4}{\sec\theta - \tan\theta}, \frac{5}{\sec\theta - \tan\theta} \right]$$

Tangent cuts $y = -\frac{5x}{4}$ at

$$M \left[\frac{4}{\sec\theta + \tan\theta}, \frac{-5}{\sec\theta + \tan\theta} \right]$$

If P is the midpoint of LM then $LP = PM$.

→ 1 Mark

→ 1 Mark

→ 1 Mark

Showing LHS = RHS → 1 Mark

Correct differential → 1 Mark

Correct eqn. of tangent → 1 Mark

Correct rearrangement to get required answer → 1 Mark

Correct coordinates of L → ½ Mark

Coordinates of M → ½ Mark

$\therefore x$- coordinate of midpoint of LM $= \frac{1}{2} \left[\frac{4}{\sec \theta - \tan \theta} + \frac{4}{\sec \theta + \tan \theta} \right]$ $= \frac{1}{2} \left[\frac{8 \sec \theta}{\sec^2 \theta - \tan^2 \theta} \right]$ $= 4 \sec \theta.$ y- coordinate of midpoint of LM $= \frac{1}{2} \left[\frac{5}{\sec \theta - \tan \theta} + \frac{-5}{\sec \theta + \tan \theta} \right]$ $= 5 \tan \theta.$ $\therefore P$ is the midpoint of LM. $\therefore LP = PM$ (b) (i) $I_n = \int_1^2 (\ln x)^n dx$ $= \left[x (\ln x)^n \right]_1^2 - \int_1^2 \frac{x \cdot n (\ln x)^{n-1}}{x} dx$ $= 2(\ln 2)^n - n I_{n-1}$ (ii) $I_3 = 2(\ln 2)^3 - 3 I_2$ $= 2(\ln 2)^3 - 3[2(\ln 2)^2 - 2 I_1]$ Now $I_1 = \int_1^2 \ln x dx$ $= \left[x \ln x - x \right]_1^2$ $= 2 \ln 2 - 2 - 0 + 1$ $= 2 \ln 2 - 1$ $\therefore I_3 = 2(\ln 2)^3 - 6(\ln 2)^2 + 12 \ln 2 - 6.$	Correctly finding x and y coordinates of midpoint Or using distance formula & conclusion → 2 Marks → 2 Marks → 1 Mark → 1 Mark → 1 Mark
---	--

Question 3

$$(a) \int \frac{x+1}{x^2+1} dx = \int \left(\frac{x}{x^2+1} + \frac{1}{x^2+1} \right) dx$$

$$= \frac{1}{2} \ln(x^2+1) + \tan^{-1} x + c$$

(b) See next page

(c) (i) Let $f(x) = x \cos x$ then $f(-x) = -x \cos(-x)$
 $= -f(x) \therefore$ it's an odd function.

$$\therefore \int_{-\pi}^{\pi} x \cos x dx = 0$$

$$(ii) \int_0^{\frac{\pi}{2}} \cos^5 \theta d\theta = \int_0^{\frac{\pi}{2}} \cos \theta (\cos^4 \theta) d\theta$$

$$= \int_0^{\frac{\pi}{2}} \cos \theta (1 - \sin^2 \theta)^2 d\theta$$

$$= \int_0^{\frac{\pi}{2}} \cos \theta - 2 \cos \theta \sin^2 \theta + \cos \theta \sin^4 \theta d\theta$$

$$= \left[\sin \theta - \frac{2}{3} \sin^3 \theta + \frac{1}{5} \sin^5 \theta \right]_0^{\frac{\pi}{2}}$$

$$= 1 - \frac{2}{3} + \frac{1}{5} = \frac{8}{15}$$

$$(d) (i) PF = \sqrt{(x-4)^2 + y^2}$$

$$PM = |x-1| \therefore PF^2 = 4 PM^2$$

$$\therefore x^2 - 8x + 16 + y^2 = 4x^2 - 8x + 4$$

$$\therefore 3x^2 - y^2 = 12 \Rightarrow \frac{x^2}{4} - \frac{y^2}{12} = 1$$

(ii) asymptotes $y = \pm \sqrt{3}x$

$\therefore$ line $y = \sqrt{3}x$ makes an angle of $\frac{\pi}{3}$ rads with positive x -axis and line $y = -\sqrt{3}x$ makes an angle of $\frac{2\pi}{3}$ rads. $\therefore$ acute angle is $\frac{\pi}{3}$ rads.

$\rightarrow$ 1 Mark

$\rightarrow$ 1 Mark

(i) $\rightarrow$ 1 Mark

(ii) $\rightarrow$ 2 Mark

Showing it's an odd function $\rightarrow$ 1 Mark

Answer = 0 $\rightarrow$ 1 Mark

$\rightarrow$ 1 Mark

Correct integration $\rightarrow$ 1 Mark

Correct answer $\rightarrow$ 1 Mark

$\rightarrow$ 1 Mark **each** for PF and PM (by definition)

$\rightarrow$ $\frac{1}{2}$ for squaring both sides & expanding


$\rightarrow$ $\frac{1}{2}$ for correct locus.

Correct equations of asymptotes $\rightarrow$ 1 Mark

Correct answer with some explanation $\rightarrow$ 1 Mark


Q3 (b) (i) $y = |f(x)|$

1



(ii) $y = \frac{1}{f(x)}$

2



Question 4

(a) $\int_0^{\frac{\pi}{2}} \frac{dx}{2 + \sin x}$

[when $x = 0, t = 0$ and $x = \frac{\pi}{2}$ then $t = 1$]

$$= \int_0^1 \frac{\frac{2}{1+t^2}}{2 + \frac{2t}{1+t^2}} dt$$

$$= \int_0^1 \frac{2 dt}{2 + 2t^2 + 2t} = \int_0^1 \frac{dt}{t^2 + t + 1}$$

$$= \int_0^1 \frac{dt}{\left(t + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$= \frac{2}{\sqrt{3}} \left[\tan^{-1} \left(\frac{t + \frac{1}{2}}{\frac{\sqrt{3}}{2}} \right) \right]_0^1 = \frac{2}{\sqrt{3}} \left[\tan^{-1} \left(\frac{2t+1}{\sqrt{3}} \right) \right]_0^1$$

$$= \frac{2}{\sqrt{3}} \left[\tan^{-1} \sqrt{3} - \tan^{-1} \frac{1}{\sqrt{3}} \right] = \frac{\pi}{3\sqrt{3}}$$

(b) (i) $\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_a^{2a} f(x) dx$

Now $\int_a^{2a} f(x) dx = \int_a^0 f(2a-u) \cdot -du$

$$= \int_0^a f(2a-u) du = \int_0^a f(2a-x) dx$$

Since the definite integral is independent of the variable.

Hence $\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$

$$= \int_0^a [f(x) + f(2a-x)] dx$$

Change of variables & correct t substitution
→ 2 Marks

→ 1 Mark

Correct integration & substitution → 1 Mark

→ 1 Mark

Let $u = 2a - x$ then $dx = -du$ & $x = a \rightarrow u = a$ & $x = 2a \rightarrow u = 0$. → 1 Mark

→ 1 Mark

$$\begin{aligned}
 \text{(ii)} \quad & \int_0^{\pi} \frac{x \, dx}{2 + \sin x} \\
 &= \int_0^{\frac{\pi}{2}} \left[\frac{x}{2 + \sin x} + \frac{\pi - x}{2 + \sin(\pi - x)} \right] dx \\
 &= \int_0^{\frac{\pi}{2}} \left[\frac{\pi}{2 + \sin x} \right] dx \text{ since } \sin x = \sin(\pi - x) \\
 &= \pi \cdot \frac{\pi}{3\sqrt{3}} \text{ from part (a)} \\
 &= \frac{\pi^2}{3\sqrt{3}}
 \end{aligned}$$

Correct use of part (i) → 1 Mark

→ 1 Mark

→ 1 Mark

$$\begin{aligned}
 \text{(c) (i)} \quad & \int_0^{\frac{\pi}{4}} \sec \theta \, d\theta = \left[\ln(\sec \theta + \tan \theta) \right]_0^{\frac{\pi}{4}} \\
 &= \ln \left(\sec \frac{\pi}{4} + \tan \frac{\pi}{4} \right) - \ln(\sec 0 + \tan 0) \\
 &= \ln(\sqrt{2} + 1)
 \end{aligned}$$

Correct integration → 1 Mark

Correct answer → 1 Mark

$$\begin{aligned}
 \text{(ii)} \quad & \int_0^{\frac{\pi}{4}} \sec^3 \theta \, d\theta = \int \sec \theta \cdot \frac{d}{d\theta} \tan \theta \, d\theta \\
 &= \left[\sec \theta \tan \theta \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \tan \theta \sec \theta \tan \theta \, d\theta \\
 &= \sqrt{2} - \int_0^{\frac{\pi}{4}} \sec \theta \tan^2 \theta \, d\theta \\
 &= \sqrt{2} - \int_0^{\frac{\pi}{4}} (\sec^3 \theta - \sec \theta) \, d\theta \\
 &\therefore 2 \int_0^{\frac{\pi}{4}} \sec^3 \theta \, d\theta = \sqrt{2} + \int_0^{\frac{\pi}{4}} \sec \theta \, d\theta \\
 &= \sqrt{2} + \ln(\sqrt{2} + 1) \\
 &\therefore \int_0^{\frac{\pi}{4}} \sec^3 \theta \, d\theta = \frac{1}{2} \left[\sqrt{2} + \ln(\sqrt{2} + 1) \right]
 \end{aligned}$$

→ 1 Mark

→ 1 Mark

→ 1 Mark using part (i)