

Intervalo de confianza	Estadístico de prueba	Criterio de rechazo
1 $\left[\bar{X} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \bar{X} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right]$	$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$	$ Z > z_{\alpha/2}$ $Z > z_{\alpha}$ $Z < -z_{\alpha}$
2 $\left[\bar{X} - t_{\alpha/2, n-1} \frac{S}{\sqrt{n}}, \bar{X} + t_{\alpha/2, n-1} \frac{S}{\sqrt{n}} \right]$	$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$	$ T > t_{\alpha/2, n-1}$ $T > t_{\alpha, n-1}$ $T < -t_{\alpha, n-1}$
3 $\left[\bar{X} - z_{\alpha/2} \frac{S}{\sqrt{n}}, \bar{X} + z_{\alpha/2} \frac{S}{\sqrt{n}} \right]$	$Z = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$	$ Z > z_{\alpha/2}$ $Z > z_{\alpha}$ $Z < -z_{\alpha}$
4 $\left[(\bar{X}_1 - \bar{X}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}, (\bar{X}_1 - \bar{X}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \right]$	$Z = \frac{(\bar{X}_1 - \bar{X}_2) - \Delta_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$	$ Z > z_{\alpha/2}$ $Z > z_{\alpha}$ $Z < -z_{\alpha}$
5 $\left[(\bar{X}_1 - \bar{X}_2) - z_{\alpha/2} \sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}, (\bar{X}_1 - \bar{X}_2) + z_{\alpha/2} \sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}} \right]$	$Z = \frac{(\bar{X}_1 - \bar{X}_2) - \Delta_0}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}}$	$ Z > z_{\alpha/2}$ $Z > z_{\alpha}$ $Z < -z_{\alpha}$
6 $\left[(\bar{X}_1 - \bar{X}_2) - t_{\alpha/2, n_1+n_2-2} S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}, (\bar{X}_1 - \bar{X}_2) + t_{\alpha/2, n_1+n_2-2} S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \right]$ con $S_p^2 = \frac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{n_1+n_2-2}$	$T = \frac{(\bar{X}_1 - \bar{X}_2) - \Delta_0}{S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$	$ T > t_{\alpha/2, n_1+n_2-2}$ $T > t_{\alpha, n_1+n_2-2}$ $T < -t_{\alpha, n_1+n_2-2}$
7 $\left[(\bar{X}_1 - \bar{X}_2) - t_{\alpha/2, \nu} \sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}, (\bar{X}_1 - \bar{X}_2) + t_{\alpha/2, \nu} \sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}} \right]$ con $\nu = \frac{\left(\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}\right)^2}{\frac{(S_1^2/n_1)^2}{n_1-1} + \frac{(S_2^2/n_2)^2}{n_2-1}}$	$T^* = \frac{(\bar{X}_1 - \bar{X}_2) - \Delta_0}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}}$	$ T^* > t_{\alpha/2, \nu}$ $T^* > t_{\alpha, \nu}$ $T^* < -t_{\alpha, \nu}$
8 $\left[\bar{D} - t_{\alpha/2, n-1} \frac{S_D}{\sqrt{n}}, \bar{D} + t_{\alpha/2, n-1} \frac{S_D}{\sqrt{n}} \right]$	$T = \frac{\bar{D} - \Delta_0}{S_D/\sqrt{n}}$	$ T > t_{\alpha/2, n-1}$ $T > t_{\alpha, n-1}$ $T < -t_{\alpha, n-1}$
9 $\left[\frac{(n-1)S^2}{\chi_{\alpha/2, n-1}^2}, \frac{(n-1)S^2}{\chi_{1-\alpha/2, n-1}^2} \right]$	$V = \frac{(n-1)S^2}{\sigma_0^2}$	$V > \chi_{\alpha/2, n-1}^2$ ó $V < \chi_{1-\alpha/2, n-1}^2$ $V > \chi_{\alpha, n-1}^2$ $V < \chi_{1-\alpha, n-1}^2$
10 $\left[\frac{S_1^2}{S_2^2} f_{1-\alpha/2, n_2-1, n_1-1}, \frac{S_1^2}{S_2^2} f_{\alpha/2, n_2-1, n_1-1} \right]$	$F = \frac{S_1^2}{S_2^2}$	$F > f_{\alpha/2, n_1-1, n_2-1}$ ó $F < f_{1-\alpha/2, n_1-1, n_2-1}$ $F > f_{\alpha, n_1-1, n_2-1}$ $F < f_{1-\alpha, n_1-1, n_2-1}$
11 $\left[\hat{P} - z_{\alpha/2} \sqrt{\frac{\hat{P}(1-\hat{P})}{n}}, \hat{P} + z_{\alpha/2} \sqrt{\frac{\hat{P}(1-\hat{P})}{n}} \right]$	$Z = \frac{\hat{P} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$	$ Z > z_{\alpha/2}$ $Z > z_{\alpha}$ $Z < -z_{\alpha}$
12 $\left[\hat{P}_1 - \hat{P}_2 - z_{\alpha/2} \sqrt{\frac{\hat{P}_1(1-\hat{P}_1)}{n_1} + \frac{\hat{P}_2(1-\hat{P}_2)}{n_2}}, \hat{P}_1 - \hat{P}_2 + z_{\alpha/2} \sqrt{\frac{\hat{P}_1(1-\hat{P}_1)}{n_1} + \frac{\hat{P}_2(1-\hat{P}_2)}{n_2}} \right]$	$Z = \frac{\hat{P}_1 - \hat{P}_2}{\sqrt{\hat{P}(1-\hat{P})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$ $\hat{P} = \frac{n_1 \hat{P}_1 + n_2 \hat{P}_2}{n_1 + n_2}$	$ Z > z_{\alpha/2}$ $Z > z_{\alpha}$ $Z < -z_{\alpha}$