

Formal Verification of Cycle Integral Properties from First Principles

Thiago Henrique Ramos da Mata

Independent Researcher

thiago.henrique.mata@gmail.com

<https://thiagomata.com/>

<https://github.com/thiagomata>

Abstract

In previous articles, we defined bounded Lists, Integrals of Lists, and unbounded Cycles of Integers from scratch, relying only on core type constructs and recursion, with no prior knowledge of Scala’s collections required. From that, we proved and formally verified some properties related to them. This article uses that as a foundation to define Integral of Cycles using two presentations: the canonical recursive `CycleIntegral` definition and a `ModCycleIntegral` closed-form definition. For both presentations, we formally verify the sum property (integral equals cumulative cycle sum) and the step property (difference between consecutive values equals the corresponding cycle element) using the Stainless verification system. We also prove that the recursive and modulo definitions are extensionally equivalent. All properties are expressed and proved within a minimal framework using only elementary arithmetic, recursion, and pure Scala code. This work bridges mathematical foundations and executable verification, offering a self-contained, verifiable approach for reasoning about infinite periodic accumulations.

Licensed under the Creative Commons Attribution 4.0 International License (CC BY 4.0):

<https://creativecommons.org/licenses/by/4.0/>.

1 Introduction

Cycles are a powerful concept in computer science and mathematics, representing unbounded lists that repeat a finite sequence of elements. When we integrate cycles, we obtain a list of cumulative sums with some unique properties that we will explore in this article.

$$L = [l_0, l_1, l_2, \dots, l_{n-1}] \mid l_n \in \mathbb{S}, L \in \mathbb{L}$$

$$\begin{aligned} \text{Cycle}(L) &= [l_0, l_1, l_2, \dots, l_{n-1}, l_0, l_1, \dots] \\ &= [v_0, v_1, v_2, \dots] \mid v_i = L[i \bmod n] \end{aligned}$$

$$\text{Integral}(L, \text{init}) = [y_0, y_1, y_2, \dots, y_{n-1}] \mid y_k = \sum_{i=0}^k x_i + \text{init}$$

$$\text{CycleIntegral}(L, \text{init}) = [w_0, w_1, w_2, \dots] \mid w_k = \sum_{i=0}^k \text{Cycle}(L)_i + \text{init}$$

In this article, we present a discrete definition of Cycle Integral over finite integer lists, defined recursively and verified some of its properties using the Stainless system. Our approach follows a zero-prior-knowledge philosophy, building on a previously verified foundation for recursive list and

integral structures and summation. The result is a verified, from-scratch implementation of the cycle integral suitable as a foundation for higher-level numeric reasoning over unbounded lists.

This article verifies:

- Two equivalent definitions: recursive and modulo-based — Subsections 3.1–3.3
- Core properties: next position, same difference after cycle, sum of mod values, strictly increasing, positivity — Subsections 4.1–4.5
- Persistent and periodic properties: how a fixed cycle integral’s residues behave forever, and how it advances across full periods — Subsections 5.1–5.6
- Deriving new cycle integrals: expansion, index shifts, rotation, survivor filtering, and merge-based reconstruction — Subsections 6.1–6.10

1.1 Related work

Lean’s Mathlib provides a general formal theory of periodic functions. It proves that a periodic function remains periodic under integer multiples of a period, and that a finite sum of periodic functions is periodic [7]. Mathlib also represents a finite cycle as a list modulo cyclic rotation [6]. These results give formal context for the finite period and shift structure used by the present construction.

This article studies a different object: the cumulative integral of a concrete periodic integer list. Its central verified results are the equivalence of a recursive integral and a quotient–remainder closed form, together with the resulting step, full-period, residue, and reconstruction properties. The existing periodicity and finite-cycle developments therefore enrich the setting without standing in for this two-presentation cycle-integral proof.

2 Preliminaries

We reuse several basic list, cycle and integral operations and their verified properties from the companion articles *Using Formal Verification to Prove Properties of Lists Recursively Defined* [5], *Formal Verification of Discrete Integration Properties from First Principles* [4], and *Formal Verification of Cyclic Lists* [3]. We also reuse some modulo properties previously defined and verified in the article *Division and Modulo from Recursive Normalization* [2].

These articles also defined and verified their properties using the same zero-prior-knowledge methodology, and are treated here as foundational primitives.

3 Cycle Integral Definitions

The cycle integral extends the finite integral to unbounded repeating sequences. Two equivalent definitions are proven.

- Recursive: recurrence on the cycle position — Subsection 3.1
- Modulo: closed-form using `div` and `mod` — Subsection 3.2
- The two definitions are extensionally equivalent — Subsection 3.3

$$\forall i \in \mathbb{N}_0, \text{init} \in \mathbb{N}_0, L \in \mathbb{L}, n = |L|$$

$$\text{CycleIntegral}(L, \text{init}) := [w_0, w_1, w_2, \dots]$$

$$0 \leq i < n \implies w_i = \sum_{j=0}^i L_j + \text{init} \quad [\text{Sum Property}]$$

$$i > 0 \implies w_i - w_{i-1} = L_{(i \bmod n)} \quad [\text{Step Property}]$$

3.1 Recursive Cycle Integral

$$\text{RecCycle}(L) = [v_0, v_1, \dots] \mid v_i = \begin{cases} L[i] & i < n \\ v_{i-n} & i \geq n \end{cases}$$

$$\text{RecCycleIntegral}(L, \text{init}) = [w_0, w_1, \dots] \mid w_i = \begin{cases} \text{init} + w_0 & i = 0 \\ v_i + w_{i-1} & i > 0 \end{cases}$$

Recursive Cycle Equivalence: Recursive Cycle is equivalent to Mod Cycle, as proven in the article [Formal Verification of Cyclic Lists](#) [3].

$$\begin{aligned} \text{RecCycle}(L)_i &= \text{ModCycle}(L)_i && [\text{Cycle Equivalence}] \\ &= L_{(i \bmod n)} && [\text{Mod Cycle Definition}] \end{aligned}$$

Sum Property:

$$v_0 = L_0 \quad [\text{Base Case}]$$

$$= L_{(0 \bmod n)} \quad [\text{By Modulo Property}]$$

$$= \sum_{j=0}^0 L_j \quad [\text{Summation Re-indexing}]$$

$$i < n \implies$$

$$w_0 = \text{init} + v_0 \quad [\text{Base Case}]$$

$$= \text{init} + \sum_{j=0}^i L_j \quad [\text{Summation Re-indexing}]$$

$$i > 0 \implies$$

$$w_i = \text{init} + v_0 + \sum_{j=1}^i v_j \quad [\text{By Definition}]$$

$$= \text{init} + v_0 + \sum_{j=1}^i L_j \quad [\text{Mod of Small Values Property}]$$

$$= \text{init} + \sum_{j=0}^i v_j \quad [\text{Summation Re-indexing}]$$

$$\therefore w_i = \text{init} + \sum_{j=0}^i L_j \quad [\text{Q.E.D.}]$$

This property is verified in the [CycleIntegralProperties::assertCycleIntegralEqualsSumSmallPositions](#). A key Scala verification excerpt is in Appendix A.1; the complete proof is linked in the source reference.

Step Property:

$$\begin{aligned}
 w_i - w_{i-1} &= v_i + w_{i-1} - w_{i-1} && [\text{By Definition}] \\
 &= v_i && [\text{Simplification}] \\
 &= L_{(i \bmod n)} && [\text{Substitution}]
 \end{aligned}$$

This property is verified in the [CycleIntegralProperties::assertDiffEqualsCycleValue](#). A key Scala verification excerpt is in Appendix A.2; the complete proof is linked in the source reference.

3.2 Modulo Cycle Integral

$$\begin{aligned}
 \text{ModCycle}(L)_i &:= L_{(i \bmod n)} = [w_0, w_1, \dots] \\
 I_k &:= \sum_{j=0}^k L_j \quad (0 \leq k < n) && [\text{Integral of } L] \\
 S &:= I_{n-1} && [\text{One full cycle sum}] \\
 \text{ModCycleIntegral}(L, \text{init})_i &:= (i \text{ div } n) \cdot S + I_{(i \bmod n)} + \text{init}
 \end{aligned}$$

Sum Property:

$$\begin{aligned}
 i < n &\implies w_i = \sum_{j=0}^i L_j + \text{init} && [\text{Claim to Prove}] \\
 i < n &\implies \\
 i \text{ div } n &= 0 && [\text{By Div of Small Values Property}] \\
 w_i &= (i \text{ div } n) \cdot S + I_{(i \bmod n)} + \text{init} && [\text{Definition}] \\
 &= 0 \cdot S + I_i + \text{init} && [\text{Substitution}] \\
 &= \sum_{j=0}^i L_j + \text{init} && [\text{By definition of } I_i] \\
 \therefore \forall i < n, \quad w_i &= \sum_{j=0}^i L_j + \text{init} && [\text{Q.E.D.}]
 \end{aligned}$$

This property is verified in the [ModCycleIntegralProperties::assertFirstValuesMatchIntegral](#). A key Scala verification excerpt is in Appendix A.3; the complete proof is linked in the source reference.

Step Property:

$$w_i - w_{i-1} = L_{i \bmod n}, \quad i > 0, n > 0 \quad [\text{Claim to Prove}]$$

Case $i \bmod n > 0$:

$$\begin{aligned}
i \bmod n &= ((i-1) \bmod n) + 1 && \text{[By Modulo Properties]} \\
i \operatorname{div} n &= (i-1) \operatorname{div} n && \text{[By Division Properties]} \\
w_i &= (i \operatorname{div} n) S + I_{i \bmod n} + \textit{init} && \text{[Definition]} \\
&= (i-1 \operatorname{div} n) S + I_{((i-1) \bmod n)+1} + \textit{init} && \text{[Div/Mod property]} \\
w_{i-1} &= (i-1 \operatorname{div} n) S + I_{(i-1) \bmod n} + \textit{init} && \text{[Definition]} \\
w_i - w_{i-1} &= I_{((i-1) \bmod n)+1} - I_{(i-1) \bmod n} && \text{[Cancellation]} \\
&= \left(\sum_{j=0}^{(i-1) \bmod n} L_j + L_{((i-1) \bmod n)+1} \right) - \sum_{j=0}^{(i-1) \bmod n} L_j && \text{[Expand sum]} \\
&= L_{((i-1) \bmod n)+1} && \text{[Cancellation]} \\
&= L_{i \bmod n} && \text{[Modulo property]}.
\end{aligned}$$

Case $i \bmod n = 0$:

$$\begin{aligned}
w_i &= (i \operatorname{div} n) S + L_0 + \textit{init} && \text{[Definition]} \\
w_{i-1} &= (i \operatorname{div} n - 1) S + I_{n-1} + \textit{init} && \text{[Div/Mod property]} \\
w_i - w_{i-1} &= (i \operatorname{div} n) S + L_0 + \textit{init} - ((i \operatorname{div} n - 1) S + I_{n-1} + \textit{init}) && \text{[Substitution]} \\
&= S + L_0 - I_{n-1} && \text{[Simplification]} \\
&= L_0 && \text{[Since } S = I_{n-1}\text{]} \\
&= L_{i \bmod n} && \text{[Modulo property]}.
\end{aligned}$$

$$\therefore w_i - w_{i-1} = L_{i \bmod n}, \quad \forall i > 0 \quad \text{[Q.E.D.]}$$

This property is verified in the `ModCycleIntegralProperties::assertSimplifiedDiffValuesMatchCycle`. A key Scala verification excerpt is in Appendix A.4; the complete proof is linked in the source reference.

3.3 Equivalence of Definitions

The nontrivial equivalence proved here is between the recursive `CycleIntegral` definition and the closed-form `ModCycleIntegral` definition.

$$\begin{aligned}
\text{CycleIntegral}(L, \textit{init}) &= \text{ModCycleIntegral}(L, \textit{init}) \\
&= [w_0, w_1, w_2, \dots] \mid w_i = \sum_{j=0}^i L_{(j \bmod n)} + \textit{init} \blacksquare
\end{aligned}$$

This property is verified in the `ModCycleIntegralProperties::assertCycleIntegralMatchModCycleDef`. A key Scala verification excerpt is in Appendix A.5; the complete proof is linked in the source reference.

4 Core Verified Properties

The fundamental properties of the cycle integral that hold for every position.

- Next position: $CI_{i+1} = CI_i + \text{Cycle}(L)_{i+1}$ — Subsection 4.1
- Full cycle shift: adding one cycle period advances by the total sum — Subsection 4.2
- Sum of mod values: the modulo definition matches the list sum — Subsection 4.3
- Strictly increasing: positive base values force a strictly growing integral — Subsection 4.4
- Positivity: a non-negative start and positive base values keep the integral positive everywhere — Subsection 4.5

4.1 Next Position

For any positive position, the cycle integral at that position equals the previous value plus the current cycle element.

$$\forall i > 0 : CI(L, init)_i = CI(L, init)_{i-1} + \text{Cycle}(L)_i$$

This follows directly from the recursive definition of `CycleIntegral`.

This property is verified in the `CycleIntegralProperties::assertNextPosition`.

4.2 Same Difference After Full Cycle

The difference between consecutive values is invariant under adding a full cycle size to both positions.

$$\forall i \geq 0 : CI(L, init)_{i+1} - CI(L, init)_i = CI(L, init)_{i+size+1} - CI(L, init)_{i+size}$$

This property is verified in the `CycleIntegralProperties::assertSameDiffAfterCycle`. A key Scala verification excerpt is in Appendix A.6; the complete proof is linked in the source reference.

4.3 Sum of Mod Values as List

The cycle integral at any position equals the sum of a constructed list containing the initial value and all cycle values up to that position (using modular indexing for positions beyond one cycle).

$$\forall i \geq 0 : CI(L, init)_i = \text{sum}([init] \\ + [\text{Cycle}(L)_0, \text{Cycle}(L)_1, \dots, \text{Cycle}(L)_i])$$

This property is verified in the `CycleIntegralProperties::assertSumModValueAsListEqualsCycleIntegralLoop`. A key Scala verification excerpt is in Appendix A.7; the complete proof is linked in the source reference.

4.4 Cycle Integral Strictly Increasing

When the initial value is non-negative and every base-list value is positive, the cycle integral is strictly increasing: a later position always produces a larger value.

$$init \geq 0 \wedge (\forall x \in L, x > 0) \wedge b > a \implies \text{CycleIntegral}(L, init)_b > \text{CycleIntegral}(L, init)_a$$

Proof. Induct on $b - a$. The base case is one positive step; the inductive step adds one more positive cycle value.

$$\begin{aligned}
b = a + 1 &\implies CI_b - CI_a = \text{Cycle}(L)_b > 0 \\
&\quad [\S 3.1 \text{ and cycle positivity}] \\
&\implies CI_b > CI_a, \\
CI_{b-1} > CI_a, \quad CI_b - CI_{b-1} = \text{Cycle}(L)_b > 0 &\implies CI_b > CI_{b-1} > CI_a \\
&\therefore CI_b > CI_a. \quad \blacksquare [\text{Q.E.D.}]
\end{aligned}$$

This property is verified in the `CycleIntegralProperties::assertCycleIntegralIncreasing`.

4.5 Cycle Integral Positivity

When the initial value is non-negative and every base-list value is positive, the cycle integral is positive at every position.

$$init \geq 0 \wedge (\forall x \in L, x > 0) \implies \text{CycleIntegral}(L, init)_i > 0$$

Proof. At the first position, the non-negative initial value and a positive cycle value give a positive integral. Every later position adds one more positive cycle value.

$$\begin{aligned}
CI_0 &= init + \text{Cycle}(L)_0 > 0, \\
CI_{i-1} > 0, \quad CI_i - CI_{i-1} = \text{Cycle}(L)_i > 0 &\implies CI_i > 0 \\
&\therefore CI_i > 0. \quad \blacksquare [\text{Q.E.D.}]
\end{aligned}$$

This property is verified in the `CycleIntegralProperties::assertCycleIntegralPositive`.

5 Persistent and Periodic Properties

These properties describe a single, fixed cycle integral: how its residues behave forever, and how it advances across full periods. All six properties are fully Stainless-verified, except Properties 5.3 and 5.4, which are direct corollaries of Property 5.2.

- Cycle-period shifts: a full period advances the integral by the cycle sum — Subsection 5.1
- General residue periodicity: any residue depends only on the position within one cycle period — Subsection 5.2
- Persistent non-zero residue: if no residue in one period is zero, none ever is — Subsection 5.3
- Persistent zero residue: if every residue in one period is zero, every residue always is — Subsection 5.4
- Gap telescoping: two consecutive gaps sum to the integral span across both — Subsection 5.5
- Residue classification: all-zero, some-zero, none-zero — Subsection 5.6

5.1 Cycle-Period Shifts

After a full cycle period, the integral advances by the cycle's total sum. This is the termination bound for scanning: a survivor is always found within one cycle period because the integral advances by a fixed, positive amount.

Both quantities below belong to the finite backing cycle that ci accumulates over, not to ci 's own output stream — which is unbounded and, by Section 4.4, strictly increasing and never repeats:

$$\begin{aligned} n &:= \text{period}(ci) && \text{[Length of the finite backing cycle]} \\ \text{periodSum}(ci) &:= \sum_{j=0}^{n-1} \text{cycle}(j) && \text{[Total of one period's gap values]} \end{aligned}$$

$$\begin{aligned} ci(\text{pos} + \text{period}(ci)) &= ci(\text{pos}) + \text{periodSum}(ci) && \text{[Full-cycle shift]} \\ ci(\text{pos} + \text{period}(ci) \cdot m) &= ci(\text{pos}) + m \cdot \text{periodSum}(ci) && \text{[Multi-cycle shift, by induction on } m\text{]} \end{aligned}$$

Proof.

Step 0 (base identity, $pos = 0$). Unfolding the recursive definition of ci across one full period, using the cycle's own periodicity ($\text{cycle}(\text{period}(ci)) = \text{cycle}(0)$, by definition of a cycle — Section 1) and reordering the resulting finite sum:

$$\begin{aligned} ci(\text{period}(ci)) - ci(0) &= \text{cycle}(1) + \dots + \text{cycle}(\text{period}(ci) - 1) \\ &\quad + \text{cycle}(\text{period}(ci)) && \text{[Telescoping the recursive definition]} \\ &= \text{cycle}(1) + \dots + \text{cycle}(\text{period}(ci) - 1) + \text{cycle}(0) && \text{[Cycle periodicity]} \\ &= \text{cycle}(0) + \text{cycle}(1) + \dots + \text{cycle}(\text{period}(ci) - 1) && \text{[Reorder terms]} \\ &= \text{periodSum}(ci) && \text{[By Definition of periodSum}(ci)\text{]} \end{aligned}$$

$$\therefore ci(\text{period}(ci)) = ci(0) + \text{periodSum}(ci) \quad \blacksquare$$

Full-cycle shift, by induction on pos .

Base Case ($pos = 0$): shown in Step 0.

Induction Step ($pos > 0$):

$$\begin{aligned} ci(\text{pos} - 1 + \text{period}(ci)) &= ci(\text{pos} - 1) + \text{periodSum}(ci) && \text{[Induction Hypothesis]} \\ \text{cycle}(\text{pos} + \text{period}(ci)) &= \text{cycle}(\text{pos}) && \text{[Cycle periodicity]} \\ ci(\text{pos} + \text{period}(ci)) &= ci(\text{pos} - 1 + \text{period}(ci)) \\ &\quad + \text{cycle}(\text{pos} + \text{period}(ci)) && \text{[By Definition]} \\ &= (ci(\text{pos} - 1) + \text{periodSum}(ci)) + \text{cycle}(\text{pos}) && \text{[Substitution]} \\ &= (ci(\text{pos} - 1) + \text{cycle}(\text{pos})) + \text{periodSum}(ci) && \text{[Regroup]} \\ &= ci(\text{pos}) + \text{periodSum}(ci) && \text{[By Definition]} \end{aligned}$$

$$\therefore \forall pos \in \mathbb{N}_0, ci(\text{pos} + \text{period}(ci)) = ci(\text{pos}) + \text{periodSum}(ci) \quad \blacksquare$$

Multi-cycle shift, by induction on m .

Base Case ($m = 0$):

$$ci(\text{pos} + \text{period}(ci) \cdot 0) = ci(\text{pos}) = ci(\text{pos}) + 0 \cdot \text{periodSum}(ci)$$

Induction Step ($m > 0$):

$$\begin{aligned}
ci(\text{pos} + \text{period}(ci) \cdot (m - 1)) &= ci(\text{pos}) \\
&\quad + (m - 1) \cdot \text{periodSum}(ci) \quad [\text{Induction Hypothesis}] \\
ci((\text{pos} + \text{period}(ci) \cdot (m - 1)) + \text{period}(ci)) &= ci(\text{pos} + \text{period}(ci) \cdot (m - 1)) \\
&\quad + \text{periodSum}(ci) \quad [\text{Full-cycle shift}] \\
ci(\text{pos} + \text{period}(ci) \cdot m) &= (ci(\text{pos}) + (m - 1) \cdot \text{periodSum}(ci)) \\
&\quad + \text{periodSum}(ci) \quad [\text{Substitution}] \\
&= ci(\text{pos}) + m \cdot \text{periodSum}(ci) \quad [\text{Arithmetic}]
\end{aligned}$$

$$\therefore \forall m \in \mathbb{N}_0, ci(\text{pos} + \text{period}(ci) \cdot m) = ci(\text{pos}) + m \cdot \text{periodSum}(ci) \quad \blacksquare$$

The full-cycle shift identity is verified twice under different names — `GapProperties::assertPeriodicShift` and `assertFullCycleShift` — both thin wrappers over the same `CycleIntegralFilterProperties::assertCIShiftEqualsSum` lemma; the article proves the one identity above rather than two. The multi-cycle generalization is verified separately as `assertMultiCycleShift`. The core Scala verification code is in Appendix A.12.

5.2 General Residue Periodicity

When the total sum of a cycle’s values is a multiple of m , the residue $\text{mod}(ci(\text{pos}), m)$ depends only on $\text{pos} \% ci.\text{period}$ — it repeats every cycle period. When m is a product of coprime values, the Chinese Remainder Theorem [1] implies the periodicity holds simultaneously for each factor: the cycle period serves as a common period for all residues. This is the arithmetic backbone of Eratosthenes’ sieve [1].

$$\text{mod}(\text{periodSum}(ci), m) = 0$$

$$\implies \text{mod}(ci(\text{pos}), m) = \text{mod}(ci(\text{pos} \bmod \text{period}(ci)), m)$$

Proof. By strong induction on pos , subtracting one full cycle period at a time.

Base Case ($\text{pos} < \text{period}(ci)$):

$$\begin{aligned}
\text{pos} \bmod \text{period}(ci) &= \text{pos} \quad [\text{Mod of a value smaller than the divisor}] \\
\text{mod}(ci(\text{pos}), m) &= \text{mod}(ci(\text{pos} \bmod \text{period}(ci)), m) \quad [\text{Substitution}]
\end{aligned}$$

Induction Step ($\text{pos} \geq \text{period}(ci)$): let $\text{previous} := \text{pos} - \text{period}(ci)$. By the one-period shift identity proven in Subsection 5.1, $ci(\text{previous} + \text{period}(ci)) = ci(\text{previous}) + \text{periodSum}(ci)$, i.e. $ci(\text{pos}) = ci(\text{previous}) + \text{periodSum}(ci)$:

$$\text{previous} \bmod \text{period}(ci) = \text{pos} \bmod \text{period}(ci) \quad [\text{Subtracting a full period does not}$$

$$\text{change the position's residue}]$$

$$\text{mod}(ci(\text{previous}), m) = \text{mod}(ci(\text{previous mod period}(ci)), m) \quad [\text{Induction Hypothesis}]$$

$$ci(\text{pos}) = ci(\text{previous}) + \text{periodSum}(ci) \quad [\text{Full-cycle shift, §5.1}]$$

$$\text{mod}(ci(\text{pos}), m) = \text{mod}(ci(\text{previous}) + \text{periodSum}(ci), m) \quad [\text{Substitution}]$$

$$\begin{aligned} \text{mod}(ci(\text{pos}), m) &= \text{mod}(ci(\text{previous}) + \text{periodSum}(ci), m) \quad [\text{Substitution}] \\ &= \text{mod}(ci(\text{previous}), m) \quad [\text{Since } \text{mod}(\text{periodSum}(ci), m) = 0] \\ &= \text{mod}(ci(\text{previous mod period}(ci)), m) \quad [\text{By Induction Hypothesis}] \\ &= \text{mod}(ci(\text{pos mod period}(ci)), m) \quad [\text{By the residue equality above}] \end{aligned}$$

$$\therefore \text{mod}(\text{periodSum}(ci), m) = 0 \implies \forall \text{ pos} \in \mathbb{N}_0,$$

$$\text{mod}(ci(\text{pos}), m) = \text{mod}(ci(\text{pos mod period}(ci)), m) \quad \blacksquare [\text{Q.E.D.}]$$

This property is verified in the `GapProperties::assertModIsPeriodic`. The full Scala verification code is in Appendix A.11.

The companion div/mod decomposition `ci(pos) == ci(pos % size) + (pos / size) * ci.sum` is verified in `GapProperties::assertCModDivFormula`.

5.3 Persistent Non-Zero Residue

Let $v \in \mathbb{N}$, $v > 0$, with $\text{mod}(\text{periodSum}(ci), v) = 0$. If none of the n residues in one full period is zero mod v , then the residue is never zero at any position, forever.

$$\text{mod}(\text{periodSum}(ci), v) = 0 \wedge (\forall k \in [0, n), \text{mod}(ci(k), v) \neq 0) \implies \forall i \in \mathbb{N}_0, \text{mod}(ci(i), v) \neq 0$$

Proof. By Subsection 5.2, $\text{mod}(ci(i), v) = \text{mod}(ci(i \bmod n), v)$ for every position i . Since $i \bmod n$ always falls in $[0, n)$, and none of those n residues is zero by hypothesis, the residue at any position i cannot be zero either.

$$\begin{aligned} \text{mod}(ci(i), v) &= \text{mod}(ci(i \bmod n), v) \quad [\S 5.2] \\ &\neq 0 \quad [\text{In-period hypothesis}] \\ \therefore \text{mod}(ci(i), v) &\neq 0. \quad \blacksquare [\text{Q.E.D.}] \end{aligned}$$

This is a direct corollary of the periodicity lemma `GapProperties::assertModIsPeriodic` used in Subsection 5.2; no separate lemma is needed once every in-period residue is checked. The full Scala verification code is in Appendix A.11.

5.4 Persistent Zero Residue

The mirror-image case: if every residue in one full period is zero mod v , then the residue stays zero at every position, forever.

$$\text{mod}(\text{periodSum}(ci), v) = 0 \wedge (\forall k \in [0, n), \text{mod}(ci(k), v) = 0) \implies \forall i \in \mathbb{N}_0, \text{mod}(ci(i), v) = 0$$

Proof. Identical to Subsection 5.3, with the inequality reversed: by Subsection 5.2, $\text{mod}(ci(i), v) = \text{mod}(ci(i \bmod n), v)$, and $i \bmod n$ always falls among the n in-period residues, all of which are zero by hypothesis.

$$\begin{aligned} \text{mod}(ci(i), v) &= \text{mod}(ci(i \bmod n), v) && [\S 5.2] \\ &= 0 && [\text{In-period hypothesis}] \\ \therefore \text{mod}(ci(i), v) &= 0. && \blacksquare [\text{Q.E.D.}] \end{aligned}$$

This is the same corollary of `GapProperties::assertModIsPeriodic` as Subsection 5.3, applied to the opposite hypothesis. The full Scala verification code is in Appendix A.11.

5.5 Gap Telescoping

Two consecutive gaps telescope to the integral difference across both steps. By applying the one-step difference lemma twice, the gap values at positions k and $k + 1$ add up to the integral span from $k - 1$ to $k + 1$. This is the step-lemma that underlies merging adjacent gap ranges.

$$ci(k + 1) - ci(k - 1) = \text{cycle}(k) + \text{cycle}(k + 1) \quad \text{for } 1 \leq k \text{ and } k + 1 < \text{period}(ci).$$

Proof. The one-step property at positions $k - 1$ and k gives

$$\begin{aligned} ci(k) - ci(k - 1) &= \text{cycle}(k), \\ ci(k + 1) - ci(k) &= \text{cycle}(k + 1). \end{aligned}$$

Adding these equalities cancels $ci(k)$, so

$$\therefore ci(k + 1) - ci(k - 1) = \text{cycle}(k) + \text{cycle}(k + 1) \quad \blacksquare [\text{Q.E.D.}]$$

This property is verified in the `CycleIntegralProperties::assertConsecutiveGapSumEqualsDiff`. The full Scala verification code is in Appendix A.10.

5.6 Cycle Residue Classification

For any cycle and modulus $d > 0$, the values of the cycle fall into exactly one of three residue categories modulo d :

$$\text{all-zero: } \forall k, \text{mod}(\text{cycle}(k), d) = 0 \quad [\text{Filter removes everything}]$$

$$\text{none-zero: } \forall k, \text{mod}(\text{cycle}(k), d) \neq 0 \quad [\text{Filter has no effect}]$$

$$\text{some-zero: } \exists k_0 : \text{mod}(\text{cycle}(k_0), d) = 0 \wedge \exists k_1 :$$

$$\text{mod}(\text{cycle}(k_1), d) \neq 0 \quad [\text{Filter removes specific positions}]$$

These three states are detected by `MemCycle.checkMod(d)` and stored in lists (`modIsZeroForAllValues`, `modIsZeroForNoneValues`, `modIsZeroForSomeValues`). The evaluation is idempotent — the cycle’s values list never changes, only the classification metadata is updated. Ten lemmas in `CycleCheckMod.scala` prove the classification is correct, mutually exclusive, and exhaustive.

These properties are verified in the `CycleCheckMod` module.

6 Deriving New Cycle Integrals

These properties describe how to build a new cycle integral from an existing one: replicating its backing list, shifting its index, rotating its gaps, filtering out multiples of a value, and reconstructing the filtered result directly. Properties 6.1, 6.4–6.10 are fully Stainless-verified; Properties 6.2 and 6.3 have mathematical proofs but are not yet Stainless-verified.

- x-fold cycle expansion: the physical period changes while the represented stream is preserved — Subsection 6.1
- Index shifts: right and left — Subsections 6.2–6.3
- Gap rotation with head adjustment: rotating the gap cycle shifts the represented integral by one position — Subsection 6.4
- Survivor filtering: exactness and structure of the scan that retains non-multiples — Subsections 6.5–6.6
- Filter-merge reconstruction: the resulting gap cycle after removing a multiple — Subsections 6.7–6.10

6.1 x-fold Cycle Expansion

Let $L^{(x)}$ be the x -fold concatenation of a list $L \in \mathbb{L}$. This operation changes the physical backing cycle, but it does not change the unbounded stream represented by the cycle.

The values that change are the finite storage properties:

$$\begin{aligned} |L^{(x)}| &= x \cdot |L| && [\text{Expanded physical period}] \\ \sum L^{(x)} &= x \cdot \sum L && [\text{Expanded physical sum}] \end{aligned}$$

The period equation is verified in `RepeatedGapIntegralProperties::assertRepeatedPeriodIsMultiplied`.

The values that do not change are the semantic stream properties:

$$\begin{aligned} L_i^{(x)} &= L_{(i \bmod |L|)} && [\text{Same cycle lookup}] \\ \text{CycleIntegral}(L^{(x)}, \text{init})_i &= \text{CycleIntegral}(L, \text{init})_i && [\text{Same integral stream}] \end{aligned}$$

The cycle lookup equation is verified in `RepeatedGapIntegralProperties::assertReplicatedCycleValueEqual`.

So the expansion is not an invariance of the finite cycle object itself. It is an invariance of the infinite stream that the cycle object represents.

$$\begin{aligned}
n &= |L|, \quad L = [v_0, \dots, v_{n-1}] && \text{[Length of original cycle]} \\
L^{(x)} &:= \underbrace{L ++ \dots ++ L}_{x \text{ copies}} && \text{[Concatenate } x \text{ copies]} \\
m &:= |L^{(x)}| = x \cdot n && \text{[Length of new cycle]} \\
T &:= \sum_{j=0}^{n-1} v_j && \text{[Original cycle sum]}
\end{aligned}$$

The list-level repetition is defined by applying the original list at the original index modulo n :

$$\begin{aligned}
L_i^{(x)} &= L_{(i \bmod n)} && \text{for } 0 \leq i < x \cdot n \\
\sum L^{(x)} &= x \cdot \sum L && \text{[Repeated sum]}
\end{aligned}$$

These list-level properties are verified by `RepeatedList` and `ListRepeatProperties`: the repeated list size is $x \cdot n$, the repeated value at index i is the original value at $i \bmod n$, and the repeated sum is x times the original sum. See `RepeatedList`, `RepeatedListProperties::assertSumMultiplier`, and `ListRepeatProperties::assertRepeatedIndex`.

$$\begin{aligned}
\text{CycleIntegral}(L^{(x)}, \text{init})_i &= (i \operatorname{div} m) \cdot T^{(x)} + I_{i \bmod m}^{(x)} + \text{init} && \text{[By Definition]} \\
&= (i \operatorname{div} (x \cdot n)) \cdot (x \cdot T) + I_{i \bmod (x \cdot n)}^{(x)} + \text{init} && \text{[Substitution]} \\
&= (i \operatorname{div} n) \cdot T + I_{i \bmod n} + \text{init} && \text{[Exact simplification]} \\
&= \text{CycleIntegral}(L, \text{init})_i && \text{[Exact value reproduction]}
\end{aligned}$$

$$\therefore \forall i \in \mathbb{N}_0 : \text{CycleIntegral}(L^{(x)}, \text{init})_i = \text{CycleIntegral}(L, \text{init})_i \quad \blacksquare$$

This property is verified in the `CycleIntegralProperties::assertRepeatedValuesIntegralMatches`. The full Scala verification code is in Appendix A.8.

6.2 Right Index Shift

Let $L' \in \mathbb{L}$ be the right shift of $L \in \mathbb{L}$ by one position, and $\text{init}' := \text{init} + L_0$ be the shifted initial value. Then the `CycleIntegral` of L' with init' reproduces the `CycleIntegral` of L with init , shifted by one position.

$$\begin{aligned}
n &= |L|, \quad L = [v_0, \dots, v_{n-1}] \\
L' &:= [v_1, v_2, \dots, v_{n-1}, v_0] \\
S' &= S && \text{[Cycle Sum Invariance]} \\
\text{init}' &:= \text{init} + L_0
\end{aligned}$$

$$\text{CycleIntegral}(L, \text{init})_{i+1} = \text{CycleIntegral}(L', \text{init}')_i \quad \forall i \in \mathbb{N}_0$$

Base Case:

$$\begin{aligned}
A &:= \text{CycleIntegral}(L, \text{init})_i \\
B &:= \text{CycleIntegral}(L', \text{init}')_i \\
A_0 &= \text{init} + L_0 \\
A_1 &= \text{init} + L_0 + L_1 \\
B_0 &= \text{init}' + L'_0 = (\text{init} + L_0) + L'_0 = \text{init} + L_0 + L_1 = A_1
\end{aligned}$$

Induction Step:

$$\begin{aligned}
B_{i-1} &= A_i && \text{[Induction Hypothesis]} \\
A_{i+1} &= A_i + L_{((i+1) \bmod n)} && \text{[By Definition]} \\
B_i &= B_{i-1} + L'_{(i \bmod n)} && \text{[By Definition]} \\
&= A_i + L_{((i+1) \bmod n)} && \text{[By Induction Hypothesis]} \\
&= A_{i+1} && \text{[By Definition]}
\end{aligned}$$

$$\therefore \forall i \in \mathbb{N}_0 : \text{CycleIntegral}(L, \text{init})_{i+1} = \text{CycleIntegral}(L', \text{init}')_i \quad \blacksquare$$

The one-period **CycleIntegral** wrapper is verified directly. If the shifted cycle uses the one-step rotation of the original backing values and the shifted initial value is advanced by the original first gap, then the shifted integral at position i equals the original integral at position $i + 1$ for every stored-period index with $i + 1 < \text{period}$.

This property is verified in `GapProperties::assertRotateOneCycleIntegralShiftsByOne`. The full Scala verification code is in Appendix A.9.

The article's mathematical statement above is the all-position version. The verified lemma proves the stored-period core; packaging the universal all-position wrapper only needs the already verified full-cycle shift law.

6.3 Left Index Shift

Let $L'' \in \mathbb{L}$ be the left shift of $L \in \mathbb{L}$ by one position ($|L| > 1$), and $\text{init}'' := \text{init} + L_0 - L_{n-1}$ be the shifted initial value. Then the **CycleIntegral** of L'' with init'' reproduces the **CycleIntegral** of L with init , shifted by one position in the opposite direction.

$$\begin{aligned}
n &= |L|, \quad L = [v_0, \dots, v_{n-1}], \quad n > 1 \\
L'' &:= [v_{n-1}, v_0, v_1, \dots, v_{n-2}] \\
S'' &= S && \text{[Cycle Sum Invariance]} \\
\text{init}'' &:= \text{init} + L_0 - L_{n-1}
\end{aligned}$$

$$\text{CycleIntegral}(L, \text{init})_i = \text{CycleIntegral}(L'', \text{init}'')_{i+1} \quad \forall i \in \mathbb{N}_0$$

Base Case:

$$\begin{aligned}
C &:= \text{CycleIntegral}(L'', \text{init}'')_i \\
C_1 &= \text{init}'' + L''_0 = (\text{init} + L_0 - L_{n-1}) + L''_0 \\
&= \text{init} + L_0 - L_{n-1} + L_{n-1} = \text{init} + L_0 = A_0
\end{aligned}$$

Induction Step:

$$\begin{aligned}
C_{i+1} &= A_i && \text{[Induction Hypothesis]} \\
A_{i+1} &= A_i + L_{((i+1) \bmod n)} && \text{[By Definition]} \\
C_{i+1} &= C_i + L''_{(i \bmod n)} && \text{[By Definition]} \\
&= A_i + L_{((i-1+n) \bmod n)} && \text{[By Induction Hypothesis]} \\
&= A_{i+1} && \text{[By Definition]}
\end{aligned}$$

$$\therefore \forall i \in \mathbb{N}_0 : \text{CycleIntegral}(L, \text{init})_{i+1} = \text{CycleIntegral}(L'', \text{init}'')_i \quad \blacksquare$$

6.4 Gap Rotation with Head Adjustment

Rotating a gap cycle by one position and adjusting the head shifts the entire integral by one position.

$$\begin{aligned} & \text{GapList}(\text{head} + \text{gaps}_0, \text{tail}(\text{gaps}) ++ (\text{gaps}_0 :: L_e))_i \\ &= \text{GapList}(\text{head}, \text{gaps})_{i+1} \quad \text{for } 0 \leq i \text{ and } i+1 < \text{period}(\text{ci}). \end{aligned}$$

Proof. At $i = 0$, the adjusted head is $\text{head} + \text{gaps}_0$, which is the original integral at position 1. For $i > 0$, assume the shifted integral at $i - 1$ equals the original integral at i . The rotated gap at i is the original gap at $i + 1$; applying the one-step property to both integrals gives

$$\begin{aligned} I'_i &= I'_{i-1} + \text{gaps}'_{i-1} \\ &= I_i + \text{gaps}_i \\ &= I_{i+1}. \\ \therefore I'_i &= I_{i+1} \quad \blacksquare \text{ [Q.E.D.]} \end{aligned}$$

This property is verified in `GapProperties::assertRotateOneShiftsIntegralByOne`. It delegates to the verified `ShiftedList.assertShiftedApplyIsOriginalPlusOne`; the full source is linked there rather than repeated inline.

For a filter value f and a range $[start, start + count)$, the survivor sequence is the sub-sequence of ci 's values at those positions whose remainder modulo f is nonzero — the cycle-integral analogue of one Eratosthenes sieve step [1], where positions divisible by f are removed and only the non-multiples remain:

$$\begin{aligned} S &:= \text{survivorValues}(ci, f, start, count) := \\ &[ci(pos) \mid start \leq pos < start + count, \text{mod}(ci(pos), f) \neq 0] \end{aligned}$$

Ten verified lemmas in `GapProperties.scala` characterize this sequence.

6.5 Survivor Exactness

S is exact by construction: it is defined as exactly the non-multiples of f in range, nothing more and nothing less. The recursive Scala implementation of `survivorValues` is verified to compute exactly this specification.

This is verified in `GapProperties::assertSurvivorValuesContainsOnlyNonMultiples` and `assertSurvivorValuesContainsNonMultipleAtPosition`.

6.6 Survivor Structure

When every value in $[start, pos)$ is a multiple of f and $ci(pos)$ itself is not, $ci(pos)$ is the first survivor, and S decomposes as that value followed by the survivors of the rest of the range, (pos, end) where $end := start + count$:

$$\begin{aligned} & (\forall q \in [start, pos), \text{mod}(ci(q), f) = 0) \wedge \text{mod}(ci(pos), f) \neq 0 \\ & \implies \text{head}(S) = ci(pos) \quad \text{[First survivor]} \\ & S = ci(pos) :: [ci(q) \mid pos < q < end, \text{mod}(ci(q), f) \neq 0] \quad \text{[Structural split]} \end{aligned}$$

At the endpoints of a range that itself survives, the first and last elements of S are exactly the endpoint values, and over one complete period the difference between them equals the cycle's total gap sum — filtering does not change how far the integral advances across a full period:

$$\begin{aligned}
& \text{mod}(ci(\text{start}), f) \neq 0 \implies \text{head}(S) = ci(\text{start}) \\
& \text{mod}(ci(\text{start} + \text{count} - 1), f) \neq 0 \implies \text{last}(S) \\
& \quad = ci(\text{start} + \text{count} - 1) \\
& \text{mod}(ci(0), f) \neq 0 \wedge \text{mod}(ci(n), f) \neq 0 \implies \text{last}(S) - \text{head}(S) = \text{periodSum}(ci)
\end{aligned}$$

The gap between two consecutive survivors, once every intervening multiple has been filtered out, is strictly positive:

$$\begin{aligned}
& \text{mod}(ci(\text{from}), f) \neq 0 \wedge \text{mod}(ci(\text{to}), f) \neq 0 \wedge (\forall q \in (\text{from}, \text{to}), \text{mod}(ci(q), f) = 0) \\
& \implies ci(\text{to}) - ci(\text{from}) > 0
\end{aligned}$$

All survivor-structure properties are verified in `GapProperties::assertFirstSurvivorAtPosition`, `assertSurvivorValuesSplitAtFirstPosition`, `assertFirstSurvivorIsHead`, `assertLastSurvivorIsLastScanned`, `assertFilteredSumEqualsOriginalSum`, and `assertMergedGapPositive`.

Filtering removes a value; the surviving values must still form a valid cycle-integral gap list. This section proves that removing one multiple by merging its two neighboring gaps produces exactly the cycle integral that a fresh construction from the survivor list would produce, and that the result contains no multiples of the filter value anywhere.

6.7 Merge Shift Law

Let ci be a cycle integral with period n , and let m be a merge index with $0 \leq m$ and $m + 1 < n$. Let ci' be a cycle integral with period $n - 1$, the same initial value, and a gap cycle equal to ci 's except that the gaps at m and $m + 1$ are combined into one:

$$\begin{aligned}
& \text{cycle}'(i) = \text{cycle}(i) && \text{for } i < m \\
& \text{cycle}'(m) = \text{cycle}(m) + \text{cycle}(m + 1) \\
& \text{cycle}'(i) = \text{cycle}(i + 1) && \text{for } i > m
\end{aligned}$$

Then ci' agrees with ci before the merge point, and agrees with ci one position ahead at and after it:

$$\begin{aligned}
& pos < m \implies ci'(pos) = ci(pos) && [\text{Before merge}] \\
& ci'(m) = ci(m + 1) && [\text{At merge}] \\
& pos > m \implies ci'(pos) = ci(pos + 1) && [\text{After merge}] \\
& ci'(n - 1) = ci(n) && [\text{Period boundary}]
\end{aligned}$$

Proof. By induction on the position, using the step property (Subsection 3.1) in each of three cases split at the merge point.

Case 1: Before the merge ($pos < m$), by induction on pos .

Base Case ($pos = 0$):

$$\begin{aligned}
& ci'(0) = \text{init} + \text{cycle}'(0) && [\text{By Definition}] \\
& = \text{init} + \text{cycle}(0) && [\text{Same initial value; unchanged gap before merge}] \\
& = ci(0) && [\text{By Definition}]
\end{aligned}$$

Induction Step ($0 < pos < m$):

$$\begin{aligned}
ci'(pos - 1) &= ci(pos - 1) && \text{[Induction Hypothesis]} \\
ci'(pos) &= ci'(pos - 1) + cycle'(pos) && \text{[By Definition]} \\
&= ci(pos - 1) + cycle(pos) && \text{[Substitution; unchanged gap before merge]} \\
&= ci(pos) && \text{[By Definition]}
\end{aligned}$$

$$\therefore pos < m \implies ci'(pos) = ci(pos) \quad \blacksquare$$

Case 2: At the merge ($pos = m$). For $m > 0$, using Case 1 at $m - 1$:

$$\begin{aligned}
ci'(m) &= ci'(m - 1) + cycle'(m) && \text{[By Definition]} \\
&= ci(m - 1) + (cycle(m) + cycle(m + 1)) && \text{[Case 1; merged-gap definition]} \\
&= (ci(m - 1) + cycle(m)) + cycle(m + 1) && \text{[Regroup]} \\
&= ci(m) + cycle(m + 1) && \text{[By Definition]} \\
&= ci(m + 1) && \text{[By Definition]}
\end{aligned}$$

For $m = 0$, the same identity holds directly, with no Case 1 dependency: $ci'(0) = init + cycle'(0) = init + cycle(0) + cycle(1) = ci(1)$.

$$\therefore ci'(m) = ci(m + 1) \quad \blacksquare$$

Case 3: After the merge ($pos > m$), by induction on pos .

Base Case ($pos = m + 1$), using Case 2:

$$\begin{aligned}
ci'(m + 1) &= ci'(m) + cycle'(m + 1) && \text{[By Definition]} \\
&= ci(m + 1) + cycle(m + 2) && \text{[Case 2; shifted-gap definition]} \\
&= ci(m + 2) && \text{[By Definition]}
\end{aligned}$$

Induction Step ($pos > m + 1$):

$$\begin{aligned}
ci'(pos - 1) &= ci(pos) && \text{[Induction Hypothesis]} \\
ci'(pos) &= ci'(pos - 1) + cycle'(pos) && \text{[By Definition]} \\
&= ci(pos) + cycle(pos + 1) && \text{[Substitution; shifted-gap definition]} \\
&= ci(pos + 1) && \text{[By Definition]}
\end{aligned}$$

$$\therefore pos > m \implies ci'(pos) = ci(pos + 1) \quad \blacksquare$$

The period boundary case is the last instance of Case 3, at $pos = n - 1$: $ci'(n - 1) = ci(n)$. This property is verified in `CycleIntegralFilterProperties::assertSameBeforeMerge`, `assertShiftAtMerge`, `assertShiftAfterMerge`, and `assertNewCIAAtSizeEqualsOld`. The full Scala verification code is in Appendix A.13.

6.8 Removing a Multiple

When the value at the merge point is a multiple of a filter value f , merging its two neighboring gaps removes it from the sequence entirely: the new integral's value at that position is the old integral's next value, which by construction is not itself a multiple.

$$\begin{aligned} \text{mod}(ci(0), f) \neq 0 \wedge \text{mod}(ci(m), f) = 0 &\implies ci'(m) = ci(m+1) \quad [\text{Multiple removed}] \\ \text{mod}(ci(m+1), f) \neq 0 &\implies \text{mod}(ci'(m), f) \neq 0 \quad [\text{Result is not a multiple}] \end{aligned}$$

Proof. The first identity is the merge shift law's "at merge" case (Subsection 6.7) applied directly: $ci'(m) = ci(m+1)$ regardless of whether $ci(m)$ was a multiple. The second follows because equal values satisfy the same modulo condition.

This property is verified in `CycleIntegralFilterProperties::assertRemoveOneMultiple` and `assertRemoveMultipleModNotZero`. The full Scala verification code is in Appendix A.14.

6.9 Direct Construction from Survivors

Rather than merging gaps one multiple at a time, a filtered cycle integral can be built directly from the survivor list (Subsection 6.5): its initial value is the first survivor, and its gap cycle is the list of differences between consecutive survivors. This direct construction reproduces exactly what repeated merging would produce.

$$\begin{aligned} S &:= \text{survivorValues}(ci, f, 0, n) \\ ci''\text{'s initial value} &= S_0, \quad \text{cycle}''(i) = S_{i+1} - S_i \\ \implies ci''(k) &= S_{k+1} \quad [\text{Direct construction matches survivors}] \end{aligned}$$

Proof. By induction on k . The base case follows directly from the integral's own definition applied to the first gap. The inductive step adds the k -th constructed gap, which by definition equals $S_{k+1} - S_k$, to the inductive hypothesis $ci''(k-1) = S_k$, giving $ci''(k) = S_{k+1}$.

This property is verified in `CycleIntegralFilterProperties::assertNewCIGeneratesFiltered`, `assertNewCIMatchesSurvivors`, and `assertGapsFromSurvivorsMatchCI`. The full Scala verification code is in Appendix A.15.

6.10 Filtered Result Has No Multiples

The cycle integral constructed directly from the survivor list contains no multiples of the filter value at any position — its values are, by construction, exactly the survivors.

$$\text{mod}(ci(0), f) \neq 0 \implies \text{mod}(ci''(k), f) \neq 0 \quad \text{for every valid } k$$

Proof. By Subsection 6.9, $ci''(k)$ equals the survivor S_{k+1} , and every element of the survivor list is already known not to be a multiple of f (Subsection 6.5).

$$\begin{aligned} ci''(k) &= S_{k+1} && [\S 6.9] \\ \text{mod}(S_{k+1}, f) &\neq 0 && [\S 6.5] \\ \therefore \text{mod}(ci''(k), f) &\neq 0. \quad \blacksquare && [\text{Q.E.D.}] \end{aligned}$$

This property is verified in `CycleIntegralFilterProperties::assertFilterMergeComposition` and `assertNextGapsValid`. The full Scala verification code is in Appendix A.16.

References

- [1] G. H. Hardy and E. M. Wright. *An Introduction to the Theory of Numbers*. Oxford University Press, 5th edition, 1979. §5.4 (Chinese Remainder Theorem), §15.1 (Sieve of Eratosthenes).
- [2] T. H. Mata. Division and Modulo from Recursive Normalization. <http://ai.viXra.org/abs/2609.0009>, 2026. ai.viXra:2609.0009.
- [3] T. H. Mata. Formal Verification of Cyclic Lists. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter4/cycle.md>, 2026.
- [4] T. H. Mata. Formal Verification of Discrete Integration Properties from First Principles. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter4/integral.md>, 2026.
- [5] T. H. Mata. Using Formal Verification to Prove Properties of Lists Recursively Defined. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter3/list.md>, 2026.
- [6] The Lean Community. Mathlib: Cycles of Lists. https://leanprover-community.github.io/mathlib4_docs/Mathlib/Data/List/Cycle.html.
- [7] The Lean Community. Mathlib: Periodic Functions. https://leanprover-community.github.io/mathlib4_docs/Mathlib/Algebra/Ring/Periodic.html.