

Structural Properties and Signed Boundaries of 2-Gaps in Sieve Sequences

Thiago Henrique Ramos da Mata

Independent Researcher

thiago.henrique.mata@gmail.com

<https://thiagomata.com/>

<https://github.com/thiagomata>

Proof status: The sequence foundation is Stainless-verified in the companion Sieve Sequence article. The signed-localization theorems introduced here are mathematically proved but not yet Stainless-verified.

Abstract

This article studies how 2-gaps evolve under successive prime filters and sharpens the boundary between complete-period survival and square-window placement. Complete-period CRT arguments prove that 2-gaps persist globally: an incoming odd prime removes exactly two copy classes of each old 2-gap. Those counts do not force a survivor into a particular interval $[Q, Q^2]$.

The full argument reduces survival to a single signed quantity. A weighted conservation law and its Cauchy–Schwarz corollary give a sharp terminal threshold on the harmful-excess energy. A closed exhaustion argument—capacity envelopes, native-period Bessel, fixed and moving cuts, and the stability-gap repair—shows that no unsigned-capacity route can clear that threshold. The new contribution is consequently signed and local. At filter 7, exact residue order gives the sharp interval bound $|b_7| \leq 18/7$, replacing a capacity estimate that grows quadratically with the window scale. More generally, when an old period is copied through an incoming prime r , the centered harmful excess in copy block j is exactly $B_j = d_t + d_{t-2}$ for two entries of the centered old-start histogram modulo r . Consequently,

$$\sum_{j=0}^{r-1} B_j^2 = 2V_r + 2 \sum_{t \bmod r} d_t d_{t-2} \leq 4V_r.$$

This composes residue energy with the weighted harmful-excess survival criterion over complete old-period blocks. It does not control the two partial boundary fragments of an arbitrary interval, nor does it prove a relative residue-energy estimate across an unbounded filter chain. Those are now the precise remaining arithmetic obligations, framed by a scale conflict that limits fixed-seed averaging and a Type-II barrier that any prime-producing sieve must overcome.

Licensed under the Creative Commons Attribution 4.0 International License (CC BY 4.0):

<https://creativecommons.org/licenses/by/4.0/>.

1 Introduction

The Sieve Sequence represents an infinite accepted-value stream using one finite cyclic gap list. That representation makes complete-period identities exact, but the twin-prime application asks for a surviving 2-gap start in the eligible square-safe window of one future head. The distinction between those scopes is the organizing principle of this article.

Version 1 established the copy-or-merge and complete-period/local-window boundary. This version retains that theorem spine and adds the signed localization results that emerged from the later quadratic investigation. The article develops the following properties in dependency order:

1. exact complete-period 2-gap count — Section 3.1;
2. exact copy-index filter frequency — Section 3.2;
3. exact batched survival — Section 3.3;
4. exact complete-period (2,4,2) cluster count — Section 3.4;
5. rotation invariance of cyclic counts — Section 3.5;
6. stable global absence of 2-gaps — Section 3.6;
7. square-safe certification — Section 4.1;
8. post-filter-3 isolation — Section 4.2;
9. exact accepted strikes and the sharp one-transition threshold — Sections 4.3–4.4;
10. why the capacity envelope is exhausted — Section 6;
11. exact weighted deletion conservation and its terminal quadratic corollary — Sections 5.1–5.2;
12. the exact filter-7 interval saving — Section 7;
13. the live frontier (accepted-boundary discrepancy and residue-collision energy estimates) — Section 8;
14. the copy-block residue-energy bridge — Section 9;
15. the classified routes and the live program — Section 10; and
16. the fixed-seed scale conflict (Section 10.1) and the Type-II barrier (Section 11.1).

The final sections separate the proved complete-block control from the open partial-boundary and cross-layer estimates. No complete-period theorem is used as a short-window lower bound.

2 Preliminaries And Evidence Boundary

For a prime stage head p , let

$$M_p = \prod_{q < p} q$$

and retain the integers coprime to M_p . Their accepted residues repeat modulo M_p ; adjacent differences in one complete period form a cyclic gap list. The verified companion article establishes accepted-value completeness, strict increase, period shift, exact survivor count, and the copy-or-merge transition rule.

This article takes those verified construction facts as inputs. It then studies mathematical properties of 2-gap populations and signed interval discrepancies. The division is deliberate: the companion article fixes and machine-checks the sieve object; this article expands the mathematical theory of that object without claiming that every new theorem has a Scala encoding.

2.1 Roles, Populations, And Scopes

The notation follows the shared research vocabulary:

- Q is a fixed **future head** used for square-safe certification;
- r is one **incoming prime**, and r_i is filter i in a complete conditioned chain to Q ;
- a **complete period** contains one full cyclic sieve pattern;
- a **local window** is an explicitly stated bounded integer interval;
- a **2-gap-start population** counts starts x , not all accepted values; and
- a **square-safe certificate** requires both x and $x + 2$ to lie strictly below Q^2 after every missing prime below Q has been installed.

For a conditioned chain, S_i denotes the local 2-gap starts immediately before filter r_i , and $N_i = |S_i|$. The final survivor population is S_m with size N_m . Complete-period populations are named separately and are never substituted for N_i without an explicit localization theorem.

2.2 Evidence Status

The following status convention is used:

- **Verified foundation:** supported by maintained Stainless source through the companion article.
- **Mathematically proved:** a complete mathematical proof is included here, but no corresponding `.holds` theorem currently exists.
- **Open:** the required estimate is stated but not proved.

These local labels map to the shared vocabulary as follows: **Verified foundation** means a Stainless-verified theorem, **Mathematically proved** means a mathematically proved theorem, and **Open** covers an open hypothesis or an unproved required estimate. The article's derivations and appendices are the authority for its mathematical claims; linked property records are supplementary navigation through the wider research program.

No theorem in this article claims infinitely many twin primes.

Each property section states its population, scope, and quantifier, then proves the result. Verified construction inputs link to maintained Scala contracts. A mathematical proof is not called Stainless-verified merely because its finite instances can be computed.

3 Complete-Period 2-Gap Properties

The first property group concerns cyclic 2-gap starts in complete periods. These identities are exact because every installed prime contributes a full residue system. They are not statements about a local square window.

3.1 Exact Global 2-Gap Count

After filter 2 is installed, every accepted value is odd, so accepted endpoints x and $x + 2$ are consecutive survivors: the intermediate value is even. For every prime stage after filter 2, this property counts all such cyclic 2-gap starts in one complete period of that stage, without constructing the gap list.

Let p be the stage head and

$$M_p = \prod_{q < p} q,$$

where q ranges over installed primes. A residue x represents a cyclic 2-gap exactly when

$$\gcd(x(x + 2), M_p) = 1.$$

The exact count is

$$G_2(p) = \prod_{\substack{3 \leq q < p \\ q \text{ prime}}} (q - 2).$$

For the filter 2, the start must be odd, leaving one allowed residue. For each installed odd prime q , the two endpoints fail precisely in the two classes

$$x \equiv 0 \pmod{q}, \quad x \equiv -2 \pmod{q}.$$

They are distinct because q is odd, so exactly $q - 2$ residues remain allowed at each installed odd prime q .

The installed primes are pairwise coprime, so CRT gives a bijection between one allowed choice at every installed prime and one residue modulo M_p . Consequently,

$$\begin{aligned} G_2(p) &= 1 \cdot \prod_{\substack{3 \leq q < p \\ q \text{ prime}}} (q - 2) \quad [\text{Chinese Remainder Theorem}] \\ &= \prod_{\substack{3 \leq q < p \\ q \text{ prime}}} (q - 2). \quad \blacksquare [\text{Q.E.D.}] \end{aligned}$$

The empty odd-prime product is 1, so the statement includes the first odd stage. The result proves complete-period global presence, not placement in any specified local window.

The derivation above proves the exact product count. A supplementary research record is available in [Exact Global 2-Gap Count](#). No `.holds` theorem currently encodes this exact product count, so this result is not yet Stainless-verified.

3.2 Exact Filter Frequency Across Repeated Copies

Expansion distributes every old 2-gap through equally spaced copies before filtering, and one incoming prime cannot choose arbitrary copies: the two endpoint strikes occur at two exact copy-index phases. This holds for lifted copies of one fixed old cyclic 2-gap, across every complete or finite run of copy indices satisfying the stated coprimality precondition.

Let M be the old period, let $(a, a + 2)$ be one old cyclic 2-gap, and let $r > 2$ be an incoming prime with $\gcd(M, r) = 1$. Copy j has endpoints

$$a + jM, \quad a + 2 + jM.$$

Let M^{-1} denote the inverse of M modulo r . The left endpoint is deleted exactly when

$$\begin{aligned} a + jM &\equiv 0 \pmod{r} && \text{[Left Endpoint Strike]} \\ j &\equiv -aM^{-1} \pmod{r}. && \text{[Multiply By } M^{-1}] \end{aligned}$$

Likewise, the right endpoint is deleted exactly when

$$\begin{aligned} a + 2 + jM &\equiv 0 \pmod{r} && \text{[Right Endpoint Strike]} \\ j &\equiv -(a + 2)M^{-1} \pmod{r}. && \text{[Multiply By } M^{-1}] \end{aligned}$$

If the two copy-index classes were equal, subtracting would give $2M^{-1} \equiv 0 \pmod{r}$, hence $2 \equiv 0 \pmod{r}$, impossible for $r > 2$. Therefore the classes are distinct.

Every residue class modulo r occurs at most $\lceil L/r \rceil$ times in a run of L consecutive copy indices. Thus

$$K_r(J) \leq 2 \left\lceil \frac{L}{r} \right\rceil.$$

For one complete run of r consecutive copy indices, each forbidden class occurs exactly once. Hence

$$K_r = 2, \quad G_{\text{survive}} = r - 2.$$

■ [Q.E.D.]

This is exact distribution across copies before and under one filter. It does not say that the $r - 2$ survivors are evenly distributed after filtering or that one lies in a chosen numerical window. The derivation above proves this exact copy-index law. A supplementary research record is available in [Exact Filter Frequency Across Repeated Copies](#). The repeated-stream foundation is verified in the companion Sieve Sequence article [1], but no `.holds` theorem currently packages these two exact copy-index classes and the finite-slice bound. This result is not yet Stainless-verified.

3.3 Exact Batched 2-Gap Survival

The one-filter copy law composes exactly. Applying several future filters as one batch counts intersections through CRT, so no survivor is subtracted twice and no intermediate floor or density approximation is needed. This holds for every finite set of distinct incoming odd primes coprime to the old period, over one complete period after the whole batch, for lifted copies of every old cyclic 2-gap in one complete combined period.

Let M be the old period and let $(a, a + 2)$ be one old cyclic 2-gap. Choose a finite set of distinct odd primes

$$\mathcal{R} = \{r_1, r_2, \dots, r_k\}, \quad \gcd\left(M, \prod_{r \in \mathcal{R}} r\right) = 1,$$

and define

$$R = \prod_{r \in \mathcal{R}} r.$$

The complete combined period modulo MR contains R lifted copies of the old pair. Exactly

$$\prod_{r \in \mathcal{R}} (r - 2)$$

survive every filter in the batch. Therefore, if the old complete period has G cyclic 2-gaps, the new complete period has

$$G_{\text{after}} = G \prod_{r \in \mathcal{R}} (r - 2).$$

For each $r \in \mathcal{R}$, the filter-frequency theorem (Section 3.2) identifies two distinct forbidden copy-index classes: one where r divides the left endpoint and one where it divides the right endpoint. Hence there are exactly $r - 2$ allowed choices modulo r . Because the primes in $\mathcal{R}$ are pairwise coprime, CRT makes these choices independent. Thus, for one old 2-gap,

$$G_{\text{one old gap}} = \prod_{r \in \mathcal{R}} (r - 2). \quad [\text{Chinese Remainder Theorem; Copy-Index Filter Frequency}]$$

The lifted copy sets belonging to distinct old cyclic starts are counted as distinct starts in the new complete period. Summing the same exact count over the G old starts gives

$$\begin{aligned} G_{\text{after}} &= \sum_{g=1}^G \prod_{r \in \mathcal{R}} (r - 2) \quad [\text{Sum Over Old 2-Gap Starts}] \\ &= G \prod_{r \in \mathcal{R}} (r - 2). \quad \blacksquare [\text{Simplification; Q.E.D.}] \end{aligned}$$

The formula is independent of the conceptual order of the filters and counts overlapping strikes correctly. Its scope is nevertheless the full combined period MR . A shorter interval can omit all allowed CRT classes, so this theorem cannot place a survivor in an eligible square-safe window. The derivation above proves the finite-batch formula and its complete-period limitation. A supplementary research record is available in [Exact Batched 2-Gap Survival](#). No corresponding .holds theorem currently packages the finite-batch product; this result is not yet Stainless-verified.

3.4 Exact Global (2,4,2) Cluster Count

The label (2,4,2) names the sequence of consecutive gaps inside a four-term cluster

$$a, \quad a + 2, \quad a + 6, \quad a + 8 :$$

gap 2 from a to $a + 2$, gap 4 from $a + 2$ to $a + 6$, gap 2 again from $a + 6$ to $a + 8$. This is the classic prime-quadruplet shape, realized for example by 11, 13, 17, 19 or 101, 103, 107, 109: two twin-prime pairs sharing their two middle values four apart. Counting how often this exact pattern survives is the natural next question once single 2-gaps (Section 3.1) are understood, since it asks whether 2-gaps keep arriving in tight pairs rather than only individually. It is also where the pattern comes from in the first place: at the earliest stage, right after filters 2 and 3 are installed, the entire accepted sequence already repeats as $\dots, 2, 4, 2, 4, 2, \dots$ forever, so (2,4,2) starts out as the universal shape of the sequence, not a rare event — this section tracks how much of that shape survives as larger primes filter it down.

The cluster's two 2-gaps are endpoint-disjoint and lie in a total span of 8. Expansion creates r copies of every old cluster; the incoming filter strikes exactly four copies and preserves the other $r - 4$ intact.

Let C_M be the cyclic cluster count at modulus M . Copy j of a cluster has endpoints

$$a + jM + \{0, 2, 6, 8\}, \quad 0 \leq j < r.$$

For each offset $h \in \{0, 2, 6, 8\}$, the endpoint $a + jM + h$ is removed exactly in the copy-index class

$$j \equiv -(a + h)M^{-1} \pmod{r}.$$

If two offsets produced the same class, their difference would be divisible by r . The nonzero pairwise differences are 2, 4, 6, 8. Neither 5 nor 7 divides any applicable difference, and every prime

$r \geq 11$ exceeds all four. Thus the four classes are distinct. Out of the r total copies created by expansion, let N_{struck} be the number of copies struck by the incoming filter (one per distinct endpoint class) and N_{intact} the number left intact. Then

$$\begin{aligned} N_{\text{struck}} &= 4 && \text{[Four Distinct Endpoint Classes]} \\ N_{\text{intact}} &= r - N_{\text{struck}} = r - 4. && \text{[Subtraction]} \end{aligned}$$

It remains to exclude newly created clusters. After filter 2, all old gaps are positive and even. A merged gap is a sum of at least two old gaps, so it cannot equal 2. It can equal 4 only as $2 + 2$. But two consecutive 2-gaps would require all three values $x, x + 2, x + 4$ to be accepted, whereas one of them is divisible by 3. Since filter 3 is installed, that is impossible. Therefore every new gap of size 2 or 4 is a copied old gap, and every new $(2, 4, 2)$ occurrence is one of the intact copies counted above. Hence

$$C_{rM} = (r - 4)C_M.$$

■ [No New Occurrences; Q.E.D.]

This recurrence holds for every complete-period stage with modulus M divisible by 6 and every incoming prime $r \geq 5$ with $r \nmid M$, counting cyclic occurrences of the gap word $(2, 4, 2)$ in one complete period after filters 2 and 3 are installed.

The wheel modulo 6 has cyclic gap word $(4, 2)$: accepted residues $1, 5 \pmod{6}$ repeat forever as $\dots, 4, 2, 4, 2, 4, 2, \dots$ before any prime beyond 3 is installed. At this base stage the sequence is *nothing but* copies of $(2, 4, 2)$ back to back, so there is exactly one cyclic $(2, 4, 2)$ occurrence per period, $C_6 = 1$ — the recurrence below tracks how that initially-universal pattern gets diluted, but never destroyed, as larger primes are installed. Iterating the recurrence over a finite installed-prime set $\mathcal{P}$ containing 2 and 3 gives

$$C(\mathcal{P}) = \prod_{\substack{p \in \mathcal{P} \\ p \geq 5}} (p - 4).$$

The absolute cluster population is positive at every stage and grows whenever an incoming prime exceeds 5. Its proportion among accepted positions is multiplied by $(r - 4)/(r - 1) < 1$, so global growth does not imply that a chosen short window contains a cluster.

The derivation above establishes the recurrence, no-creation result, closed product, and localization boundary. A supplementary research record is available in [Exact Global Count Of \$\(2, 4, 2\)\$ Two-Gap Clusters](#). No .holds theorem currently packages the cyclic cluster count; this result is not yet Stainless-verified.

3.5 Rotation Preserves Cyclic Gap Counts

Rotation chooses a new origin for the same cyclic list. It neither filters an accepted value nor merges adjacent gaps, so it preserves the number of entries having every gap value, including 2, for every nonempty finite cyclic gap list, every nonnegative rotation offset, and every gap value d . The rotation invariance itself is proved; the exact multiplicity claim is not yet Stainless-verified.

Let

$$G = (g_0, g_1, \dots, g_{T-1}), \quad T \geq 1,$$

and define rotation by offset j through

$$\text{rot}_j(G)_i = g_{(i+j) \bmod T}.$$

For every value d ,

$$\#\{i : \text{rot}_j(G)_i = d\} = \#\{i : g_i = d\}.$$

Define the index map $\varphi_j(i) = (i + j) \bmod T$. Addition by j modulo T is invertible, with inverse $\varphi_{-j}(i) = (i - j) \bmod T$. Hence φ_j is a bijection of $\{0, 1, \dots, T - 1\}$. Therefore

$$\begin{aligned} \#\{i : \text{rot}_j(G)_i = d\} &= \#\{i : g_{\varphi_j(i)} = d\} && \text{[By Definition Of Rotation]} \\ &= \#\{k : g_k = d\} && \text{[Reindex By The Bijection } k = \varphi_j(i)\text{]} \\ &= \#\{i : g_i = d\}. && \blacksquare \text{ [Rename The Bound Index; Q.E.D.]} \end{aligned}$$

Taking $d = 2$ proves exact invariance of the complete cyclic 2-gap count. Rotation may change which gap follows the displayed head and whether one gap crosses the end-to-start boundary of a linear rendering. It cannot imply that an absolute coordinate window such as $[Q, Q^2)$ contains the same number of 2-gaps, because that window is not defined only by cyclic indices.

Scala Verification Foundation And Pending Multiplicity Theorem

The maintained list implementation defines rotation as `back ++ front`. Its verified properties preserve membership in both directions and preserve list size:

```
def assertRotateContainsForward(
  list: List[BigInt], index: BigInt, x: BigInt
): Boolean = {
  require(index >= 0)
  require(list.contains(x))
  // ... verified proof ...
  ListUtils.rotateAt(list, index).contains(x)
}.holds

def assertRotateContainsBackward(
  list: List[BigInt], index: BigInt, x: BigInt
): Boolean = {
  require(index >= 0)
  require(ListUtils.rotateAt(list, index).contains(x))
  // ... verified proof ...
  list.contains(x)
}.holds

def assertRotateSameSize(
  list: List[BigInt], index: BigInt
): Boolean = {
  require(index >= 0)
  // ... verified proof ...
  ListUtils.rotateAt(list, index).size == list.size
}.holds
```

These foundations are verified in [RotationProperties::assertRotateContainsForward](#), [assertRotateContainsBackward](#), and [assertRotateSameSize](#). They establish the rotation operation and its same-elements/size behavior, but membership alone does not count duplicate entries. The exact multiplicity theorem above is maintained mathematically in [Rotation Preserves Cyclic Gap Counts](#); it is not yet Stainless-verified.

3.6 Absence Of 2-Gaps Is Stable

Later filtering can copy an old gap or merge consecutive old gaps, but it cannot create a smaller positive gap. Thus, once the complete cyclic population contains no 2-gap, no later filter can recreate one — a fact that holds for every later filter transition, and therefore every finite chain of later transitions, over every gap in one complete cyclic post-filter-2 gap list.

Let the old cyclic gaps be $g_0, \dots, g_{T-1}$. Because filter 2 is already installed, every g_i is positive and even. Under the no-2 hypothesis,

$$\forall i, \quad g_i \neq 2 \implies g_i \geq 4.$$

For two values that remain consecutive after the next filter, their new gap h has one of two forms. If no intermediate accepted value was removed, it copies one old gap. If one or more intermediate accepted values were removed, it is the sum of at least two consecutive old gaps. Therefore

$$h = g_i \implies h \geq 4 \quad [\text{Copied Gap}]$$

$$h = \sum_{j=u}^v g_j, \quad v \geq u + 1 \implies h \geq 4 + 4 = 8. \quad [\text{Merged Gaps}]$$

In either case $h \neq 2$, so

$$(\forall i, g_i \neq 2) \implies (\forall j, h_j \neq 2).$$

■ [Copy-Or-Merge Exhaustion; Q.E.D.]

Applying the same implication inductively yields

$$G_s(2) = 0 \implies G_t(2) = 0 \quad \text{for every later stage } t \geq s,$$

where $G_s(2)$ is the complete-period 2-gap count at stage s . This is a one-way extinction theorem. A positive global count does not force a 2-gap into a chosen short window, so it must not be used as a localization result.

The copy-or-merge argument above proves the inductive consequence and its global/local boundary. A supplementary research record is available in [Absence Of 2-Gaps Is Stable Under Later Filtering](#). No dedicated `.holds` theorem currently quantifies over the complete cyclic gap transition, so this result is not yet Stainless-verified.

These are complete-period statements. They explain global growth but do not locate any surviving copy in a prescribed short interval.

The verified Scala representation of the underlying repetition, filtering, and next-stage reconstruction is documented in [1]. This article does not duplicate those maintained source proofs.

Sections 3.1–3.6 above are exact complete-period identities; none of them measure a short local window, and Section 3.6 explicitly warns against reading a positive global count as a local guarantee. The figure below shows the quantity those theorems deliberately do not bound: the fraction of one stage’s own gaps that equal 2, per stage, over a real generated dataset (200 stages, the first 100,000 gaps of each). It falls sharply from 100% (stage 1, where every odd-only gap is 2) toward roughly 10% by head ≈ 1200 — consistent with $G_2(p)$ ’s product formula thinning the *complete-period* count, but shown here at the *local, per-stage* scale that Section 3’s theorems are careful not to claim.

The figure above summarizes each stage to one number; the figure below shows the same underlying data at full per-row detail, over the same 200-stage dataset. Every 2-gap renders in a fixed green; everything between one 2-gap and the next is collapsed into a single colored unit representing that distance. A row that is mostly green is a stage whose 2-gaps are still nearly back to back; a row with long colored runs is a stage where 2-gaps have spread apart. This is the row-by-row texture behind the declining-frequency curve above, not a new claim.




Figure 1: Fraction of a stage's gaps equal to 2, per stage, declining from 100% toward roughly 10%.




Figure 2: 2-focused compression heatmap: every 2-gap in green, the collapsed distance to the next one colored by frequency, one row per stage.

4 Local Certification And One-Transition Survival

Complete-period growth becomes relevant to twin primes only after a theorem places a surviving 2-gap inside an eligible square-safe window. This section first proves what such a survivor certifies, then isolates the exact local attrition conditions needed to retain one.

4.1 Safe-Window 2-Gaps Certify Twin Primes

The square bound turns acceptance into primality. Any composite below Q^2 has a prime divisor below Q , but every such divisor has already been installed as a filter. Therefore, for every prime future head Q and every accepted integer n with $Q \leq n < Q^2$ after all primes below Q are installed, a surviving accepted pair at distance 2 is not merely a putative pair; it is a twin-prime pair.

Let

$$P_Q = \prod_{r < Q} r,$$

where r ranges over primes. If

$$Q \leq n < Q^2, \quad \gcd(n, P_Q) = 1,$$

then n is prime. Indeed, suppose that n were composite. It has a prime divisor $r \leq \sqrt{n}$. The strict square bound gives $r \leq \sqrt{n} < Q$, so r is one of the factors of P_Q . Consequently $r \mid n$ and $r \mid P_Q$, contradicting $\gcd(n, P_Q) = 1$:

$$\begin{aligned} n \text{ composite} &\implies \exists r \text{ prime} : r \mid n \wedge r \leq \sqrt{n} && \text{[Small Prime Divisor]} \\ &\implies r < Q && \text{[Strict Bound } n < Q^2\text{]} \\ &\implies r \mid P_Q && \text{[By Definition]} \\ &\implies \gcd(n, P_Q) > 1 && \text{[Since } r \mid n\text{]} \\ &\implies \perp. && \text{[Contradiction]} \end{aligned}$$

Hence n is prime. Applying this result independently to both endpoints gives

$$\begin{aligned} Q \leq x, \quad x + 2 < Q^2, \quad \gcd(x(x + 2), P_Q) = 1 \\ \implies x \text{ and } x + 2 \text{ are prime} \quad \blacksquare \text{ [Q.E.D.]} \end{aligned}$$

The right-endpoint inequality must be strict: Q^2 is composite but has no prime divisor below Q . The theorem certifies one eligible survivor; it does not prove that any such survivor exists. The derivation above proves the theorem and its endpoint discipline. A supplementary research record is available in [Safe-Window 2-Gaps Certify Twin Primes](#). No `.holds` theorem currently encodes the least-prime-divisor argument, so this result is not yet Stainless-verified.

4.2 Isolation Of 2-Gaps After Filter 3

After filter 3, two 2-gaps cannot share an endpoint, for every sieve stage whose modulus M is divisible by 6. This improves the sharp destruction capacity of one later filter strike: for every later deletion of an accepted value, removing one accepted value can destroy at most one existing 2-gap.

Suppose both $(x, x + 2)$ and $(x + 2, x + 4)$ were accepted 2-gaps. The three values $x, x + 2, x + 4$ occupy all three residue classes modulo 3, so exactly one is divisible by 3. Because $3 \mid M$, that value

cannot be accepted. Thus

$$\begin{aligned}
& (x, x+2) \text{ and } (x+2, x+4) \text{ both accepted} \\
& \implies 3 \nmid x, 3 \nmid (x+2), 3 \nmid (x+4) && [\text{Acceptance}] \\
& \implies \perp && [\text{Complete Residue System Modulo 3}].
\end{aligned}$$

Equivalently, parity and filter 3 force every 2-gap start into one residue:

$$\begin{aligned}
x, x+2 \text{ accepted} &\implies x \equiv 1 \pmod{2} && [\text{Filter 2}] \\
&\implies x \equiv 5 \pmod{6}. && [\text{Filter 3; CRT}]
\end{aligned}$$

Any destroyed 2-gap must contain a removed accepted value as an endpoint. A value could belong to two 2-gaps only in the forbidden overlapping configuration above. Let $N_{\text{destroyed}}$ be the number of destroyed 2-gaps and N_{removed} the number of removed accepted values. Therefore

$$N_{\text{destroyed}} \leq N_{\text{removed}}.$$

■ [Q.E.D.]

Isolation limits destruction efficiency. It does not imply that a chosen window contains a 2-gap before or after the filter.

The isolation argument above forces $x \equiv 5 \pmod{6}$: the smallest gap that can separate two consecutive 2-gaps is therefore the classic prime-quadruplet pattern $[2, 4, 2]$, a distance of exactly 4, never less. The figure below measures this directly on the same 200-stage generated dataset used in Section 3: the average and maximum distance between consecutive 2-gaps (summed runs, not individual raw gaps), per stage. The average grows steadily ($4 \rightarrow \sim 125$) while the maximum grows much faster (~ 1450 by the last stage), diverging further as the head grows — but the average’s floor sits at exactly 4 across every stage, never lower, matching the isolation bound proved above rather than merely approximating it.




Figure 3: Average and maximum distance between consecutive 2-gaps, per stage; the average floor is exactly 4.

The overlap argument above proves its filtering consequence. A supplementary research record is available in [Isolation Of 2-Gaps After Filtering By 3](#). No dedicated `.holds` theorem currently counts incident 2-gaps per accepted endpoint, so this result is not yet Stainless-verified.

4.3 Exact Accepted Local Filter Strikes

Counting every multiple of r overstates local destruction because most multiples have already been removed by smaller filters. For every incoming prime $r \geq 5$ and its next prime future head Q , the remaining multiples in that square-safe window admit an exact prime-multiplier description, proved here using Bertrand's postulate as an external dependency.

Let

$$M_r = \prod_{s < r} s, \quad K = \left\lfloor \frac{Q^2 - 1}{r} \right\rfloor,$$

where s ranges over primes and Q is the next prime after r . Before filter r is applied, the accepted multiples of r in $[Q, Q^2)$ are exactly

$$r\ell \quad \text{with } \ell \text{ prime and } r \leq \ell \leq K.$$

Consequently their number is

$$A(r, Q) = \pi(K) - \pi(r - 1).$$

Every multiple of r in the window has the form $r\ell$ with $2 \leq \ell \leq K$, because Bertrand's postulate gives $r < Q < 2r$. Since $\gcd(r, M_r) = 1$, this value survived the earlier filters exactly when $\gcd(\ell, M_r) = 1$.

Bertrand's bound also gives

$$\begin{aligned} K &< \frac{Q^2}{r} && \text{[By Definition Of } K\text{]} \\ &< 4r && \text{[Bertrand: } Q < 2r\text{]} \\ &\leq r^2. && \text{[Since } r \geq 5\text{]} \end{aligned}$$

If an accepted multiplier $\ell < K + 1 \leq r^2$ were composite, its least prime divisor would be at most $\sqrt{\ell} < r$ and would divide M_r , contradicting $\gcd(\ell, M_r) = 1$. Thus every accepted multiplier is prime and at least r . Conversely, every prime $\ell \geq r$ is coprime to M_r . Therefore

$$\begin{aligned} &\#\{n \in [Q, Q^2) : r \mid n, \gcd(n, M_r) = 1\} \\ &= \#\{\ell : r \leq \ell \leq K, \ell \text{ prime}\} && \text{[Accepted Multiplier Characterization]} \\ &= \pi(K) - \pi(r - 1) && \text{[Prime-Counting Definition]} \\ &= A(r, Q). && \blacksquare \text{ [Q.E.D.]} \end{aligned}$$

This is the exact count of accepted values struck by filter r in the stated window. A struck value need not be a 2-gap endpoint, so $A(r, Q)$ is only an upper bound on the number of destroyed local 2-gaps.

The derivation above proves the exact characterization, including its Bertrand dependency. A supplementary research record is available in [Exact Accepted Local Filter Strikes](#). No `.holds` theorem currently contains the prime-counting argument; this result is not yet Stainless-verified.

4.4 Sharp Local 2-Gap Survival Threshold

The proved theorem is the conditional implication below; its local-abundance antecedent remains open. It holds for every incoming prime $r \geq 5$, its next prime future head Q , and the single transition that installs filter r , over pre-filter and post-filter 2-gap starts whose two endpoints remain inside one eligible square-safe window.

The exact accepted-strike count becomes a sharp deterministic survival test. If the eligible window initially contains more 2-gaps than filter r has accepted values to remove, endpoint isolation forces at least one gap to survive.

Let $G_{\text{local}}(r, Q)$ count the pre-filter 2-gaps $(x, x + 2)$ satisfying

$$Q \leq x, \quad x + 2 < Q^2,$$

and let $G_{\text{surviving}}(r, Q)$ count those still present after filter r . With

$$K = \left\lfloor \frac{Q^2 - 1}{r} \right\rfloor, \quad A(r, Q) = \pi(K) - \pi(r - 1),$$

let $G_{\text{destroyed}}(r, Q)$ count the destroyed eligible 2-gaps. The 2-gap isolation and accepted local strikes properties give

$$\begin{aligned} G_{\text{destroyed}}(r, Q) &\leq \#\{\text{accepted values struck in } [Q, Q^2)\} && \text{[2-Gap Isolation]} \\ &= A(r, Q). && \text{[Accepted Local Strikes]} \end{aligned}$$

Therefore

$$\begin{aligned} G_{\text{surviving}}(r, Q) &= G_{\text{local}}(r, Q) - G_{\text{destroyed}}(r, Q) && \text{[Population Accounting]} \\ &\geq G_{\text{local}}(r, Q) - A(r, Q). && \text{[Substitution]} \end{aligned}$$

Both counts are integers, so

$$G_{\text{local}}(r, Q) > A(r, Q) \implies G_{\text{surviving}}(r, Q) > 0 \quad \blacksquare \text{ [Integer Positivity; Q.E.D.]}$$

Equivalently, $A(r, Q) + 1$ eligible pre-filter gaps suffice. The theorem does not prove this local-abundance antecedent. Iterating it through later filters would require a fresh eligible population bound at every transition.

The derivation above proves the conditional theorem and its exact boundary. A supplementary research record is available in [Sharp Local 2-Gap Survival Threshold](#). No `.holds` theorem currently encodes the local populations or the prime-counting threshold; this result is not yet Stainless-verified.

Local Harmful-Excess Notation

Let Q be a future prime head and define the eligible start window

$$W_Q = \{x \in \mathbb{Z} : Q \leq x \wedge x + 2 < Q^2\}.$$

For a fixed filter r , let $K_r(I)$ count the incoming 2-gap starts in an interval I destroyed because r divides one endpoint. If $N_r(I)$ is the incoming population, define its centered harmful excess by

$$b_r(I) = K_r(I) - \frac{2N_r(I)}{r}.$$

The density term $2N_r(I)/r$ is not itself an upper bound. The sign and size of $b_r(I)$ encode the interval-order information discarded by separate capacity estimates.

References

- [1] T. H. Mata. Formal Verification of Sieve Sequence Stages and Their Transitions. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter6/sieve-sequence.md>, 2026.