

Formal Verification of Cycle Integral Properties from First Principles

Thiago Henrique Ramos da Mata

Independent Researcher

thiago.henrique.mata@gmail.com

<https://thiagomata.com/>

<https://github.com/thiagomata>

Abstract

In previous articles, we defined bounded Lists, Integrals of Lists, and unbounded Cycles of Integers from scratch, relying only on core type constructs and recursion, with no prior knowledge of Scala’s collections required. From that, we proved and formally verified some properties related to them. This article uses that as a foundation to define Integral of Cycles using two presentations: the canonical recursive `CycleIntegral` definition and a `ModCycleIntegral` closed-form definition. For both presentations, we formally verify the sum property (integral equals cumulative cycle sum) and the step property (difference between consecutive values equals the corresponding cycle element) using the Stainless verification system. We also prove that the recursive and modulo definitions are extensionally equivalent. All properties are expressed and proved within a minimal framework using only elementary arithmetic, recursion, and pure Scala code. This work bridges mathematical foundations and executable verification, offering a self-contained, verifiable approach for reasoning about infinite periodic accumulations.

Licensed under the Creative Commons Attribution 4.0 International License (CC BY 4.0):

<https://creativecommons.org/licenses/by/4.0/>.

1 Introduction

Cycles are a powerful concept in computer science and mathematics, representing unbounded lists that repeat a finite sequence of elements. When we integrate cycles, we obtain a list of cumulative sums with some unique properties that we will explore in this article.

$$L = [l_0, l_1, l_2, \dots, l_{n-1}] \mid l_n \in \mathbb{S}, L \in \mathbb{L}$$

$$\begin{aligned} \text{Cycle}(L) &= [l_0, l_1, l_2, \dots, l_{n-1}, l_0, l_1, \dots] \\ &= [v_0, v_1, v_2, \dots] \mid v_i = L[i \bmod n] \end{aligned}$$

$$\text{Integral}(L, \text{init}) = [y_0, y_1, y_2, \dots, y_{n-1}] \mid y_k = \sum_{i=0}^k x_i + \text{init}$$

$$\text{CycleIntegral}(L, \text{init}) = [w_0, w_1, w_2, \dots] \mid w_k = \sum_{i=0}^k \text{Cycle}(L)_i + \text{init}$$

In this article, we present a discrete definition of Cycle Integral over finite integer lists, defined recursively and verified some of its properties using the Stainless system. Our approach follows a zero-prior-knowledge philosophy, building on a previously verified foundation for recursive list and

integral structures and summation. The result is a verified, from-scratch implementation of the cycle integral suitable as a foundation for higher-level numeric reasoning over unbounded lists.

This article verifies:

- Two equivalent definitions: recursive and modulo-based — Subsections 3.1–3.3
- Core properties: next position, same difference after cycle, sum of mod values, strictly increasing, positivity — Subsections 4.1–4.5
- Persistent and periodic properties: how a fixed cycle integral’s residues behave forever, and how it advances across full periods — Subsections 5.1–5.6
- Deriving new cycle integrals: expansion, index shifts, rotation, survivor filtering, and merge-based reconstruction — Subsections 6.1–6.10

1.1 Related work

Lean’s Mathlib provides a general formal theory of periodic functions. It proves that a periodic function remains periodic under integer multiples of a period, and that a finite sum of periodic functions is periodic [6]. Mathlib also represents a finite cycle as a list modulo cyclic rotation [5]. These results give formal context for the finite period and shift structure used by the present construction.

This article studies a different object: the cumulative integral of a concrete periodic integer list. Its central verified results are the equivalence of a recursive integral and a quotient–remainder closed form, together with the resulting step, full-period, residue, and reconstruction properties. The existing periodicity and finite-cycle developments therefore enrich the setting without standing in for this two-presentation cycle-integral proof.

2 Preliminaries

We reuse several basic list, cycle and integral operations and their verified properties from the companion articles *Using Formal Verification to Prove Properties of Lists Recursively Defined* [4], *Formal Verification of Discrete Integration Properties from First Principles* [3], and *Formal Verification of Cyclic Lists* [2]. We also reuse some modulo properties previously defined and verified in the article *Division and Modulo from Recursive Normalization* [1].

These articles also defined and verified their properties using the same zero-prior-knowledge methodology, and are treated here as foundational primitives.

3 Cycle Integral Definitions

The cycle integral extends the finite integral to unbounded repeating sequences. Two equivalent definitions are proven.

- Recursive: recurrence on the cycle position — Subsection 3.1
- Modulo: closed-form using `div` and `mod` — Subsection 3.2
- The two definitions are extensionally equivalent — Subsection 3.3

$$\forall i \in \mathbb{N}_0, \text{init} \in \mathbb{N}_0, L \in \mathbb{L}, n = |L|$$

$$\text{CycleIntegral}(L, \text{init}) := [w_0, w_1, w_2, \dots]$$

$$0 \leq i < n \implies w_i = \sum_{j=0}^i L_j + \text{init} \quad [\text{Sum Property}]$$

$$i > 0 \implies w_i - w_{i-1} = L_{(i \bmod n)} \quad [\text{Step Property}]$$

3.1 Recursive Cycle Integral

$$\text{RecCycle}(L) = [v_0, v_1, \dots] \mid v_i = \begin{cases} L[i] & i < n \\ v_{i-n} & i \geq n \end{cases}$$

$$\text{RecCycleIntegral}(L, \text{init}) = [w_0, w_1, \dots] \mid w_i = \begin{cases} \text{init} + w_0 & i = 0 \\ v_i + w_{i-1} & i > 0 \end{cases}$$

Recursive Cycle Equivalence: Recursive Cycle is equivalent to Mod Cycle, as proven in the article [Formal Verification of Cyclic Lists](#) [2].

$$\begin{aligned} \text{RecCycle}(L)_i &= \text{ModCycle}(L)_i && [\text{Cycle Equivalence}] \\ &= L_{(i \bmod n)} && [\text{Mod Cycle Definition}] \end{aligned}$$

Sum Property:

$$v_0 = L_0 \quad [\text{Base Case}]$$

$$= L_{(0 \bmod n)} \quad [\text{By Modulo Property}]$$

$$= \sum_{j=0}^0 L_j \quad [\text{Summation Re-indexing}]$$

$$i < n \implies$$

$$w_0 = \text{init} + v_0 \quad [\text{Base Case}]$$

$$= \text{init} + \sum_{j=0}^i L_j \quad [\text{Summation Re-indexing}]$$

$$i > 0 \implies$$

$$w_i = \text{init} + v_0 + \sum_{j=1}^i v_j \quad [\text{By Definition}]$$

$$= \text{init} + v_0 + \sum_{j=1}^i L_j \quad [\text{Mod of Small Values Property}]$$

$$= \text{init} + \sum_{j=0}^i v_j \quad [\text{Summation Re-indexing}]$$

$$\therefore w_i = \text{init} + \sum_{j=0}^i L_j \quad [\text{Q.E.D.}]$$

This property is verified in the [CycleIntegralProperties::assertCycleIntegralEqualsSumSmallPositions](#). A key Scala verification excerpt is in Appendix A.1; the complete proof is linked in the source reference.

Step Property:

$$\begin{aligned}
 w_i - w_{i-1} &= v_i + w_{i-1} - w_{i-1} && \text{[By Definition]} \\
 &= v_i && \text{[Simplification]} \\
 &= L_{(i \bmod n)} && \text{[Substitution]}
 \end{aligned}$$

This property is verified in the [CycleIntegralProperties::assertDiffEqualsCycleValue](#). A key Scala verification excerpt is in Appendix A.2; the complete proof is linked in the source reference.

3.2 Modulo Cycle Integral

$$\begin{aligned}
 \text{ModCycle}(L)_i &:= L_{(i \bmod n)} = [w_0, w_1, \dots] \\
 I_k &:= \sum_{j=0}^k L_j \quad (0 \leq k < n) && \text{[Integral of L]} \\
 S &:= I_{n-1} && \text{[One full cycle sum]} \\
 \text{ModCycleIntegral}(L, \text{init})_i &:= (i \text{ div } n) \cdot S + I_{(i \bmod n)} + \text{init}
 \end{aligned}$$

Sum Property:

$$\begin{aligned}
 i < n &\implies w_i = \sum_{j=0}^i L_j + \text{init} && \text{[Claim to Prove]} \\
 i < n &\implies \\
 i \text{ div } n &= 0 && \text{[By Div of Small Values Property]} \\
 w_i &= (i \text{ div } n) \cdot S + I_{(i \bmod n)} + \text{init} && \text{[Definition]} \\
 &= 0 \cdot S + I_i + \text{init} && \text{[Substitution]} \\
 &= \sum_{j=0}^i L_j + \text{init} && \text{[By definition of } I_i\text{]} \\
 \therefore \forall i < n, \quad w_i &= \sum_{j=0}^i L_j + \text{init} && \text{[Q.E.D.]}
 \end{aligned}$$

This property is verified in the [ModCycleIntegralProperties::assertFirstValuesMatchIntegral](#). A key Scala verification excerpt is in Appendix A.3; the complete proof is linked in the source reference.

Step Property:

$$w_i - w_{i-1} = L_{i \bmod n}, \quad i > 0, n > 0 \quad \text{[Claim to Prove]}$$

Case $i \bmod n > 0$:

$$\begin{aligned}
i \bmod n &= ((i-1) \bmod n) + 1 && \text{[By Modulo Properties]} \\
i \operatorname{div} n &= (i-1) \operatorname{div} n && \text{[By Division Properties]} \\
w_i &= (i \operatorname{div} n) S + I_{i \bmod n} + \textit{init} && \text{[Definition]} \\
&= (i-1 \operatorname{div} n) S + I_{((i-1) \bmod n)+1} + \textit{init} && \text{[Div/Mod property]} \\
w_{i-1} &= (i-1 \operatorname{div} n) S + I_{(i-1) \bmod n} + \textit{init} && \text{[Definition]} \\
w_i - w_{i-1} &= I_{((i-1) \bmod n)+1} - I_{(i-1) \bmod n} && \text{[Cancellation]} \\
&= \left(\sum_{j=0}^{(i-1) \bmod n} L_j + L_{((i-1) \bmod n)+1} \right) - \sum_{j=0}^{(i-1) \bmod n} L_j && \text{[Expand sum]} \\
&= L_{((i-1) \bmod n)+1} && \text{[Cancellation]} \\
&= L_{i \bmod n} && \text{[Modulo property]}.
\end{aligned}$$

Case $i \bmod n = 0$:

$$\begin{aligned}
w_i &= (i \operatorname{div} n) S + L_0 + \textit{init} && \text{[Definition]} \\
w_{i-1} &= (i \operatorname{div} n - 1) S + I_{n-1} + \textit{init} && \text{[Div/Mod property]} \\
w_i - w_{i-1} &= (i \operatorname{div} n) S + L_0 + \textit{init} - ((i \operatorname{div} n - 1) S + I_{n-1} + \textit{init}) && \text{[Substitution]} \\
&= S + L_0 - I_{n-1} && \text{[Simplification]} \\
&= L_0 && \text{[Since } S = I_{n-1}\text{]} \\
&= L_{i \bmod n} && \text{[Modulo property]}.
\end{aligned}$$

$$\therefore w_i - w_{i-1} = L_{i \bmod n}, \quad \forall i > 0 \quad \text{[Q.E.D.]}$$

This property is verified in the `ModCycleIntegralProperties::assertSimplifiedDiffValuesMatchCycle`. A key Scala verification excerpt is in Appendix A.4; the complete proof is linked in the source reference.

3.3 Equivalence of Definitions

The nontrivial equivalence proved here is between the recursive `CycleIntegral` definition and the closed-form `ModCycleIntegral` definition.

$$\begin{aligned}
\text{CycleIntegral}(L, \textit{init}) &= \text{ModCycleIntegral}(L, \textit{init}) \\
&= [w_0, w_1, w_2, \dots] \mid w_i = \sum_{j=0}^i L_{(j \bmod n)} + \textit{init} \blacksquare
\end{aligned}$$

This property is verified in the `ModCycleIntegralProperties::assertCycleIntegralMatchModCycleDef`. A key Scala verification excerpt is in Appendix A.5; the complete proof is linked in the source reference.

4 Core Verified Properties

The fundamental properties of the cycle integral that hold for every position.

- Next position: $CI_{i+1} = CI_i + \text{Cycle}(L)_{i+1}$ — Subsection 4.1
- Full cycle shift: adding one cycle period advances by the total sum — Subsection 4.2
- Sum of mod values: the modulo definition matches the list sum — Subsection 4.3
- Strictly increasing: positive base values force a strictly growing integral — Subsection 4.4
- Positivity: a non-negative start and positive base values keep the integral positive everywhere — Subsection 4.5

4.1 Next Position

For any positive position, the cycle integral at that position equals the previous value plus the current cycle element.

$$\forall i > 0 : CI(L, init)_i = CI(L, init)_{i-1} + \text{Cycle}(L)_i$$

This follows directly from the recursive definition of `CycleIntegral`.

This property is verified in the `CycleIntegralProperties::assertNextPosition`.

4.2 Same Difference After Full Cycle

The difference between consecutive values is invariant under adding a full cycle size to both positions.

$$\forall i \geq 0 : CI(L, init)_{i+1} - CI(L, init)_i = CI(L, init)_{i+size+1} - CI(L, init)_{i+size}$$

This property is verified in the `CycleIntegralProperties::assertSameDiffAfterCycle`. A key Scala verification excerpt is in Appendix A.6; the complete proof is linked in the source reference.

4.3 Sum of Mod Values as List

The cycle integral at any position equals the sum of a constructed list containing the initial value and all cycle values up to that position (using modular indexing for positions beyond one cycle).

$$\forall i \geq 0 : CI(L, init)_i = \text{sum}([init] \\ + [\text{Cycle}(L)_0, \text{Cycle}(L)_1, \dots, \text{Cycle}(L)_i])$$

This property is verified in the `CycleIntegralProperties::assertSumModValueAsListEqualsCycleIntegralLoop`. A key Scala verification excerpt is in Appendix A.7; the complete proof is linked in the source reference.

4.4 Cycle Integral Strictly Increasing

When the initial value is non-negative and every base-list value is positive, the cycle integral is strictly increasing: a later position always produces a larger value.

$$init \geq 0 \wedge (\forall x \in L, x > 0) \wedge b > a \implies \text{CycleIntegral}(L, init)_b > \text{CycleIntegral}(L, init)_a$$

Proof. Induct on $b - a$. The base case is one positive step; the inductive step adds one more positive cycle value.

$$\begin{aligned}
b = a + 1 &\implies CI_b - CI_a = \text{Cycle}(L)_b > 0 \\
&\quad [\S 3.1 \text{ and cycle positivity}] \\
&\implies CI_b > CI_a, \\
CI_{b-1} > CI_a, \quad CI_b - CI_{b-1} = \text{Cycle}(L)_b > 0 &\implies CI_b > CI_{b-1} > CI_a \\
&\therefore CI_b > CI_a. \quad \blacksquare [\text{Q.E.D.}]
\end{aligned}$$

This property is verified in the `CycleIntegralProperties::assertCycleIntegralIncreasing`.

4.5 Cycle Integral Positivity

When the initial value is non-negative and every base-list value is positive, the cycle integral is positive at every position.

$$init \geq 0 \wedge (\forall x \in L, x > 0) \implies \text{CycleIntegral}(L, init)_i > 0$$

Proof. At the first position, the non-negative initial value and a positive cycle value give a positive integral. Every later position adds one more positive cycle value.

$$\begin{aligned}
CI_0 &= init + \text{Cycle}(L)_0 > 0, \\
CI_{i-1} > 0, \quad CI_i - CI_{i-1} = \text{Cycle}(L)_i > 0 &\implies CI_i > 0 \\
&\therefore CI_i > 0. \quad \blacksquare [\text{Q.E.D.}]
\end{aligned}$$

This property is verified in the `CycleIntegralProperties::assertCycleIntegralPositive`.

References

- [1] T. H. Mata. Division and Modulo from Recursive Normalization. <http://ai.viXra.org/abs/2609.0009>, 2026. ai.viXra:2609.0009.
- [2] T. H. Mata. Formal Verification of Cyclic Lists. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter4/cycle.md>, 2026.
- [3] T. H. Mata. Formal Verification of Discrete Integration Properties from First Principles. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter4/integral.md>, 2026.
- [4] T. H. Mata. Using Formal Verification to Prove Properties of Lists Recursively Defined. <https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter3/list.md>, 2026.
- [5] The Lean Community. Mathlib: Cycles of Lists. https://leanprover-community.github.io/mathlib4_docs/Mathlib/Data/List/Cycle.html.
- [6] The Lean Community. Mathlib: Periodic Functions. https://leanprover-community.github.io/mathlib4_docs/Mathlib/Algebra/Ring/Periodic.html.