Relaxed Almost-Prime Production in Sieve Sequences

Thiago Henrique Ramos da Mata
Independent Researcher
thiago.henrique.mata@gmail.com
ORCID: 0009-0002-7366-939X
https://github.com/thiagomata
DOI: https://doi.org/10.5281/zenodo.22955792

Proof status: The Sieve Sequence foundation is Stainless-verified in the companion chapter article; the algebraic results introduced here are proved mathematically. The properties studied here are not yet Stainless-verified.

Abstract

The square-safe Sieve Sequence certifies that every survivor $p \in [Q, Q^2)$ is prime. Requiring $p+2$ to survive the same filters would ask for a twin-prime pair. We study a deliberately weaker target instead: require $p+2$ to avoid primes below $z = Q^{2\alpha}$ for some fixed $\alpha > 1/3$. Positivity then implies that $p+2$ has at most two prime factors for all sufficiently large Q.

We prove three exact algebraic results for this relaxed weight. The one-divisor count has an explicit local factor and one periodic boundary remainder. The natural pre-sieved divisor remainder is exactly a discrepancy of primes in the progression -2 modulo d. Finally, the scalar-centered final weight decomposes into nonprincipal character products $\chi(m)\chi(n)$. A quadratic character modulo 3 correlates with the full relaxed survivor count on the complete reduced wheel, refuting the shortcut that scalar-density centering creates arbitrary-coefficient Type-II orthogonality.

These results do not prove positivity of the relaxed weight. They identify its correct Type-I comparison, refute one over-strong Type-II formulation, and isolate the remaining need for an averaged prime-progression theorem followed by a locally adapted bilinear estimate.

Licensed under the Creative Commons Attribution 4.0 International License (CC BY 4.0):
https://creativecommons.org/licenses/by/4.0/.

1 Introduction

The twin-prime program requires both endpoints to survive every missing prime filter below a future head. This article investigates a distinct and weaker program: the first endpoint is square-safe prime, while the shifted endpoint is required only to avoid primes below a smaller threshold. The conclusion is a prime plus an integer with at most two prime factors, not a twin-prime pair.

The proof develops the following results in dependency order:

1. relaxed-weight positivity implies prime-plus-P_2 production — Section 3;
2. the exact divisor local factor and periodic remainder — Section 4;
3. the exact shifted-divisor prime-progression discrepancy — Section 5;
4. the exact bilinear character decomposition — Section 6; and

5. the refuted scalar-density Type-II shortcut — Section 7.

Sections 8–9 state the remaining analytic program and the exact claim boundary. No complete-period identity is substituted for an averaged short-interval distribution theorem.

1.1 Scope And Evidence Status

Let Q be a prime head and put

$$X = Q^2, \qquad W = P(Q) = \prod_{p<Q} p.$$

The installed-wheel survivors in the square-safe interval are

$$S_Q = \{n \in [Q, Q^2) : \gcd(n, W) = 1\}.$$

Every $n \in S_Q$ is prime: if it were composite and smaller than Q^2, it would have a prime divisor smaller than Q.

We do not prove a new existence theorem here. Classical Chen theory already supplies infinitely many primes p for which $p + 2$ has at most two prime factors. The project-specific question is whether positivity can be derived from the Sieve Sequence's own relaxed weights.

We prove the implication in Section 3 and the divisor local factor, cofactor progression discrepancy, and bilinear character obstruction properties in Sections 4–6 mathematically; none of the four is yet Stainless-verified. No statement in this article should be described as formally verified.

The maintained Sieve Sequence construction is Stainless-verified in the companion chapter article. This article uses that construction as an input but does not call its new number-theoretic properties verified. Each property section below states its population and scope before the proof, and links its supplementary working note at the end.

1.2 Relation To Known Results

This article does not reprove Chen's theorem [1], which already establishes infinitely many primes p such that $p + 2$ has at most two prime factors. The machinery invoked below is standard in the linear-sieve literature: the sifted sequence and level-of-distribution framework of Halberstam and Richert [3], with Type-I/Type-II terminology as in Iwaniec and Kowalski [4] and Friedlander and Iwaniec [2]. What is project-specific is the population: the square-safe interval $[Q, Q^2)$ anchored at the head, the nested wheels $2 \mid Z \mid W$, and the pre-sieved shifted sequence $\mathcal{A}_Q = \{n + 2 : n \in S_Q\}$. The modulo-3 obstruction proved in Section 7 is a specific fixed-character correlation on the complete reduced wheel; the classical parity barrier — a sieve's general inability to distinguish the parity of the number of prime factors — is the broader context for why such local obstructions defeat scalar-density models, but the two statements are not identical and are not conflated here.

2 Preliminaries And The Relaxed Candidate Weight

The sieve vocabulary used throughout:

- A P_2 **number** is a positive integer n with $\Omega(n) \le 2$, where Ω counts prime factors with multiplicity.

- A **lower-bound sieve** is a sieve method that produces a lower bound for the count of elements of a sequence surviving a set of divisibility conditions; the quantity $\sum_{Q\le n<Q^2} a_Q(n)$ of this article is the sifted sum such a method would bound from below.
- A **Type-I estimate** controls divisor sums: it bounds $\sum_d \tau(d)\,|A_d - A_1/\varphi(d)|$ for divisors d up to some level D, i.e. it uses distribution of the sequence on arithmetic progressions. For this article the relevant remainders are the prime-progression discrepancies of Section 5.
- A **Type-II estimate** controls bilinear sums $\sum_{m,n} \xi_m \kappa_n\, a_{mn}$ with arbitrary bounded coefficients, i.e. distribution of the sequence along products; Section 6 identifies the exact character modes such an estimate must control for this weight.

Choose a fixed exponent

$$\frac{1}{3} < \alpha < \frac{1}{2}$$

and define

$$z = X^\alpha = Q^{2\alpha}, \qquad Z = P(z).$$

The upper restriction $\alpha < 1/2$ is useful because it gives $z < Q$, hence $Z \mid W$. Define

$$a_Q(n) = \mathbf{1}_{\gcd(n,W)=1}\mathbf{1}_{\gcd(n+2,Z)=1}.$$

The candidate asks whether, for some fixed $\alpha > 1/3$ and infinitely many heads Q,

$$\sum_{Q\le n<Q^2} a_Q(n) > 0.$$

This is weaker than twin-prime positivity because $n+2$ is not required to survive every prime below Q.

3 Relaxed Positivity Implies Prime-Plus-Almost-Prime Production

Theorem 1 (Relaxed positivity implies prime-plus-P_2 production). For every fixed exponent $1/3 < \alpha < 1/2$ and every sufficiently large prime future head Q, over the integers in that head's square-safe interval weighted by a_Q:

$$a_Q(n) = 1 \quad \Longrightarrow \quad n \text{ is prime and } \Omega(n+2) \le 2.$$

The implication itself is proved; positivity for infinitely many heads remains open, and neither this implication nor the properties it depends on is yet Stainless-verified.

The relaxed weight keeps enough filtering to certify the first endpoint as prime and to bound the factorization depth of the second. It does not certify the second endpoint as prime.

Suppose $a_Q(n) = 1$ for some $Q \le n < Q^2$. By the installed-wheel factor, $\gcd(n, W) = 1$, so square-safe certification proves that n is prime. By the relaxed factor, $\gcd(n+2, P(z)) = 1$, so every prime factor of $n+2$ is at least $z = X^\alpha$.

Because $3\alpha > 1$, there is $X_0(\alpha)$ such that $X^{3\alpha} > X + 1$ whenever $X \ge X_0(\alpha)$. If $n+2$ had at least three prime factors counted with multiplicity, then

$$\begin{aligned}
\Omega(n+2) \geq 3 \Longrightarrow n+2 &\geq z^3 && \text{[Three Factor Lower Bound]} \\
&= X^{3\alpha} && \text{[By Definition Of } z\text{]} \\
&> X+1 && \text{[Since } 3\alpha > 1 \text{ And } X \geq X_0(\alpha)\text{]}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
n < Q^2 = X &\Longrightarrow n \leq X-1 && \text{[Integer Bound]} \\
&\Longrightarrow n+2 \leq X+1. && \text{[Add 2]}
\end{aligned}$$

The inequalities contradict each other. Therefore

$$a_Q(n) = 1 \Longrightarrow n \text{ is prime and } \Omega(n+2) \leq 2. \qquad \blacksquare \text{ [Q.E.D.]}$$

Consequently,

$$\sum_{Q \leq n < Q^2} a_Q(n) > 0 \Longrightarrow \exists n : \ n \text{ prime and } \Omega(n+2) \leq 2.$$

This is prime-plus-almost-prime production. It is neither a twin-prime certificate nor a proof that the relaxed sum is positive for any unbounded family of heads.

The project-specific candidate and this conditional proof are also recorded, with the same derivation, in Chen-Type Almost-Prime Survivor [5]. The first-endpoint input's proof is also recorded in Safe-Window 2-Gaps Certify Twin Primes. Neither that input nor the factor-count argument above is yet encoded as a `.holds` theorem.

4 Exact Divisor Local Factor And Boundary Remainder

Theorem 2 (Exact divisor local factor and boundary remainder). For every pair of squarefree prime wheels W, Z with $2 \mid W$, every integer interval $[L, U)$ with $L < U$, and every $m \geq 1$, over relaxed-weight integers in that interval that are also divisible by the fixed integer m:

$$\mathcal{N}_m[L,U) = \rho(m)\ell_m + E_m[L,U), \qquad |E_m[L,U)| \leq R-1.$$

Before studying divisor averages, the comparison density must account for how the tested divisor meets the two wheels. Let W and Z be squarefree with $2 \mid W$, and define

$$\mathcal{N}_m[L,U) = \#\{n \in [L,U) : m \mid n, \ \gcd(n,W) = 1, \ \gcd(n+2,Z) = 1\}.$$

Writing $n = mk$ converts the numerical interval to

$$K_m[L,U) = \left[\left\lceil \frac{L}{m} \right\rceil, \left\lceil \frac{U}{m} \right\rceil\right) \cap \mathbb{Z}, \qquad \ell_m = |K_m[L,U)|.$$

If $\gcd(m,W) > 1$, choose a prime $p \mid \gcd(m,W)$. Every $n = mk$ then has $p \mid n$ and cannot be coprime to W. Thus

$$\mathcal{N}_m[L,U) = 0.$$

Adopt the convention $\rho(m) = E_m[L,U) = 0$ whenever $\gcd(m,W) > 1$: with this convention the displayed identity $\mathcal{N}_m[L,U) = \rho(m)\ell_m + E_m[L,U)$ holds trivially in this branch, so the theorem's

quantifier over every $m \geq 1$ is exact rather than implicitly restricted to the coprime case developed below.

Assume now $\gcd(m, W) = 1$. For every prime $p \mid WZ$, let $\lambda_p(m)$ count the allowed residues of k modulo p. Direct local analysis gives

$$\lambda_p(m) = \begin{cases} p-1, & p \mid W,\ p \nmid Z, \\ 1, & p = 2,\ p \mid W,\ p \mid Z, \\ p-2, & p > 2,\ p \mid W,\ p \mid Z, \\ p, & p \mid Z,\ p \nmid W,\ p \mid m, \\ p-1, & p \mid Z,\ p \nmid W,\ p \nmid m. \end{cases}$$

Each row follows from an exact residue count:

- If $p \mid W$ and $p \nmid Z$, the hypothesis $\gcd(m, W) = 1$ makes m invertible modulo p. The installed condition forbids only $k \equiv 0 \pmod{p}$, leaving $p - 1$ classes.
- If $p \mid W$ and $p \mid Z$, the two conditions forbid $k \equiv 0 \pmod{p}$ and $k \equiv -2m^{-1} \pmod{p}$. They coincide for $p = 2$, leaving one class, and are distinct for $p > 2$, leaving $p - 2$ classes.
- If $p \mid Z$ and $p \nmid W$, then p is odd because $2 \mid W$. When $p \mid m$, one has $mk + 2 \equiv 2 \not\equiv 0 \pmod{p}$ for every k, so all p classes survive. When $p \nmid m$, invertibility of m leaves exactly one forbidden class and therefore $p - 1$ allowed classes.

These cases are exhaustive because every prime dividing WZ belongs to the installed wheel only, both wheels, or the relaxed wheel only. This proves the local table.

Put

$$R = \prod_{p \mid WZ} p, \qquad \rho(m) = \prod_{p \mid WZ} \frac{\lambda_p(m)}{p}.$$

CRT proves that every complete block of R consecutive k values contains exactly $R\rho(m)$ allowed values.

Let the k interval have length $\ell_m = qR + s$, with $0 \leq s < R$, and let C_m count allowed values in its final s positions. Then

$$\begin{aligned} \mathcal{N}_m[L, U) &= qR\rho(m) + C_m && \text{[Complete Periods Plus Remainder]} \\ &= \rho(m)\ell_m + (C_m - s\rho(m)). && \text{[Substitution]} \end{aligned}$$

Defining $E_m[L, U) = C_m - s\rho(m)$ gives

$$\mathcal{N}_m[L, U) = \rho(m)\ell_m + E_m[L, U), \qquad |E_m[L, U)| \leq s \leq R - 1. \qquad \blacksquare \text{ [Q.E.D.]}$$

In the candidate range $Z \mid W$, all divisors coprime to W have the same density

$$\rho_{Q,z} = \frac{1}{2} \prod_{2 < p < z} \left(1 - \frac{2}{p}\right) \prod_{z \leq p < Q} \left(1 - \frac{1}{p}\right).$$

This is sieve dimension two below z and dimension one from z to Q. The exact pointwise remainder bound is not a Type-I theorem because the primorial R is much larger than the square-safe interval.

No maintained Scala theorem currently models both squarefree wheels, all five local cases, CRT composition, and the arbitrary interval remainder; we leave it unverified rather than represent it with speculative code. Proving it would require establishing the local table one prime at a time and then using a verified CRT product lemma. The complete mathematical proof is also recorded, with the same derivation, in Relaxed Almost-Prime Weight Has An Exact Divisor Local Factor [8].

5 Shifted Divisor Discrepancy

Theorem 3 (Shifted divisor discrepancy reduces to prime progressions). For every squarefree installed wheel W with $2 \mid W$, every odd squarefree divisor $d \mid W$, and the square-safe interval I:

$$r_d(I) = A_d(I) - \frac{A_1(I)}{\varphi(d)} = \pi(I; d, -2) - \frac{\pi(I)}{\varphi(d)}.$$

This is proved over every finite integer interval $I = [L, U)$, for installed-wheel survivors in that interval whose shift $n + 2$ is constrained by d. The exact reduction is proved; the accumulated prime-progression estimate remains open.

The lower-bound sieve should be applied before the final relaxed filtering step. Its base sequence is

$$\mathcal{A}_Q = \{n + 2 : n \in S_Q\}.$$

For an odd squarefree divisor $d \mid W$, define

$$A_d(I) = \#\{n \in I : \gcd(n, W) = 1,\ d \mid n + 2\}, \qquad A_1(I) = \#\{n \in I : \gcd(n, W) = 1\}.$$

In one complete residue system modulo W, every prime dividing d fixes $n = -2$, while every other wheel prime permits all nonzero classes. Therefore

$$\begin{aligned} A_d([a, a + W)) &= \prod_{\substack{p \mid W \\ p \nmid d}} (p - 1) && \text{[CRT]} \\ &= \frac{\varphi(W)}{\varphi(d)} && \text{[Squarefree Products]} \\ &= \frac{A_1([a, a + W))}{\varphi(d)}. && \blacksquare \text{ [Q.E.D.]} \end{aligned}$$

Thus the centered word

$$h_d(n) = \mathbf{1}_{\gcd(n,W)=1} \left(\mathbf{1}_{d \mid n+2} - \frac{1}{\varphi(d)} \right)$$

is W-periodic with zero mean. For every interval I,

$$r_d(I) := A_d(I) - \frac{A_1(I)}{\varphi(d)}$$

is exactly the sum of h_d over the one incomplete wheel remainder after complete blocks are removed. More explicitly, write

$$|I| = qW + t, \qquad 0 \le t < W.$$

Partition I from its left endpoint into q complete consecutive blocks of length W and one final block of length t. Periodicity and the complete wheel identity give

$$\begin{aligned} r_d(I) &= \sum_{n \in I} h_d(n) && \text{[By Definition]} \\ &= \sum_{u=0}^{q-1} \sum_{j=0}^{W-1} h_d(L + uW + j) + \sum_{j=0}^{t-1} h_d(L + qW + j) && \text{[Block Decomposition]} \\ &= \sum_{j=0}^{t-1} h_d(L + qW + j). && \text{[Complete Blocks Have Zero Mean]} \end{aligned}$$

Since $|h_d(n)| \le 1$, the exact representation gives only the pointwise bound

$$|r_d(I)| \le t \le W - 1.$$

For a primorial W, this magnitude bound is too large to be a Type-I theorem; the useful fact is the exact signed remainder.

In the square-safe interval, wheel survivors are primes. Hence

$$r_d(I) = \pi(I; d, -2) - \frac{\pi(I)}{\varphi(d)}. \qquad \blacksquare \text{ [Q.E.D.]}$$

The missing Type-I input is consequently an averaged theorem of the form

$$\sum_{\substack{d \le D \\ d \mid P(z)/2}} \tau_B(d) \max_I \left| \pi(I; d, -2) - \frac{\pi(I)}{\varphi(d)} \right| \ll \frac{Q^2}{(\log Q)^A},$$

with a range D and interval family strong enough for the chosen lower-bound sieve. This displayed estimate is an open target, not a theorem of this article.

The complete-period CRT identity and periodic remainder are suitable for future formalization. The prime-progression interpretation also depends on square-safe certification. No maintained theorem currently connects all these pieces for arbitrary squarefree d; the accumulated analytic inequality lies outside what has been formalized either way. The complete mathematical reduction is also recorded, with the same derivation, in Relaxed Cofactor Divisor Sum Is A Prime-Progression Discrepancy [9].

6 Exact Bilinear Character Decomposition

Theorem 4 (Exact bilinear character decomposition). For every pair of squarefree nested wheels $2 \mid Z \mid W$, every factor pair with $\gcd(mn, W) = 1$, every finite factor domain $\mathcal{D}$, and arbitrary coefficients ξ_m, κ_n, the scalar-centered relaxed weight at products $x = mn$ decomposes exactly into nonprincipal character products $\chi(m)\chi(n)$ as displayed below.

Assume $2 \mid Z \mid W$ and put $Z_{\text{odd}} = Z/2$. Conditional on $\gcd(x, W) = 1$, the complete-wheel relaxed density is

$$\vartheta_Z = \prod_{p \mid Z_{\mathrm{odd}}} \left(1 - \frac{1}{p-1}\right).$$

Center the final weight by

$$w(x) = \mathbf{1}_{\gcd(x,W)=1} \left(\mathbf{1}_{\gcd(x+2,Z)=1} - \vartheta_Z\right).$$

If $\gcd(mn, W) = 1$, then m, n are odd units modulo every divisor of Z_{odd}, so $mn + 2$ is odd. The even terms in the Möbius coprimality identity vanish, while expansion of the squarefree Euler product for ϑ_Z gives

$$\mathbf{1}_{\gcd(mn+2,Z)=1} = \sum_{d \mid Z_{\mathrm{odd}}} \mu(d) \mathbf{1}_{mn \equiv -2 \pmod d},$$
$$\vartheta_Z = \sum_{d \mid Z_{\mathrm{odd}}} \frac{\mu(d)}{\varphi(d)}.$$

Subtracting yields the exact pointwise decomposition

$$w(mn) = \mathbf{1}_{\gcd(m,W)=\gcd(n,W)=1} \sum_{d \mid Z_{\mathrm{odd}}} \mu(d) \left(\mathbf{1}_{n \equiv -2m^{-1} \pmod d} - \frac{1}{\varphi(d)}\right).$$

Character orthogonality on the reduced residue group gives

$$\mathbf{1}_{mn \equiv -2 \pmod d} - \frac{1}{\varphi(d)} = \frac{1}{\varphi(d)} \sum_{\substack{\chi \pmod d \\ \chi \neq \chi_0}} \overline{\chi(-2)} \chi(m) \chi(n).$$

This is the genuine arbitrary-coefficient bilinear family. It is not removed by subtracting one scalar density. Indeed, for every finite factor domain $\mathcal{D}$ and arbitrary coefficients ξ_m, κ_n, substitution gives

$$\sum_{(m,n) \in \mathcal{D}} \xi_m \kappa_n w(mn) = \sum_{d \mid Z_{\mathrm{odd}}} \frac{\mu(d)}{\varphi(d)} \sum_{\substack{\chi \pmod d \\ \chi \neq \chi_0}} \overline{\chi(-2)} \cdot \sum_{\substack{(m,n) \in \mathcal{D} \\ \gcd(m,W)=\gcd(n,W)=1}} \xi_m \kappa_n \chi(m) \chi(n). \quad \blacksquare \text{ [Q.E.D.]}$$

The formula diagonalizes the local congruence modes. It does not estimate them: the geometry of $\mathcal{D}$, for example a hyperbolic restriction on mn, still couples the two variables.

The proof uses Möbius inversion and finite character orthogonality, neither of which currently has a Stainless representation for this weight in the project. The exact finite algebra is proved mathematically above and also recorded, with the same derivation, in Relaxed Almost-Prime Bilinear Remainder Has A Character Obstruction [7].

7 Refuted Route — Scalar-Density Type-II Orthogonality

Theorem 5 (Scalar-density Type-II orthogonality fails on the complete wheel). For every pair of squarefree wheels $2 \mid Z \mid W$ with $3 \mid Z$, the quadratic character modulo 3 gives bounded product coefficients with

$$\left|\sum_{m,n\in G_W}\chi_3(m)\chi_3(n)w(mn)\right| = \sum_{m,n\in G_W} a(mn).$$

This section refutes one auxiliary statement, using bounded character coefficients over product pairs on the complete reduced wheel $G_W \times G_W$. The refutation does not touch candidate #25 itself or short-domain locally adapted Type-II estimates.

The failed shortcut claimed that scalar centering makes the complete-wheel weight orthogonal to all bounded product coefficients, or at least gives a universal strict contraction $c < 1$ relative to the relaxed survivor count. The quadratic character modulo 3 disproves both claims.

Assume $3 \mid Z$. Let χ_3 be the nonprincipal real character modulo 3 and let

$$G_W = (\mathbb{Z}/W\mathbb{Z})^\times.$$

Choose bounded product coefficients

$$\xi_m = \chi_3(m), \qquad \kappa_n = \chi_3(n).$$

If the relaxed weight accepts mn, then $mn + 2$ is nonzero modulo 3. Because mn is a unit, necessarily $mn \equiv 2 \pmod 3$, and therefore

$$\xi_m\kappa_n = \chi_3(mn) = -1.$$

Hence the relaxed indicator has correlation

$$\sum_{m,n\in G_W}\xi_m\kappa_n a(mn) = -\sum_{m,n\in G_W} a(mn). \qquad \text{[Constant Character Sign]}$$

The scalar comparison is

$$b(x) = \vartheta_Z \mathbf{1}_{\gcd(x,W)=1}.$$

CRT balances the reduced residues between the two unit classes modulo 3, so $\sum_{m\in G_W}\chi_3(m) = 0$. Since every product of two wheel units is again a wheel unit,

$$\begin{aligned}\sum_{m,n\in G_W}\xi_m\kappa_n b(mn) &= \vartheta_Z\left(\sum_{m\in G_W}\chi_3(m)\right)\left(\sum_{n\in G_W}\chi_3(n)\right) && \text{[Scalar Comparison]}\\ &= 0. && \text{[CRT Balance]}\end{aligned}$$

Because $w = a - b$, subtraction gives

$$\left|\sum_{m,n\in G_W}\xi_m\kappa_n w(mn)\right| = \sum_{m,n\in G_W} a(mn). \qquad \blacksquare \text{ [Q.E.D.]}$$

Thus scalar-density centering does not produce complete-wheel orthogonality or any strict universal contraction against arbitrary product coefficients: the ratio to the survivor count is exactly 1.

This counterexample does not refute relaxed-weight positivity. It refutes only the proof shortcut that treats the final scalar-centered periodic weight as locally pseudorandom. A short hyperbolic factor domain is not a complete reduced wheel, so the counterexample also does not decide every locally adapted Type-II estimate.

The exact failed statement, counterexample, and retry boundary are archived in Scalar-Density Type-II Orthogonality For The Relaxed Weight [10]. The same character calculation is derived from the bilinear character decomposition proved in Section 6 above. No empirical sample is used in the refutation.

8 The Correct Remaining Program

The proved results impose an order on the remaining work.

First, match a theorem on primes in arithmetic progressions to the exact shifted-divisor remainder

$$\pi(I; d, -2) - \frac{\pi(I)}{\varphi(d)}.$$

The theorem must cover the divisor range and interval uniformity required by the chosen lower-bound sieve. A complete-period CRT count cannot replace that average.

Second, formulate Type II before the final relaxed sifting step, or use a comparison sequence that already includes every fixed local character mode. The scalar-centered final indicator is not an admissible pseudorandom model.

Third, verify that the proved Type-I and bilinear ranges actually make the lower-bound almost-prime weight positive. Neither local factors nor orthogonality formulas imply positivity on their own.

9 Claim Boundary

This article proves:

1. positivity of the relaxed weight implies a prime plus an integer with at most two prime factors;
2. the exact one-divisor local factor and periodic boundary remainder;
3. the exact shifted-divisor reduction to prime-progressions;
4. the exact inverse-residue and nonprincipal-character bilinear family; and
5. failure of scalar-density Type-II orthogonality on the complete wheel.

It does not prove:

- positivity for any unbounded family of heads;
- the accumulated prime-progression estimate;
- a locally adapted Type-II estimate;
- a new proof of Chen's theorem; or
- any twin-prime statement.

10 Conclusion

The relaxed program has a proved conditional conclusion. For fixed $1/3 < \alpha < 1/2$ and sufficiently large Q,

$$a_Q(n) = 1 \Longrightarrow n \text{ is prime and } \Omega(n+2) \le 2.$$

The divisor local factor property gives the exact one-divisor comparison. Wheel-sharing divisors vanish; coprime divisors have

$$\begin{aligned} \mathcal{N}_m[L,U) &= \rho(m)\ell_m + E_m[L,U), && \text{[Exact Interval Decomposition]} \\ |E_m[L,U)| &\le R-1, && \text{[Periodic Boundary]} \\ \rho_{Q,z} &= \frac{1}{2}\prod_{2<p<z}\left(1-\frac{2}{p}\right)\prod_{z\le p<Q}\left(1-\frac{1}{p}\right). && \text{[Nested-Wheel Specialization]} \end{aligned}$$

The cofactor progression discrepancy property identifies the natural pre-sieved Type-I remainder exactly:

$$\begin{aligned} r_d(I) &= A_d(I) - \frac{A_1(I)}{\varphi(d)} && \text{[Shifted Divisor Discrepancy]} \\ &= \pi(I;d,-2) - \frac{\pi(I)}{\varphi(d)}. && \text{[Square-Safe Prime Identity]} \end{aligned}$$

The bilinear character obstruction property gives the exact nonprincipal bilinear spectrum

$$\mathbf{1}_{mn\equiv -2 \pmod d} - \frac{1}{\varphi(d)} = \frac{1}{\varphi(d)}\sum_{\chi\neq\chi_0}\overline{\chi(-2)}\chi(m)\chi(n),$$

and the modulo-3 character refutes scalar-only centering:

$$\left|\sum_{m,n\in G_W}\chi_3(m)\chi_3(n)w(mn)\right| = \sum_{m,n\in G_W}a(mn).$$

The next theorem must therefore add genuine averaged prime-distribution information and use a locally adapted bilinear formulation. Even those two estimates must still be inserted into a lower-bound almost-prime identity that proves positivity. The article establishes the algebraic reductions and one failed shortcut; it does not supply that final analytic argument, a new proof of Chen's theorem, or a twin-prime result.

References

[1] J. R. Chen. On the representation of a larger even integer as the sum of a prime and the product of at most two primes. *Scientia Sinica*, 16:157–176, 1973.

[2] J. Friedlander and H. Iwaniec. *Opera de Cribro*, volume 57 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, 2010.

[3] H. Halberstam and H.-E. Richert. *Sieve Methods*. Academic Press, London, 1974.

[4] H. Iwaniec and E. Kowalski. *Analytic Number Theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, 2004.

[5] Thiago Henrique Ramos da Mata. Chen-Type Almost-Prime Survivor. https://github.com/thiagomata/prime-numbers/blob/master/candidates/chen-type-almost-prime-survivor.md, 2026.

[6] Thiago Henrique Ramos da Mata. Formal Verification of Sieve Sequence Stages and Their Transitions. https://github.com/thiagomata/prime-numbers/blob/master/articles/chapter6/sieve-sequence.md, 2026.

[7] Thiago Henrique Ramos da Mata. Relaxed Almost-Prime Bilinear Remainder Has A Character Obstruction. https://github.com/thiagomata/prime-numbers/blob/master/properties/sieve-sequence/relaxed-almost-prime-bilinear-character-obstruction.md, 2026.

[8] Thiago Henrique Ramos da Mata. Relaxed Almost-Prime Weight Has An Exact Divisor Local Factor. https://github.com/thiagomata/prime-numbers/blob/master/properties/sieve-sequence/relaxed-almost-prime-divisor-local-factor.md, 2026.

[9] Thiago Henrique Ramos da Mata. Relaxed Cofactor Divisor Sum Is A Prime-Progression Discrepancy. https://github.com/thiagomata/prime-numbers/blob/master/properties/sieve-sequence/relaxed-cofactor-divisor-sum-is-prime-progression-discrepancy.md, 2026.

[10] Thiago Henrique Ramos da Mata. Scalar-Density Type-II Orthogonality For The Relaxed Weight. https://github.com/thiagomata/prime-numbers/blob/master/candidates/refuted/relaxed-weight-scalar-density-type-ii.md, 2026.

A Evidence And Verification Status

Result	Mathematical status	Stainless status	Cross-reference
Relaxed positivity implies prime-plus-P_2	Conditional implication proved; positivity for infinitely many heads open	Not yet verified	Candidate #25
Exact divisor local factor	Proved, including all five local cases and arbitrary-interval remainder	Not yet verified	Divisor Local Factor
Shifted divisor discrepancy	Exact reduction proved; accumulated prime-progression estimate open	Not yet verified	Cofactor Progression Discrepancy
Bilinear character decomposition	Exact pointwise and arbitrary-domain decompositions proved	Not yet verified	Bilinear Character Obstruction
Scalar-density Type-II shortcut	[Refuted] on the complete reduced wheel; short locally adapted domains remain open	Not applicable to a false statement	Archived refutation

The operational Sieve Sequence construction and its square-safe inputs are documented separately in Formal Verification of Sieve Sequence Stages and Their Transitions [6]. The new arithmetic results in this article remain mathematically proved; none has a Stainless representation.