

The ACT measurement of the lensing power spectrum

May 9, 2023

In April 2023, the Atacama Cosmology Telescope collaboration has released a measurement of a **signal dominated lensing map** over a quarter of the sky, and its **associated power spectrum**. The goal of this tutorial is to compute the underlying theoretical lensing power spectrum in the Λ CDM model and compare it to the data.




Figure 1: Top: ACT DR6 CMB lensing mass map presented in this work. The map covers 9400 deg^2 or sky fraction $f_{\text{sky}} = 0.23$ with a signal-to-noise significantly greater than unity over a wide range of scales. Bright orange corresponding to peaks of the dark-matter dominated mass distribution and dark purple regions corresponding to voids. Bottom: the ACT DR6 lensing potential power spectrum bandpowers for our baseline (combined temperature and polarization) analysis

Useful quantities

$$\begin{aligned}
1 \text{ Mpc} &= 3.086 \times 10^{22} \text{ m} \\
G &= 6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2} \\
m_H &= 1.6735575 \times 10^{-27} \text{ kg} \\
c &= 3 \times 10^8 \text{ m s}^{-1}
\end{aligned} \tag{1}$$

1 Energy densities

The first step for predicting the form and amplitude of the Λ CDM lensing power spectrum is to compute the densities of the different components in the Universe.

We will work in the LCDM framework with the following cosmological parameters:

$$\begin{aligned}
T_{\text{CMB}} &= 2.7255 \text{ K} \\
H_0 &= 67.5 \text{ km/s/Mpc} \\
\Omega_b h^2 &= 0.022 \\
\Omega_{\text{cdm}} h^2 &= 0.122 \\
\Omega_m &= \Omega_b + \Omega_{\text{cdm}}
\end{aligned} \tag{2}$$

Ω_b stands for Omega baryons, and is defined as the ratio of the current baryon density over the current critical density of the Universe

$$\Omega_b = \frac{\rho_b^0}{\rho_c^0}$$

the critical density today is defined such as

$$\rho_c^0 = \frac{3H_0^2}{8\pi G} \tag{3}$$

and h is the reduced Hubble factor

$$h = \frac{H_0}{100} \tag{4}$$

1. Compute the numerical value of H_0 in s^{-1} unit.
2. Compute the numerical values of ρ_b^0 and ρ_{cdm}^0 in kg/m^3 , assume that all baryonic matter is composed of hydrogen, how many hydrogen atoms is there per cubic meter in the universe ?
3. Photons at equilibrium with temperature T_{CMB} follow a Bose-Einstein distribution $f_\gamma(\nu|T_{\text{CMB}}) = \left[\exp\left(\frac{h\nu}{k_B T_{\text{CMB}}} - 1\right) \right]^{-1}$, the energy density of photons is given by

$$\rho_\gamma^0 c^2 = 2 \int \frac{d^3 p}{(2\pi\hbar)^3} h\nu f_\gamma(\nu|T_{\text{CMB}}) \tag{5}$$

for photons, $p = \frac{h\nu}{c}$, setting $x = \frac{h\nu}{k_B T_{\text{CMB}}}$ show that the integral can be written

$$\rho_\gamma^0 c^2 = \frac{(k_B T_{\text{CMB}})^4}{\hbar^3 c^3 \pi^2} \int_0^\infty dx x^3 [\exp(x - 1)]^{-1} \tag{6}$$

Using $\Gamma(s)\zeta(s) = \int_0^\infty \frac{t^{s-1}}{e^t - 1} dt$ and $\Gamma(4) = 3!$, $\zeta(4) = \frac{\pi^4}{90}$ Show that

$$\rho_\gamma^0 = \frac{\pi^2 (k_B T_{\text{CMB}})^4}{15 \hbar^3 c^5} \tag{7}$$




Figure 2: comoving distance as a function of redshift in the LCDM Universe.

4. Compute the numerical values of Ω_γ .
5. We also need to include the contribution from neutrinos, the formula for the neutrinos density is similar to the one of photons with some corrections, a factor $\frac{7}{8}$ coming from the fact that neutrinos are fermions and a factor $(4/11)^{4/3}$ coming from the fact that neutrinos are a bit colder than photons because they decouple before the electron-positron annihilation (which reheat the plasma) $T_\nu = (4/11)^{1/3}T_\gamma$, we also need to take into account the fact that there are three species of neutrinos $N_{\text{eff}} = 3.046$ (departure from exactly 3 comes from QED correction)

$$\rho_\nu^0 = N_{\text{eff}} \frac{7}{8} \frac{\pi^2 (k_B T_\nu)^4}{15 \hbar^3 c^5} = N_{\text{eff}} \frac{7}{8} (4/11)^{4/3} \rho_\gamma^0 \quad (8)$$

Compute the numerical value of $\Omega_{\text{rad}} = \Omega_\gamma + \Omega_\nu$.

6. Get the value of Ω_Λ assuming that $\Omega_m + \Omega_{\text{rad}} + \Omega_\Lambda = 1$, note that this relation is only true if the universe is flat (which is one of our assumptions here).

2 Distances

The next step is to evaluate distances in the FLRW Universe

There are a lot of useful distance definitions in an expanding Universe, here we will use comoving distance. Imagine a pair of galaxies, their physical distance will increase due to the expansion of the Universe, but their comoving distance, which is measured in coordinates following the expansion of the Universe will stay fixed. The comoving distance between us and an object at redshift z can be computed as (see paragraph 1 of [notes](#))

$$\chi = \int_0^z \frac{cdz'}{H(z')} \quad (9)$$

Where

$$H(z) = H_0 \sqrt{\Omega_{\text{rad}}(1+z)^4 + \Omega_m(1+z)^3 + \Omega_\Lambda} \quad (10)$$

1. Compute the numerical value of the co-moving distance between us and the last scattering surface, assuming the CMB is emitted at $z_* = 1100$ and give its value in Mpc. What is its value in Giga light-year?, compare it to the age of the Universe which is 13.787 Giga year.

2. Compute the comoving distance for an array of 10^4 redshifts, logarithmically spaced¹ between $z_{\min} = 10^{-4}$ and $z_{\max} = 10^4$, you should get something looking like Fig 2.
3. Something that will be useful for later is a way to get the redshift associated to a given comoving distance, so basically inverting Eq 9. A simple way to create this is to interpolate the array created in the previous question to get $z(\chi)$. you can use the python function:

```

1 import numpy as np
2 from scipy.interpolate import InterpolatedUnivariateSpline
3
4 def interpolate_z_and_chi(H0, Omega_r, Omega_m, Omega_Lambda, logmin_z
   =-4, logmax_z=4, nz=10**4):
5
6     """
7     This function return the interpolation of z(chi)
8
9     Parameters
10    -----
11    H0: float
12         Hubble constant value in km/s/Mpc
13    Omega_r: float
14         Omega radiation
15    Omega_m: float
16         Omega matter
17    Omega_lambda: float
18         Omega dark energy
19    logmin_z: float
20         Minimal value of log(z) for the interpolation
21    logmax_z: float
22         Maximum value of log(z) for the interpolation
23    nz: integer
24         number of z values
25    """
26
27    redshift = np.logspace(logmin_z, logmax_z, nz)
28    chi = np.zeros(nz)
29    for i, z in enumerate(redshift):
30        chi[i] = comoving_distance(z, H0, Omega_r, Omega_m, Omega_Lambda)
31
32    z_of_chi = InterpolatedUnivariateSpline(chi, redshift)
33
34    return z_of_chi

```

What is the redshift corresponding to a comoving distance of 1 Gpc ?

3 Lensing

The lensing can be described from a remapping of the CMB photons from a position

$$\mathbf{x}_{\text{true}} = \chi_{\text{LSS}} \begin{pmatrix} \theta_S^1 \\ \theta_S^2 \\ 1 \end{pmatrix} \quad (11)$$

¹You can form the redshift array using `np.logspace(-4,4,10000)`




Figure 3: Path of a light ray without (left) and with (right) lensing.




Figure 4: drawing for question (a)

to a position

$$\mathbf{x}_{\text{obs}} = \chi_{\text{LSS}} \begin{pmatrix} \theta^1 \\ \theta^2 \\ 1 \end{pmatrix} \quad (12)$$

Where θ_S^1, θ_S^2 refers to the true source position, while θ^1, θ^2 refers to its observed position (see Fig 3). χ_{LSS} is the comoving distance to the last scattering surface. After a lengthy derivation (see paragraph 2 of [notes](#)), The relationship between the two positions is given by

$$\theta_S^i = \theta^i + \Delta\theta^i \quad (13)$$

$$(14)$$

$$\Delta\theta^i = \delta^{ij} \frac{\partial}{\partial\theta^j} \phi^L(\boldsymbol{\theta}) \quad (15)$$

$$\phi^L(\boldsymbol{\theta}) = -2 \int_0^{\chi_{\text{LSS}}} \frac{d\chi_1}{\chi_1} \psi(\mathbf{x}(\boldsymbol{\theta}, \chi_1), \eta_0 - \chi_1) \left(1 - \frac{\chi_1}{\chi_{\text{LSS}}}\right) \quad (16)$$

$\phi^L(\boldsymbol{\theta})$ is the lensing potential, its angular gradient will encode the change of position between true and observed position of the source. We see that it is given as an integral along the line of sight of the gravitational potential ψ times a geometric factor $\left(\frac{\chi_{\text{LSS}} - \chi_1}{\chi_1 \chi_{\text{LSS}}}\right)$ accounting for where the deflection happens.

- (a) (Optional) Let's check that the deflection has the correct sign, for simplicity let's think about it with a single lens and only two dimensions. The situation is represented in Figure 4. Let's imagine that the center of the lens sit at x_L , what will be the sign of $\Delta\theta = \theta_S - \theta$ for a source located at $x_S = x_L + \delta x$ with δx positive) ?

Tip: you can use that for $x > x_L$ on the lens plane, $\frac{\partial}{\partial\theta} \psi > 0$ since ψ is negative for over-density, and $\psi_{\text{min}} = \psi(x_L)$

The lensing power is given by (see paragraph 3 of [notes](#))

$$C_\ell^L = 4 \int_0^{\chi_{\text{LSS}}} d\chi_1 \left(\frac{\chi_{\text{LSS}} - \chi_1}{\chi_1^2 \chi_{\text{LSS}}} \right)^2 P_\psi \left(k = \frac{\ell + 1/2}{\chi_1}, z(\chi_1) \right) \quad (17)$$

This is the quantity we want to evaluate in this tutorial, we already have a way to compute χ so what we need is an expression for $P_\psi \left(k = \frac{\ell + 1/2}{\chi_1}, z(\chi_1) \right)$. In order to get this expression, we need to solve the Einstein-Boltzmann equations for perturbations in the Universe, I have done it for you using the public code CAMB

1. First download the tar file at the following [url](#), and untar it, the data consist of a power spectrum of the form $k^2 P_\psi(k, z)$ for different values of k and z . We will make an interpolation table using the following function

```

1 import numpy as np
2 from scipy.interpolate import interp2d
3
4 def get_P_z_and_k_interp(mode):
5     """
6     This function read and interpolate k^2 P_{\psi}(k, z)
7
8     Parameters
9     -----
10    mode: str

```

```

11         either "lcdm" or "baryons_only"
12
13     """
14
15     P_z_and_k = np.load(f"data/P_k_and_z_{mode}.npz")
16     k_array = np.load("data/k_array.npy")
17     z_array = np.load("data/z_array.npy")
18     P_z_and_k_interp = interp2d(z_array, k_array, P_z_and_k.flatten())
19     return z_array, k_array, P_z_and_k_interp

```

I have precomputed $k^2 P_\psi(k, z)$ for two modes, 'lcdm' and 'baryons_only', we will first do the computation with the lcdm mode.

2. Plot $k^2 P_\psi$ as a function of k at different redshifts $z = [0, 0.5, 1, 4, 20]$
3. The integral for computing C_ℓ^L goes from 0 to the comoving distance to the last scattering surface χ_{LSS} , you can see that it diverges at 0, in reality this doesn't affect the result because it only affect the $\ell = 0$ term. However it can lead to numerical instability so instead of doing the integral $\int_0^{\chi_{\text{LSS}}} d\chi_1$ we will do $\int_{\chi_{\text{min}}}^{\chi_{\text{LSS}}} d\chi_1$ with $\chi_{\text{min}} = 10$.
Create an evenly spaced array χ of comoving distance between χ_{min} and χ_{LSS} containing 100 values.
4. Create an array of the corresponding redshift values $z(\chi)$ using the interpolation function that we have defined in Section 2.
5. Ok time to do the integral, we will compute the lensing power spectrum for $\ell \in [2, 2000]$, looking at the formula we see that the lensing potential will be computed as

```

1 l_array = np.arange(2, 2000 + 1, dtype=np.float64)
2 cl_phi = np.zeros(l_array.shape)
3 for i, l in enumerate(l_array):
4     k_array = (1 + 0.5) / chi_array
5     cl_phi[i] = 0
6     for z, chi, k in zip(z_array, chi_array, k_array):
7         cl_phi[i] += 4 * dchis * P_z_and_k_interp(z, k) * ((chi_star -
            chi) / (chi ** 2 * chi_star)) ** 2 / k ** 4

```

Plot the resulting C_ℓ^L multiplied by $\frac{\ell^{\frac{5}{2}}(\ell+1)^2}{2\pi}$ and compare it with Figure 1.

6. In the same tarball as for question 1, you will find the measured lensing power spectrum "result_act.dat" with format, $(\ell_b, \frac{\ell_b^{\frac{5}{2}}(\ell_b+1)^2}{2\pi} C_{\ell_b}^L, \frac{\ell_b^{\frac{5}{2}}(\ell_b+1)^2}{2\pi} \sigma_{\ell_b}^L)$, where $\sigma_{\ell_b}^L$ are the errors on the measurement and ℓ_b the central value of the ℓ bin. Plot these data on the top of the theory curve. Note that there are small deviations compared to Figure 1, this is due to some approximations we made along the way in this derivation.
7. Redo the computation for the mode 'baryons_only', and conclude.