State-of-the-Art Workup and Initial **Management of Newly Diagnosed Molar Pregnancy and Postmolar** Gestational Trophoblastic Neoplasia

Kevin M. Elias, MD; Ross S. Berkowitz, MD; and Neil S. Horowitz, MD

ABSTRACT

Gestational trophoblastic disease refers to a series of interrelated tumors arising from the placenta, including benign molar pregnancies as well as the malignant conditions termed gestational trophoblastic neoplasia (GTN). GTN most commonly follows a molar pregnancy but may develop after any gestation. The wide availability of first trimester ultrasound and serum human chorionic gonadotropin (hCG) measurement has changed the presentation of molar pregnancy in recent decades from a second trimester to a first trimester disease, such that most patients have few symptoms at diagnosis. With identification of molar pregnancy at earlier gestations, accurate diagnosis increasingly relies on expert histopathology coupled with ancillary molecular and genetic techniques. However, earlier diagnosis has not changed the risk of postmolar GTN. Although most molar pregnancies are treated with dilation and curettage, hysterectomy may be appropriate in select cases when future fertility is not desired. After treatment of molar pregnancy, close surveillance with serial hCG monitoring is essential to diagnose GTN and identify the need for chemotherapy. Physicians following hCG levels should understand the performance characteristics of the test, including common causes of false-positive and false-negative results. After a diagnosis of postmolar GTN is made, selection of single-agent or multiagent chemotherapy depends on accurate assignment of the clinical stage and risk stratification by the International Federation of Gynecology and Obstetrics (FIGO) prognostic scoring system. Surgical treatment of postmolar low-risk GTN, including both second uterine curettage and hysterectomy, may decrease subsequent need for or duration of chemotherapy. Cure rates for postmolar low-risk GTN approach 100%, and subsequent pregnancy outcomes for patients reflect those of the general population.

J Natl Compr Canc Netw 2019;17(11):1396-1401 doi: 10.6004/jnccn.2019.7364

New England Trophoblastic Disease Center, Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Brigham and Women's Hospital, Dana-Farber Cancer Institute, Harvard Medical School, Boston, Massachusetts

Gestational trophoblastic disease includes a spectrum of interrelated tumors, including complete hydatidiform mole (CHM) and partial hydatidiform mole (PHM), invasive mole, choriocarcinoma, placental site trophoblastic tumor (PSTT), and epithelioid trophoblastic tumor (ETT). 1-3 After CHM and PHM, gestational trophoblastic neoplasia (GTN) develops in 15% to 20% and 1% to 4% of patients, respectively, with local uterine invasion with or without metastases.3,4 GTN most commonly follows a molar pregnancy but may develop after any gestation.

Molar Pregnancy: Complete Versus Partial

CHM and PHM differ based on their chromosomal pattern, gross morphology, and histopathology (Table 1).5 CHM is generally diploid, and the molar chromosomes are generally entirely of paternal origin.5 CHM is associated with more generalized trophoblastic hyperplasia and hydropic swelling of the chorionic villi and is not associated with fetal or embryonic tissues. PHM is generally triploid, in which the extra set of chromosomes is of paternal origin. PHM is associated with more focal trophoblastic hyperplasia and hydropic swelling of villi and is often associated with fetal or embryonic tissues. Rarely, in familial recurrent molar pregnancy CHM has a biparental chromosomal pattern and is associated with a mutation in the genes NLRP7 or KHDC3L.6,7

Presentation of Molar Pregnancy

The classic description of CHM includes excessive uterine enlargement, theca lutein ovarian cysts, hyperemesis, preeclampsia, and hyperthyroidism. The medical complications of theca lutein ovarian cysts, preeclampsia, and hyperthyroidism are primarily associated with marked trophoblastic proliferation with particularly high human chorionic gonadotropin (hCG) levels and excessive uterine size. In recent years, CHM is more commonly diagnosed earlier in the first trimester, and these dramatic signs and symptoms are much less frequent.8-11 Although excessive uterine size, preeclampsia, hyperemesis, and hyperthyroidism used to occur in 51%, 27%, 26%, and p57 expression

Table 1. Genetic and Histopathologic Features of Molar Pregnancy		
	Complete	Partial
Karyotype	Generally diploid or tetraploid; generally all chromosomes paternal	Generally triploid; extra set of chromosomes is paternal
Hydropic villi	Diffuse	Focal
Trophoblastic hyperplasia	Diffuse	Focal
Scalloping of villi	Absent	Present
Fetal or embryonic tissue	Absent	Present

7% of cases, respectively, during the 1970s and 1980s, in recent years these symptoms have declined to 28%, 1.3%, 8%, and 0%, respectively.⁸

Positive

Negative

The availability of both ultrasonography and hCG measurement has led to the earlier diagnosis of CHM. If a patient presents with irregular bleeding during pregnancy, ultrasound and hCG measurement are commonly obtained and may diagnose an abnormal pregnancy, leading to pregnancy termination. Although approximately 50% of complete moles are associated with pre-evacuation hCG levels >100,000 mIU/mL, a single hCG level is seldom helpful in differentiating a complete mole from another type of pregnancy. Early diagnosis of CHM has also led to a change in pathology, with less prominent trophoblastic hyperplasia and hydropic villi. ¹² Early CHM can be more difficult to distinguish pathologically from PHM and hydropic abortion. The absence of p57 expression on the maternal allele can be helpful in diagnosing early CHM. Importantly, early diagnosis of CHM has not led to a reduction in the development of GTN, although it has resulted in fewer medical and surgical complications.8-11

In previous years, PHM rarely presented with the classic signs and symptoms of CHM. Patients with PHM generally presented with irregular bleeding and were usually thought to have a missed or incomplete abortion. ^{13,14} In a series of 81 patients with PHM, 74 (91%) were thought to have a missed or incomplete abortion before uterine evacuation. ¹³ In more recent years, however, similar to CHM, PHM is generally diagnosed in the first trimester. Sun et al ¹⁵ reported that, at Brigham and Women's Hospital from 1994 through 2013, the median gestational age at uterine evacuation for CHM and PHM was 9 and 12 weeks, respectively.

Ultrasound and Molar Pregnancy

Ultrasound is the most precise method of diagnosing CHM and PHM before evacuation. Classic sonographic findings of CHM include a heterogeneous uterine mass with cystic spaces, no identifiable fetus or embryo, and no amniotic fluid. Ovarian theca lutein cysts may also be present. However, the classic ultrasound features may be subtle or absent in an early first trimester CHM. Estimates of the sensitivity of pelvic ultrasound in diagnosing complete mole range from 70% to 90%, and increase with gestational age. A study of 1,053 patients reported that an ultrasound finding of a molar pregnancy had a specificity of 74% for the diagnosis of CHM or PHM.

PHM is more frequently misdiagnosed as an incomplete or missed abortion compared with CHM, because PHM is associated with the presence of fetal or embryonic tissue and amniotic fluid. PHM is diagnosed as an incomplete or missed abortion in 15% to 60% of cases. ^{16–18} Ultrasound findings suggesting PHM include the presence of cystic spaces in the placenta, an increase in the transverse diameter of the gestational sac, and the presence of fetal and embryonic tissue and amniotic fluid. ¹⁹

Measuring hCG level at ultrasound may help distinguish CHM or PHM from a missed abortion. An hCG level >100,000 mIU/mL in the first trimester should strongly raise the suspicion of a molar pregnancy.²⁰

Treatment of Molar Pregnancy

Once molar pregnancy is suspected, the patient should be evaluated for the presence of possible medical complications, such as preeclampsia, electrolyte imbalance from hyperemesis, anemia, and hyperthyroidism. After the patient is stabilized, a decision must be made concerning the most appropriate method of evacuation.

Patients wishing to preserve fertility should undergo suction curettage through either manual or electric vacuum aspiration regardless of uterine size.²¹ For uteri larger than 14 weeks' size, evacuation under ultrasound guidance is recommended to ensure complete emptying of the uterine cavity. Medical induction of labor or hysterotomy are strongly discouraged because both techniques increase maternal morbidity, especially risk of hemorrhage. If the patient has a very high hCG level (>500,000 mIU/mL) or is suggestive of being hyperthyroid, anesthesia or surgery may precipitate thyroid storm. For these patients, a beta-adrenergic blocker should be administered at anesthesia. Patients with high hCG levels and excessive uterine enlargement may also be at increased risk for respiratory distress during and after evacuation. This risk is due to multiple factors, including heavy bleeding with high fluid replacement, trophoblastic pulmonary embolization, preeclampsia, and thyroid storm with high-output heart failure. The anesthesia and recovery room staff should be alerted to these possible complications. If the patient is Rh-negative, Rho (D) immunoglobulin should be administered after the procedure.

Hysterectomy may be considered in patients who do not desire future fertility, particularly among women of advanced maternal age. The risks of developing GTN after suction evacuation of CHM in patients aged >40 and >50 years are 54% and 60%, respectively; hysterectomy appears to reduce this risk and the subsequent need for chemotherapy. Hysterectomy may therefore be particularly reasonable to consider in patients aged >40 years with CHM.^{22,23} If the ovaries have prominent theca lutein cysts, these can be aspirated and preserved.

Prophylactic Chemotherapy

Patients with CHM with very high hCG values and other signs of marked trophoblastic growth, such as excessive uterine size (>20 weeks) or theca lutein cysts, are at high risk (40%–50%) of developing GTN. Patients with CHM who present with medical complications (eg, preeclampsia, marked hyperthyroidism, respiratory insufficiency) and those with repeat molar pregnancy are also at increased risk for GTN. Prophylactic chemotherapy at evacuation of high-risk CHM has reduced the incidence of GTN from approximately 40% to 10% to 15% in randomized controlled trials.^{24–26} We consider using chemoprophylaxis in patients with high-risk CHM when there is a significant concern about the reliability or availability of hCG follow-up.

Role of hCG Surveillance

After uterine evacuation, patients should be enrolled in a serum hCG surveillance protocol. Serum hCG levels are a highly sensitive marker for trophoblastic proliferation. While on surveillance, patients should use a reliable form of contraception to avoid confusing a new pregnancy with GTN.

Patients under hCG surveillance should have serum hCG levels checked weekly until the value is below the reference range for the assay. Traditionally, this level has been 5 mIU/mL. However, recent higher-sensitivity assays may produce values even <1 mIU/mL. Because most studies on hCG surveillance were performed with a cutoff value of 5 mIU/mL, we have continued to use this as the clinically actionable result.

Similarly, it is important that patients obtain serial hCG measurements at the same reference laboratory. This laboratory should use an assay that can detect all of the various isoforms of hCG (eg, total hCG, nicked hCG, free beta subunit).²⁷ Detection of hCG beta and hCG beta core fragment are particularly variable among assays.²⁸ Reference ranges, calibration, and assay sensitivity may differ markedly among laboratories, which can lead to either false-negative results or overdiagnosis of persistent disease.²⁹ This is especially true when results are compared among different laboratories. When patients present to our center, we always confirm an hCG result

on our local assay. However, for patients who will be monitored in the community setting, we advise continued use of the same reference laboratory.²⁸

Markedly elevated hCG values can also produce falsely low hCG results through a phenomenon known as the "hook effect," which generally occurs when the hCG level exceeds 1,000,000 mIU/mL. This occurs because most hCG tests are based on sandwich enzyme-linked immunosorbent assay or radioimmuno-assay techniques. Excess analyte saturates both the fixed, solid-phase antibodies and the labeled, soluble antibodies, preventing sandwich formation. If a hook effect is suspected, a true hCG value can be obtained by performing a serial dilution of the sample before analysis. ³¹

In cases of persistent low-level hCG, false-positive results or "phantom hCG" must also be considered.32 One common cause of phantom hCG is heterophile antibodies.33 In this case, the patient may have antimouse antibodies that interfere with the immunoassay. Heterophile antibodies can be excluded by obtaining a urine hCG test result, because heterophile antibodies do not cross the glomerulus. Heterophile antibodies can also be controlled for by using heterophile-blocking tubes. Another common cause of persistent low-level elevations in hCG is pituitary hCG, especially in perimenopausal women,^{29,34} in whom the prevalence of detectable hCG levels up to 14 mIU/mL is 8%.35 Pituitary hCG can be distinguished from real hCG by placing a patient on oral contraceptive pills, which will suppress pituitary hCG production.

Duration of hCG Surveillance

Patients should be followed up with weekly serum hCG measurements until the result is below the reference range for 3 weeks, and then with monthly testing thereafter. The American College of Obstetricians and Gynecologists recommends continuing postmolar hCG surveillance until 6 months of "normal" values are obtained. However, the optimal duration of hCG surveillance has been debated. In studies performed at Brigham and Women's Hospital, we found that the risk of GTN for both CHM and PHM is <1% after a single normal hCG value. The similar findings were identified in Brazil, where the risk of GTN after hCG normalization was 0.6% after CHM and 0.1% after PHM.

In the largest series, >20,000 patients with molar pregnancy from the Charing Cross Hospital in London underwent testing. After a single normal value, the risk of GTN after CHM was approximately 0.25%, and after PHM was approximately 0.03%. Based on these data, our practice has been to shorten the duration of hCG surveillance to 3 months for CHM and 1 month for PHM. However, this recommendation is only for patients experiencing spontaneous hCG remission. For patients

diagnosed with and treated for GTN, we continue hCG surveillance for 1 year.

Diagnosis of Postmolar GTN

The 2002 International Federation of Gynecology and Obstetrics (FIGO) guidelines present hCG monitoring criteria for the diagnosis of persistent neoplasia after evacuation of a molar pregnancy.41 Progressively increasing hCG levels across 3 values at least 14 days apart (ie, days 1, 7, and 14) or plateaued hCG values within a 10% range across 4 values at least 21 days apart (ie, days 1, 7, 14, and 21) establish a diagnosis of GTN. The 2002 FIGO guidelines also recommend treatment of GTN if hCG levels remain elevated >6 months after uterine evacuation. However, this last recommendation has been challenged by evidence from British and Brazilian referral centers for gestational trophoblastic disease suggesting that among women with persistently elevated but decreasing hCG levels at 6 months postevacuation, >80% of patients may experience spontaneous remission without chemotherapy. 42,43 Notably, extended surveillance does not appear associated with an increased risk of metastases, chemotherapy resistance, or need for multiagent chemotherapy. 42,43 Based on recent studies, FIGO no longer recommends starting chemotherapy when hCG is still elevated but declining 6 months after molar evacuation.3,42

Staging of GTN

Once the criteria for postmolar GTN are met, FIGO stage and score should be determined.44 Staging consists of 2 parts: a FIGO anatomic staging and a FIGO risk score to determine the risk of disease progression and resistance to single-agent chemotherapy. The score is calculated by adding the total points assigned to various clinical and tumor risk factors. Women with a FIGO score ≤6 are considered to have low-risk disease and can be treated with single-agent chemotherapy. 41,45,46 It is critically important to use a plain chest radiograph rather than a CT scan when staging patients, because approximately 30% to 40% of CT scans can show micrometastases that do not influence outcome and could lead to more aggressive chemotherapy than is needed.⁴⁷ Additionally, the hCG value used in the scoring system is the hCG level at persistence, not at the value at initial molar diagnosis or before evacuation of the mole. Again, failure to use the correct hCG value can result in an inappropriately elevated FIGO risk score. 45

Treatment of Low-Risk Postmolar GTN

Most patients who are followed up with serial hCG levels after evacuation of a molar pregnancy and are determined to have postmolar GTN based on FIGO criteria will have low-risk GTN (FIGO score <7). Treatment of low-risk GTN consists of single-agent chemotherapy,

typically with either methotrexate or actinomycin D. Multiple dosing and infusion schedules for both agents have been investigated, but given the heterogeneity of the data, no clearly preferred regimen has been determined. At our institution, we use the 8-day methotrexate regimen due to its low toxicity, ease of administration, and large reported clinical experience. Efficacy for both methotrexate and actinomycin D is similar, with complete remission rates of approximately 70% to 80%. Metaanalysis and randomized trials suggest that actinomycin D may have a higher rate of cure and require fewer cycles of therapy to achieve remission; however, much of these data come from studies comparing actinomycin D versus weekly intramuscular or infusional methotrexate rather than the more effective 5-day or 8-day regimens. 48-54 Weekly intramuscular methotrexate is no longer recommended for low-risk GTN because of lower reported efficacy rates compared with the other methotrexate dosing schedules. The improved toxicity profile of methotrexate can make it the preferred agent. Alternative singleagent chemotherapy options include etoposide and 5-fluorouracil. 55,56 Chemotherapy is continued until hCG normalization with an additional 3 cycles of consolidation chemotherapy administered to minimize recurrence.⁵⁷ For patients who do not experience remission or who develop toxicity to methotrexate or actinomycin D, an alternative single agent is often used with excellent results. However, transition to multiagent chemotherapy may be necessary if the hCG level at persistence is particularly elevated or if clear progression of disease is seen.⁵⁸

Role of Second Dilation and Curettage

The potential role of second dilation and curettage rather than chemotherapy has been investigated in several studies with conflicting results. In a recent prospective study, approximately 40% of women undergoing a second dilation and curettage experienced cure without the need for chemotherapy.⁵⁹ FIGO score (5 or 6) and age (>39 or <19 years) have been identified as risk factors for failure of second curettage to avoid the need for chemotherapy.^{59,60}

Role of Hysterectomy

Women with FIGO stage I nonmetastatic GTN who do not desire fertility preservation may be treated with hysterectomy. The best approach (open vs minimally invasive) has not been established. Given the high rate of occult pulmonary metastases, one cycle of single-agent chemotherapy administered perioperatively may be considered to achieve sustained remission. Hysterectomy is also the treatment of choice for women with the relatively chemotherapy resistant histologies of PSTT or ETT. These are uncommon tumors, and PSTT and ETT are rarely diagnosed during postmolar surveillance.

Those diagnosed with PSTT or ETT later than 48 months from antecedent pregnancy are at high risk for recurrence and poor outcome, leading some clinicians to offer postoperative high-dose chemotherapy with stem cell support.64,65

Follow-Up and Future Pregnancy

Regardless of treatment method, women with low-risk GTN should have serum hCG levels monitored monthly for 1 year after remission. During this time, effective birth control must be used. The type of birth control used does not seem to increase rates of recurrence.⁶⁶ GTN and use of chemotherapy do not seem to impact future fertility. Other than an increased risk of second molar pregnancy, subsequent pregnancies after GTN have been shown to have obstetrical outcomes similar to those for women without a history of GTN.67-70 However, for any subsequent pregnancies, we recommend an early trimester ultrasound to assess for normal early development and to check hCG level approximately 6 to 8 weeks postpartum to ensure that it has normalized and a that second trophoblastic event did not develop.

Submitted July 1, 2019; accepted for publication September 30, 2019.

Disclosures: The authors have disclosed that they have no financial interests, arrangements, affiliations, or commercial interests with the manufacturers of any products discussed in this article or their competitors. Research at the New England Trophoblastic Disease Center is supported by the Donald P. Goldstein, MD, Trophoblastic Tumor Registry Endowment and the Dyett Family Trophoblastic Disease Research and Registry Endowment.

Correspondence: Ross S. Berkowitz, MD, Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115. Email: rberkowitz@partners.org

References

- Lurain JR. Gestational trophoblastic disease II: classification and management of gestational trophoblastic neoplasia. Am J Obstet Gynecol 2011:204:11-18.
- Seckl MJ, Sebire NJ, Berkowitz RS. Gestational trophoblastic disease. Lancet 2010;376:717-729.
- Ngan HYS, Seckl MJ, Berkowitz RS, et al. Update on the diagnosis and management of gestational trophoblastic disease. Int J Gynaecol Obstet 2018;143(Suppl 2):79-85.
- Berkowitz RS, Goldstein DP. Clinical practice. Molar pregnancy. N Engl J Med 2009;360:1639-1645.
- Szulman AE, Surti U. The syndromes of hydatidiform mole. I. Cytogenetic and morphologic correlations. Am J Obstet Gynecol 1978;131:665-671.
- Nguyen NMP, Khawaikie Y, Mechtouf N, et al. The genetics of recurrent 6. hydatidiform moles: new insights and lessons from a comprehensive analysis of 113 patients. Mod Pathol 2018;31:1116-1130.
- Moglabey YB, Kircheisen R, Seoud M, et al. Genetic mapping of a maternal locus responsible for familial hydatidiform moles. Hum Mol Genet 1999;8:667-671.
- Soto-Wright V, Bernstein M, Goldstein DP, Berkowitz RS. The changing clinical presentation of complete molar pregnancy. Obstet Gynecol 1995;
- Hou JL, Wan XR, Xiang Y, et al. Changes of clinical features in hydatidiform mole: analysis of 113 cases. J Reprod Med 2008;53:629-633
- 10. Braga A, Moraes V, Maestá I, et al. Changing trends in the clinical presentation and management of complete hydatidiform mole among Brazilian women. Int J Gynecol Cancer 2016;26:984-990.
- 11. Sun SY, Melamed A, Goldstein DP, et al. Changing presentation of complete hydatidiform mole at the New England Trophoblastic Disease Center over the past three decades: does early diagnosis alter risk for gestational trophoblastic neoplasia? Gynecol Oncol 2015;138:46-49.
- Mosher R, Goldstein DP, Berkowitz R, et al. Complete hydatidiform mole: comparison of clinicopathologic features, current and past. J Reprod Med
- 13. Berkowitz RS, Goldstein DP, Bernstein MR. Natural history of partial molar pregnancy. Obstet Gynecol 1985;66:677-681.
- Joneborg U, Marions L. Current clinical features of complete and partial hydatidiform mole in Sweden. J Reprod Med 2014;59:51-55.
- Sun SY, Melamed A, Joseph NT, et al. Clinical presentation of complete hydatidiform mole and partial hydatidiform mole at a regional trophoblastic disease center in the United States over the past 2 decades. Int J Gynecol Cancer 2016;26:367-370.
- Fowler DJ, Lindsay I, Seckl MJ, et al. Routine pre-evacuation ultrasound diagnosis of hydatidiform mole: experience of more than 1000 cases from a regional referral center. Ultrasound Obstet Gynecol 2006;27:56-60.
- Johns J, Greenwold N, Buckley S, et al. A prospective study of ultrasound screening for molar pregnancies in missed miscarriages. Ultrasound Obstet Gynecol 2005;25:493-497.

- Benson CB, Genest DR, Bernstein MR, et al. Sonographic appearance of first trimester complete hydatidiform moles. Ultrasound Obstet Gynecol 2000:16:188-191
- 19. Fine C, Bundy AL, Berkowitz RS, et al. Sonographic diagnosis of partial hydatidiform mole. Obstet Gynecol 1989;73:414-418.
- Seckin KD, Baser E, Yeral I, et al. The impact of ultrasonographic lesion size and initial human chorionic gonadotropin values on treatment success in cases with complete hydatidiform mole. Eur Rev Med Pharmacol Sci 2013;17:3381-3384.
- Padrón L, Rezende Filho J, Amim Junior J, et al. Manual compared with electric vacuum aspiration for treatment of molar pregnancy. Obstet Gynecol 2018;131:652-659.
- Elias KM, Goldstein DP, Berkowitz RS. Complete hydatidiform mole in women older than age 50. J Reprod Med 2010;55:208-212.
- 23. Elias KM, Shoni M, Bernstein M, et al. Complete hydatidiform mole in women aged 40 to 49 years. J Reprod Med 2012;57:254-258.
- 24. Kim DS, Moon H, Kim KT, et al. Effects of prophylactic chemotherapy for persistent trophoblastic disease in patients with complete hydatidiform mole. Obstet Gynecol 1986;67:690-694.
- Limpongsanurak S. Prophylactic actinomycin D for high-risk complete hydatidiform mole. J Reprod Med 2001;46:110-116.
- Wang Q, Fu J, Hu L, et al. Prophylactic chemotherapy for hydatidiform mole to prevent gestational trophoblastic neoplasia. Cochrane Database Syst Rev 2017;9:CD007289.
- Sturgeon CM, Berger P, Bidart JM, et al. Differences in recognition of the 1st WHO international reference reagents for hCG-related isoforms by diagnostic immunoassays for human chorionic gonadotropin. Clin Chem 2009;55:1484-1491.
- 28. Harvey RA, Mitchell HD, Stenman UH, et al. Differences in total human chorionic gonadotropin immunoassay analytical specificity and ability to measure human chorionic gonadotropin in gestational trophoblastic disease and germ cell tumors. J Reprod Med 2010;55:285–295.
- Cole LA, Shahabi S, Butler SA, et al. Utility of commonly used commercial human chorionic gonadotropin immunoassays in the diagnosis and management of trophoblastic diseases. Clin Chem 2001;47:308-315.
- Levavi H, Neri A, Bar J, et al. "Hook effect" in complete hydatidiform molar pregnancy: a falsely low level of beta-HCG. Obstet Gynecol 1993; 82(4 Pt 2, Suppl):720-721.
- Cormano J, Mackay G, Holschneider C. Gestational trophoblastic disease diagnosis delayed by the hook effect. Obstet Gynecol 2015;126:811-814.
- Kohorn El. Persistent low-level "real" human chorionic gonadotropin: a clinical challenge and a therapeutic dilemma. Gynecol Oncol 2002;85: 315-320
- González Aguilera B, Syrios P, Gadisseur R, et al. Persistent low levels of serum hCG due to heterophilic mouse antibodies: an unrecognized pitfall in the diagnosis of trophoblastic disease. Gynecol Endocrinol 2016;32: 439-441

- Cole LA, Khanlian SA, Giddings A, et al. Gestational trophoblastic diseases: 4. Presentation with persistent low positive human chorionic gonadotropin test results. Gynecol Oncol 2006;102:165–172.
- Patel KK, Qavi AJ, Hock KG, et al. Establishing reference intervals for hCG in postmenopausal women. Clin Biochem 2017;50:234–237.
- Committee on Practice Bulletins-Gynecology, American College of Obstetricians and Gynecologists. ACOG Practice Bulletin #53. Diagnosis and treatment of gestational trophoblastic disease. Obstet Gynecol 2004; 103:1365–1377.
- Wolfberg AJ, Feltmate C, Goldstein DP, et al. Low risk of relapse after achieving undetectable HCG levels in women with complete molar pregnancy. Obstet Gynecol 2004;104:551–554.
- Wolfberg AJ, Growdon WB, Feltmate CM, et al. Low risk of relapse after achieving undetectable HCG levels in women with partial molar pregnancy. Obstet Gynecol 2006;108:393–396.
- Braga A, Maestá I, Matos M, et al. Gestational trophoblastic neoplasia after spontaneous human chorionic gonadotropin normalization following molar pregnancy evacuation. Gynecol Oncol 2015;139:283–287.
- Coyle C, Short D, Jackson L, et al. What is the optimal duration of human chorionic gonadotrophin surveillance following evacuation of a molar pregnancy? A retrospective analysis on over 20,000 consecutive patients. Gynecol Oncol 2018;148:254–257.
- 41. FIGO Oncology Committee. FIGO staging for gestational trophoblastic neoplasia 2000. Int J Gynaecol Obstet 2002;77:285–287.
- Agarwal R, Teoh S, Short D, et al. Chemotherapy and human chorionic gonadotropin concentrations 6 months after uterine evacuation of molar pregnancy: a retrospective cohort study. Lancet 2012;379:130–135.
- Braga A, Torres B, Burlá M, et al. Is chemotherapy necessary for patients with molar pregnancy and human chorionic gonadotropin serum levels raised but falling at 6months after uterine evacuation? Gynecol Oncol 2016;143:558–564.
- Edge ST, Greene FL, Byrd DR, et al, eds. AJCC Staging Manual 8th ed. Chicago, IL: American Joint Committee on Cancer; 2017.
- Abu-Rustum NR, Yashar CM, Bean S, et al. NCCN Clinical Practice Guidelines in Oncology: Gestational Trophoblastic Neoplasia. Version 2.2019. Accessed August 9, 2018. To view the most recent version, visit NCCN.org.
- Ngan HY, Bender H, Benedet JL, et al. Gestational trophoblastic neoplasia, FIGO 2000 staging and classification. Int J Gynaecol Obstet 2003; 83(Suppl 1):175–177.
- Gamer EI, Garrett A, Goldstein DP, et al. Significance of chest computed tomography findings in the evaluation and treatment of persistent gestational trophoblastic neoplasia. J Reprod Med 2004;49:411–414.
- Lawrie TA, Alazzam M, Tidy J, et al. First-line chemotherapy in low-risk gestational trophoblastic neoplasia. Cochrane Database Syst Rev 2016;6: CD007102.
- Maestá I, Nitecki R, Horowitz NS, et al. Effectiveness and toxicity of firstline methotrexate chemotherapy in low-risk postmolar gestational trophoblastic neoplasia: the New England Trophoblastic Disease Center experience. Gynecol Oncol 2018;148:161–167.
- Lurain JR, Elfstrand EP. Single-agent methotrexate chemotherapy for the treatment of nonmetastatic gestational trophoblastic tumors. Am J Obstet Gynecol 1995;172:574–579.
- McNeish IA, Strickland S, Holden L, et al. Low-risk persistent gestational trophoblastic disease: outcome after initial treatment with low-dose methotrexate and folinic acid from 1992 to 2000. J Clin Oncol 2002;20: 1838–1844.
- Petrilli ES, Twiggs LB, Blessing JA, et al. Single-dose actinomycin-D treatment for nonmetastatic gestational trophoblastic disease: a prospective phase II trial of the Gynecologic Oncology Group. Cancer 1987; 60:2173–2176.

- Osborne RJ, Filiaci V, Schink JC, et al. Phase III trial of weekly methotrexate or pulsed dactinomycin for low-risk gestational trophoblastic neoplasia: a Gynecologic Oncology Group study. J Clin Oncol 2011;29: 825–831
- Yarandi F, Mousavi A, Abbaslu F, et al. Five-day intravascular methotrexate versus biweekly actinomycin-D in the treatment of low-risk gestational trophoblastic neoplasia: a clinical randomized trial. Int J Gynecol Cancer 2016;26:971–976.
- Matsui H, litsuka Y, Seki K, et al. Comparison of chemotherapies with methotrexate, VP-16 and actinomycin-D in low-risk gestational trophoblastic disease: remission rates and drug toxicities. Gynecol Obstet Invest 1998:46:5–8.
- Sung HC, Wu PC, Yang HY. Reevaluation of 5-fluorouracil as a single therapeutic agent for gestational trophoblastic neoplasms. Am J Obstet Gynecol 1984:150:69–75.
- Lybol C, Sweep FC, Harvey R, et al. Relapse rates after two versus three consolidation courses of methotrexate in the treatment of low-risk gestational trophoblastic neoplasia. Gynecol Oncol 2012;125:576–579.
- Sita-Lumsden A, Short D, Lindsay I, et al. Treatment outcomes for 618 women with gestational trophoblastic tumours following a molar pregnancy at the Charing Cross Hospital, 2000-2009. Br J Cancer 2012;107: 1810–1814.
- Osborne RJ, Filiaci VL, Schink JC, et al. Second curettage for low-risk nonmetastatic gestational trophoblastic neoplasia. Obstet Gynecol 2016; 128:535–542.
- Garner EI, Feltmate CM, Goldstein DP, et al. The curative effect of a second curettage in persistent trophoblastic disease: a retrospective cohort survey. Gynecol Oncol 2005;99:3–5.
- Bolze PA, Mathe M, Hajri T, et al. First-line hysterectomy for women with low-risk non-metastatic gestational trophoblastic neoplasia no longer wishing to conceive. Gynecol Oncol 2018;150:282–287.
- Clark RM, Nevadunsky NS, Ghosh S, et al. The evolving role of hysterectomy in gestational trophoblastic neoplasia at the New England Trophoblastic Disease Center. J Reprod Med 2010;55:194–198.
- Goldstein DP, Berkowitz RS. Prophylactic chemotherapy of complete molar pregnancy. Semin Oncol 1995;22:157–160.
- Horowitz NS, Goldstein DP, Berkowitz RS. Placental site trophoblastic tumors and epithelioid trophoblastic tumors: biology, natural history, and treatment modalities. Gynecol Oncol 2017;144:208–214.
- Frijstein MM, Lok CAR, Short D, et al. The results of treatment with highdose chemotherapy and peripheral blood stem cell support for gestational trophoblastic neoplasia. Eur J Cancer 2019;109:162–171.
- 66. Dantas PRS, Maestá I, Filho JR, et al. Does hormonal contraception during molar pregnancy follow-up influence the risk and clinical aggressiveness of gestational trophoblastic neoplasia after controlling for risk factors? Gynecol Oncol 2017;147:364–370.
- Vargas R, Barroilhet LM, Esselen K, et al. Subsequent pregnancy outcomes after complete and partial molar pregnancy, recurrent molar pregnancy, and gestational trophoblastic neoplasia: an update from the New England Trophoblastic Disease Center. J Reprod Med 2014;59: 188–194.
- Matsui H, litsuka Y, Suzuka K, et al. Early pregnancy outcomes after chemotherapy for gestational trophoblastic tumor. J Reprod Med 2004; 49:531–534.
- Sebire NJ, Fisher RA, Foskett M, et al. Risk of recurrent hydatidiform mole and subsequent pregnancy outcome following complete or partial hydatidiform molar pregnancy. BJOG 2003;110:22–26.
- Williams J, Short D, Dayal L, et al. Effect of early pregnancy following chemotherapy on disease relapse and fetal outcome in women treated for gestational trophoblastic neoplasia. J Reprod Med 2014; 59:248–254.