

# Clinical Features Suggestive of Meningitis in Children: A Systematic Review of Prospective Data

## abstract

**CONTEXT:** Clinical diagnosis of pediatric meningitis is fundamental; therefore, familiarity with evidence underscoring clinical features suggestive of meningitis is important.

**OBJECTIVE:** To seek evidence supporting accuracy of clinical features of pediatric bacterial meningitis.

**METHODS:** A review of Medline, Embase, the Cumulative Index to Nursing and Allied Health Literature, Web of Science, and PubMed was conducted for all articles of relevance. Articles contained prospective data of clinical features in children with laboratory-confirmed bacterial meningitis and in comparison groups of those without it. Two authors independently assessed quality and extracted data to calculate accuracy data of clinical features.

**RESULTS:** Of 14 145 references initially identified, 10 met our inclusion criteria. On history, a report of bulging fontanel (likelihood ratio [LR]: 8.00 [95% confidence interval (CI): 2.4–26]), neck stiffness (7.70 [3.2–19]), seizures (outside febrile-convulsion age range) (4.40 [3.0–6.4]), or reduced feeds (2.00 [1.2–3.4]) raised concern about the presence of meningitis. On examination, jaundice (LR: 5.90 [95% CI: 1.8–19]), being toxic or moribund (5.80 [3.0–11]), meningeal signs (4.50 [2.4–8.3]), neck stiffness (4.00 [2.6–6.3]), bulging fontanel (3.50 [2.0–6.0]), Kernig sign (3.50 [2.1–5.7]), tone up (3.20 [2.2–4.5]), fever of  $>40^{\circ}\text{C}$  (2.90 [1.6–5.5]), and Brudzinski sign (2.50 [1.8–3.6]) independently raised the likelihood of meningitis. The absence of meningeal signs (LR: 0.41 [95% CI: 0.30–0.57]) and an abnormal cry (0.30 [0.16–0.57]) independently lowered the likelihood of meningitis. The absence of fever did not rule out meningitis (LR: 0.70 [95% CI: 0.53–0.92]).

**CONCLUSIONS:** Evidence for several useful clinical features that influence the likelihood of pediatric meningitis exists. No isolated clinical feature is diagnostic, and the most accurate diagnostic combination is unclear. *Pediatrics* 2010;126:952–960

**AUTHORS:** Sarah Curtis, MD, FRCPC,<sup>a,b,c</sup> Kent Stobart, MD, MSc, FRCPC,<sup>b,c</sup> Ben Vandermeer, BSc, MSc,<sup>d</sup> David L. Simel, MD, MHS,<sup>e,f</sup> and Terry Klassen, MD, FRCPC, MSc<sup>a,b,c,d</sup>

<sup>a</sup>Division of Pediatric Emergency Medicine, <sup>b</sup>Department of Pediatrics, University of Alberta, Edmonton, Alberta, Canada; <sup>c</sup>Women and Children's Health Research Institute, Edmonton, Alberta, Canada; <sup>d</sup>Alberta Research Centre for Health Evidence, Department of Pediatrics, University of Alberta, Edmonton, Alberta, Canada; <sup>e</sup>Department of Medicine, Durham Veterans Affairs Medical Center, Durham, North Carolina; and <sup>f</sup>Department of Medicine, Duke University, Durham, North Carolina

### KEY WORDS

bacterial meningitis, children, meta-analysis, systematic review, diagnosis, sensitivity, specificity, likelihood ratio, accuracy, physical examination, history, signs, symptoms

### ABBREVIATIONS

CSF—cerebrospinal fluid

LP—lumbar puncture

QUADAS—Quality Assessment for Diagnostic Accuracy Studies

LR—likelihood ratio

CI—confidence interval

Drs Curtis, Stobart, and Klassen came up with the study concept and design; Drs Curtis and Stobart acquired the data; Dr Curtis, Mr Vandermeer, and Dr Simel analyzed and interpreted the data; Dr Curtis drafted the manuscript; Drs Curtis and Stobart, Mr Vandermeer, and Drs Klassen and Simel critically revised the manuscript for important intellectual content; Drs Curtis and Vandermeer performed statistical analysis; Drs Curtis and Klassen provided administrative, technical, or material support; and Drs Stobart and Klassen supervised the study.

[www.pediatrics.org/cgi/doi/10.1542/peds.2010-0277](http://www.pediatrics.org/cgi/doi/10.1542/peds.2010-0277)

doi:10.1542/peds.2010-0277

Accepted for publication Aug 10, 2010

Address correspondence to Sarah Curtis, MD, FRCPC, Aberhart Centre, Room 7217A, 11402 University Ave, Edmonton, Alberta, Canada T6G 2J3. E-mail: [scurtis@ualberta.ca](mailto:scurtis@ualberta.ca)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

Meningitis can be difficult to diagnose clinically, particularly in young infants who do not seem to reliably display the classic features of the disease. Cerebrospinal fluid (CSF) analysis through lumbar puncture (LP) is the most important laboratory diagnostic test. However, LP is invasive and painful and can be challenging to perform and anxiety-provoking for caregivers. It has been commonly associated with adverse events such as headache and backache and rarely associated with infection, cerebral herniation, and subdural and spinal epidural hemorrhage.<sup>1</sup> Furthermore, CSF analysis is not readily accessible in many regions of the world. Thus, it may not be desirable or feasible to perform an LP on every child who presents with the non-specific symptoms that may be attributable to bacterial meningitis but are much more commonly associated with less serious conditions.

Delay in or failure of diagnosis of meningitis is reflected in reviews of medical malpractice in the pediatric setting. Missed meningitis is the most common diagnosis involved in pediatric emergency malpractice claims and has been associated with the highest median indemnity payments and defense payments for pediatricians.<sup>2,3</sup> Malpractice cases that involve children younger than 2 years and cases in which the child died were most often related to the diagnosis of meningitis. Because incidence rates decline with vaccination uptake, the opportunity for recognition of and familiarity with the clinical features of this disease for practicing physicians and trainees is becoming increasingly rare. However, this devastating disease has an ongoing potential to resurface with occasional outbreaks of known or new organisms.

Ideally, primary clinical assessment should provide an estimate of the probability of disease and help to de-

termine if further diagnostic testing is required. Identification and use of those features that raise the pretest probability of disease in contradistinction to those that do not should improve efficiency and accuracy of clinical assessment. To our knowledge, a systematic synthesis of prospective data pertaining to clinical features suggestive of meningitis has not yet been performed despite the importance of this disease in clinical training and practice.

## METHODS

### Literature Search and Selection

Using a structured search strategy, a review of Medline, Embase, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Web of Science, PubMed, and the Cochrane databases was conducted in June 2009, without time limitations, for all articles of relevance. A meningitis, a diagnostic accuracy, and a pediatric string of search terms were used. Included studies had to describe pertinent historical and physical features of children with LP-confirmed bacterial meningitis and prospectively collected data amenable to calculation of accuracy estimates. Similar data from an LP-negative comparison group also had to be present.

### Assessment of Quality

Two authors assessed quality by using the Quality Assessment for Diagnostic Accuracy Studies (QUADAS)<sup>4</sup> checklist and the guidelines for assigning quality levels of evidence.<sup>5</sup> The QUADAS checklist was developed for quality assessment in systematic reviews of diagnostic-test-accuracy studies. It is a 14-item checklist with “yes,” “no”, or “unclear” options and examines inclusion population, selection criteria, and the descriptions, timing, independence, and blinding of index and reference tests. Studies were also assessed for the execution of the tests, the con-

sistent use of a single good reference standard (LP), availability of results for all patients, and details of CSF analysis.

### Data Extraction

For both signs and symptoms, if the same word was used to describe a clinical finding in multiple studies, it was assumed that the test was similar enough to combine numerically. The decision to combine terms was reached by consensus after consideration of which terms may reasonably be combined without losing their core meaning.


### Data Analysis

The sensitivity, specificity, and likelihood ratios (LRs) with 95% confidence intervals (CIs) were calculated for symptoms and signs. When data were deemed clinically and methodologically similar enough to warrant meta-analysis, Review Manager (RevMan)<sup>6</sup> was used to calculate summary measures using the generic inverse-variance function. Heterogeneity was estimated by using the  $I^2$  statistic, which measures the amount of variance attributable to between-study variance as opposed to within-study variance.<sup>7</sup>

## RESULTS

Figure 1 shows the study flow and selection process. One author screened 14 145 titles and abstracts, which resulted in 760 potentially relevant articles; ultimately, 10 articles met our inclusion criteria (Table 1).<sup>8–17</sup> All studies had a quality level of evidence of 1 or 2 (level 1:  $n = 4$ ; level 2:  $n = 6$ ) and scored  $\geq 10$  on the QUADAS checklist.

CSF analysis was the gold standard for defining the presence of meningitis. The CSF definition of meningitis varied in detail but included a combination of CSF culture positivity or CSF pleocytosis along with either blood culture positivity or CSF latex agglutination posi-



**FIGURE 1**  
Study flow diagram.

tivity (Table 2). Normal CSF test results and negative microbiologic study results excluded bacterial meningitis.

Eighteen symptom descriptors and 48 sign descriptors were found and extracted for meta-analysis. Of these descriptors, only 5 symptoms and 21 signs resulted in significant data (Table 3). Nonsignificant findings for both positive and negative LRs are listed in Table 4.

Features were considered to be signs if described by the physician or symptoms if reported on history by caregivers. No studies evaluated the precision of clinical findings, so the focus of the review was on diagnostic accuracy. Only 2 articles reported combinations of findings.<sup>12,17</sup>

### Prevalence of Meningitis

The study (point) prevalence of meningitis varied widely from region to region (Table 1). The high prevalence of meningitis reflects the selected nature

of the type of patient studied or seasonal outbreaks of particular pathogens in the various regions of the world. Study inclusion criteria represented 2 categories of children: (1) children with seizure and fever<sup>8,9</sup>; and (2) children with a clinical suspicion of invasive bacterial disease or meningitis.<sup>10–17</sup> Thus, the LRs for the following symptoms and signs should be applied only to these child populations.

### Accuracy of Features of the Clinical History Suggestive of Meningitis

When a caregiver reported that his or her child had a bulging fontanel or neck stiffness, the likelihood of meningitis increased nearly eightfold<sup>10,17</sup> (Table 3). If a child had experienced a seizure but the age of the child was outside that of the typical age range for febrile seizures, the likelihood of meningitis was increased fourfold.<sup>12</sup> A lack of irritability lowered the odds of

the disease by half, but the presence of irritability did not strongly signify the presence of meningitis.<sup>17</sup> A child with a history of reduced feeds<sup>12,13,17</sup> had a somewhat increased likelihood of meningitis.

### Accuracy of Features of the Physical Examination Suggestive of Meningitis

#### Seizures

The presence of complex seizures doubled the risk of meningitis<sup>9–11</sup> (Table 3). When the seizure type was listed as “nonspecific”<sup>13,15</sup> or when multiple seizures<sup>8</sup> were described, the likelihood was weaker. Other seizure descriptors were described in primary studies, but data were not statistically significant (Table 4).

#### Meningeal Signs

The definition of “meningeal signs” varied (eg, stiffness or rigidity or meningeal irritation or Brudzinski or Kernig sign), and the presence of any 1 of them had a summary LR of 4.50.<sup>8,11,13,16,17</sup> The absence of meningeal signs was more consistent and decreased the likelihood of meningitis. When meningeal signs were defined only as “neck stiffness,” the results were more heterogeneous, but the LRs were comparable to the more general term. Only Walsh-Kelly et al<sup>16</sup> evaluated Kernig and Brudzinski signs in isolation. The presence of either sign increased the likelihood of meningitis, whereas the absence of either sign lowered the likelihood.

The presence of a bulging fontanel increased the risk of meningitis in an infant 3.5 times, but when absent, the risk of meningitis decreased only slightly.<sup>12,13,15–17</sup>

#### Mental Status or Appearance

The descriptors of a “change in mental status,”<sup>9,12,15,17</sup> “restless or irritable or agitated,”<sup>15,17</sup> “lethargic or

**TABLE 1** Studies That Met Inclusion Criteria for Accuracy of Clinical Features Suggestive of Bacterial Meningitis in Children

Study	Quality Level	Quality QUADAS Score, of 14	Country	Setting	Tester	Population Study Prevalence, <i>n</i> (%) <sup>a</sup>	Age and Age Subgroup	Inclusion Criteria	No. of Index Tests	Description of Index Tests
Akpede et al <sup>8</sup> (1992)	2	12	Benin, Nigeria	Pediatric ED	Unknown	522 (4.2)	1 mo to 6 y; subgroups: 1–6 mo, 6 mo to 2 y, 2–6 y	Seizure and fever	8	Insufficient description
Akpede and Sykes <sup>9</sup> (1993)	2	10	Benin, Nigeria	Pediatric ED	Unknown	92 (15)	6–14 y	Seizure and fever	9	Insufficient description
Akpede <sup>10</sup> (1995)	2	12	Maduguri, Nigeria	Pediatric ED	Unknown	341 (19)	1 mo to 15 y	Fever and seizure or “acutely ill”	15	Insufficient description
Lembo and Marchant <sup>14</sup> (1991)	2	11.5	Cleveland, OH	ED/acute care clinic of hospital	Managing physician	160 (6)	6 mo (median)	Fever and “could have meningitis”	2 groups of 4 signs and 4 symptoms	Insufficient description
Lehmann et al <sup>13</sup> (1999)	2	13	Goroka, Papua, New Guinea	Base hospital	Pediatrician	642 (18)	1 mo to 5 y	Suspected meningitis	14	Insufficient description
Walsh-Kelly et al <sup>16</sup> (1992)	1	13	Wisconsin	Pediatric ED	Pediatric emergency physicians	547 (10)	1 wk to 17 y (mean: 30 mo); subgroups: 0–6 mo, 7–12 mo, 13–18 mo, >18 mo	All children undergoing LP	8	Well-defined index tests
Weber et al <sup>17</sup> (2002)	1	13	Banjul, the Gambia	Pediatric hospital	Attending physicians	322 (16)	2–33 mo	Suspected invasive bacterial disease and member of vaccine trial	26	Insufficient description
Berkley et al <sup>11</sup> (2001)	1	13	Kilifi, Kenya	District hospital	Research clinician	906 (5)	1 mo to 13 y	Admitted children with suspected meningitis	3	Insufficient description
Berkley et al <sup>12</sup> (2004)	1	13	Kilifi, Kenya	District hospital	Clinical officers and pediatric residents	999 (9)	3–40 mo (median: 22 mo)	All pediatric admissions with any signs of meningitis, impaired consciousness and prostration, and seizures (other than simple febrile)	30	Insufficient description
Sigauque et al <sup>15</sup> (2008)	2	12	Manhiça, Mozambique	District hospital	Trained medical officer	475 (15)	0–15 y (13% neonatal)	Admitted children with neurologic signs compatible with meningitis	9	Insufficient description

ED indicates emergency department.

<sup>a</sup> *N* population with both LP results and index-test results available.

drowsy,”<sup>9,17</sup> or being “unconscious or comatose”<sup>8–11,13,16,17</sup> had comparably weak summary LRs that ranged from 1.40 to 1.90. A “toxic or moribund” appearance had a high LR of 5.80, the absence of which would halve the risk of meningitis.<sup>16</sup> The presence of an “abnormal cry” increased the likelihood of meningitis, but its absence had a larger impact on likelihood of meningitis (LR: 0.30).<sup>17</sup>

### Other Miscellaneous Signs

The presence of a high fever ( $\geq 40^{\circ}\text{C}$ )<sup>9,10</sup> was useful with a summary LR of 2.90, but the LR for temperatures of  $< 40^{\circ}\text{C}$  (or not otherwise specified) had a CI that included 1.00. It should be noted that the absence of fever did not rule out meningitis.<sup>13,15,17</sup>

Several other signs have each been evaluated in only 1 study, and their LR results require validation. Among 341 patients with a meningitis prevalence of 19%, the only patients with petechiae ( $n = 4$ ) all had meningitis.<sup>10</sup> Similarly, the presence of jaundice was also notable as a sign of meningitis (positive LR of 5.90) but was less useful for ruling out the disease.<sup>17</sup>

“Tone up” had a clinically useful LR of 3.20.<sup>17</sup> The absence of high tone reduced the likelihood of meningitis by half. The feature of having “staring eyes” had an LR of 2.40, the absence of which only decreased the likelihood of disease by one-third.<sup>13</sup> “Can’t or won’t feed” seemed to be clinically useful with an LR of 2.10, whereas normal feeding reduced the likelihood of meningitis somewhat.<sup>17</sup>

## DISCUSSION

Information on efficient use of clinical findings is extremely important for clinicians. Useful features for estimation of probability of meningitis are those features that demonstrate the strongest LRs for presence or absence of disease. The LR of a clinical feature is

**TABLE 2** LP (Gold Standard) Definitions Used in Each Study

Study	Bacterial Meningitis	Presumed Bacterial Meningitis	Aseptic	No Meningitis
Akpede et al <sup>8</sup>	> 5 WBCs per $\mu$ L CSF and bacterium on CSF Gram-stain and/or CSF culture	No bacterium in CSF; > 5 WBCs per $\mu$ L and protein > 80 mg/dL and glucose < 40% blood glucose in CSF	—	—
Akpede and Sykes <sup>9</sup>	> 5 WBCs per $\mu$ L CSF and bacterium on CSF Gram-stain and/or CSF culture	—	—	—
Akpede <sup>10</sup>	Cloudy/turbid CSF, > 10 WBCs per $\mu$ L, organism on Gram-stain/CSF culture and blood culture, and CSF/blood glucose < 50% and CSF protein > 0.8 g/L	—	—	—
Lembo and Marchant <sup>14</sup>	Bacterium in CSF culture or bacterium-specific Gram-stain antigen detection in CSF	—	CSF pleocytosis > 10 cells per $\mu$ L and < 1000 RBCs per $\mu$ L, sterile blood and CSF cultures and negative CSF Gram-stain and bacterial antigen tests when no oral antibiotic treatment	—
Lehmann et al <sup>13</sup>	Bacterium in CSF culture	CSF culture-negative but bacterium in blood culture or Hib antigen in urine and either CSF glucose < 50 mg/dL or CSF WBCs > 7 $\mu$ L with 70% polymorphs	—	—
Walsh-Kelly et al <sup>16</sup>	CSF latex agglutination-positive or Gram-stain-positive or CSF culture-positive	—	CSF WBCs > 10/ $\mu$ L neonate; > 5/ $\mu$ L child with negative CSF Gram-stain and/or latex agglutination and normal CSF culture	—
Weber et al <sup>17</sup>	CSF culture-positive	CSF culture-negative; antigen detection-positive or WBCs > 5/ $\mu$ L	—	CSF culture-negative, CSF antigen-negative, WBCs < 5/ $\mu$ L
Berkley et al <sup>11</sup>	CSF culture-positive or positive latex agglutination test or organism on CSF Gram-stain	CSF WBCs > 5/ $\mu$ L or CSF/blood glucose ratios of $\leq$ 0.1	—	CSF WBCs $\leq$ 10/ $\mu$ L, glucose CSF/blood $\geq$ 0.67, CSF protein $\leq$ 0.45 g/L
Berkley et al <sup>12</sup>	Positive CSF culture or positive CSF latex agglutination test or bacteria seen on Gram-stain or a CSF leukocyte count of > 50 cells per $\mu$ L or a CSF/blood glucose ratio of < 0.1	—	—	—
Sigauque et al <sup>15</sup>	Bacterial growth in CSF culture	—	—	—

WBCs indicates white blood cells; RBCs, red blood cells; Hib, *Haemophilus influenzae* type b.

the probability of that finding in patients with disease divided by the probability of the same feature in patients without disease (LRs range from 0 to infinity). Features with LR equal to 1.00 have no diagnostic value, because it is equally likely to find the feature in those with the disease as in those without the disease. Features with LR of > 1.00 support the diagnosis of interest in magnitude of increasing numerical value. For features with LR between 0 and 1.00, the smaller the LR, the less likely the disease.<sup>18,19</sup>

Valuable features found in this review are listed in Table 3. On history, in order of decreasing magnitude, a caregivers' report of neck stiffness, bulging fontanel, seizures (outside the febrile-convulsion range), or reduced feeds raise concern about the presence of meningitis. On physical examination, in order of decreasing magnitude, the presence of jaundice, being toxic or moribund, or having meningeal signs, neck stiffness, bulging fontanel, Kernig sign, tone up, fever of > 40°C, or Brudzinski sign all raise the probability of meningitis to varying degrees in the patient. Several other clinical features with LR between 1.30 and 2.40, are less strong but warrant further study. Note that the sign petechiae is strong with an LR of 37.00 but was surprisingly only examined in a single small prospective study, and only 4 patients displayed the feature. Thus, relevance of this well-known feature is currently uncertain, and systematic prospective evaluations of it among large numbers of patients would provide clarity.

As an example of applicability, assuming statistical independence, a pretest probability of disease of 10%, and using the LR nomogram,<sup>20</sup> a combination of the presence of meningeal signs (LR: 4.50), a bulging fontanel (LR: 3.50), and a high fever (LR: 2.90) (thus, a combined LR of 45.60) raises an infant's



**TABLE 3** Accuracy of Clinical Features

	No. of Patients	Summary Sensitivity, %	Summary Specificity, %	Summary LR <sup>+</sup> (95% CI)	I <sup>2</sup> , %	Summary LR <sup>−</sup> (95% CI)	I <sup>2</sup> , %
<b>Symptom</b>							
Bulging fontanel <sup>17</sup>	284	14	98	8.0 (2.40–26.00)	—	0.88 (0.79–0.98)	—
Neck stiffness or bulging fontanelle <sup>10</sup>	341	20	98	7.7 (3.20–19.00)	—	0.82 (0.73–0.93)	—
History of seizures outside febrile-convulsions age range <sup>12</sup>	965	32	93	4.4 (3.00–6.40)	—	0.73 (0.64–0.85)	—
Reduced feeds <sup>12,13,17</sup>	1890	52	70	2.0 (1.20–3.40)	92	0.66 (0.54–0.81)	54
Irritability <sup>17</sup>	284	82	34	1.3 (1.10–1.50)	—	0.52 (0.28–0.97)	—
<b>Sign</b>							
Petechiae <sup>10</sup>	341	6	100	37 (2.00–680.00)	—	0.94 (0.88–0.99)	—
Jaundice <sup>17</sup>	2059	6	99	5.9 (1.80–19.00)	—	0.95 (0.89–1.00)	—
Toxic or moribund <sup>16</sup>	172	49	92	5.8 (3.00–11.00)	—	0.56 (0.42–0.73)	—
Meningeal signs <sup>8,11,13,16,17a</sup>	2399	64	89	4.5 (2.40–8.30)	94	0.41 (0.30–0.57)	69
Neck stiffness <sup>11–13,15–17</sup>	3118	51	89	4.0 (2.60–6.30)	90	0.56 (0.43–0.72)	87
Bulging fontanel <sup>12,13,15–17</sup>	2247	36	90	3.5 (2.00–6.00)	84	0.74 (0.61–0.89)	82
Kernig sign <sup>16</sup>	172	53	85	3.5 (2.10–5.70)	—	0.56 (0.41–0.75)	—
Tone up <sup>17</sup>	284	59	82	3.2 (2.20–4.50)	—	0.50 (0.36–0.70)	—
Fever > 40°C <sup>9,10</sup>	433	19	93	2.9 (1.60–5.50)	62	0.81 (0.55–1.20)	66
Brudzinski sign <sup>16</sup>	172	66	74	2.5 (1.80–3.60)	—	0.46 (0.31–0.68)	—
Staring eyes <sup>13</sup>	640	42	82	2.4 (1.80–3.20)	—	0.70 (0.60–0.82)	—
Can't or won't feed <sup>17</sup>	284	61	70	2.1 (1.50–2.80)	—	0.56 (0.39–0.79)	—
Complex seizures <sup>9–11b</sup>	1400	27	82	2.0 (1.20–3.40)	84	0.86 (0.70–1.10)	45
Lethargic or drowsy <sup>8,17</sup>	376	40	79	1.9 (1.30–2.90)	48	0.58 (0.20–1.70)	67
Unconscious or coma <sup>8–11,13,15–17</sup>	3313	23	86	1.8 (1.20–2.70)	69	0.94 (0.85–1.10)	83
Abnormal cry <sup>17</sup>	284	84	52	1.8 (1.50–2.10)	—	0.30 (0.16–0.57)	—
Restless/irritable/agitated <sup>15,17</sup>	758	37	79	1.6 (1.20–2.10)	0	0.77 (0.50–1.20)	81
Multiple seizures <sup>8</sup>	522	64	57	1.5 (1.10–2.10)	—	0.62 (0.36–1.30)	—
Seizures, nonspecific <sup>13,15</sup>	1095	54	63	1.4 (1.20–1.70)	0	0.75 (0.48–1.20)	87
Change in mental status <sup>9,12,15,17c</sup>	1815	72	47	1.4 (1.20–1.70)	74	0.54 (0.34–0.87)	71
Fever (°C not otherwise specified) <sup>15–17</sup>	885	76	34	1.2 (0.98–1.40)	62	0.7 (0.53–0.92)	66

I<sup>2</sup> is a measure of heterogeneity.

<sup>a</sup> Stiffness or rigidity or meningeal irritation or Brudzinski or Kernig sign.

<sup>b</sup> Focal or multiple or >15-minute duration or complex.

<sup>c</sup> Lethargic/agitated/impaired consciousness.

probability of meningitis to 84%. Although the presence or absence of these findings, in combination or separately, hardly confirms or refutes a diagnosis of meningitis, they raise the probability high enough that an LP must be performed.

Each physician routinely incorporates a sense of the probability of disease through careful consideration of the clinical assessment, experience, and estimates of disease prevalence in the population. All of the studies included patients with a suspicion of meningitis or severe illness. The point prevalence of meningitis ranged from 4.2% to 19% across these studies; each prevalence reflects the clinical impression of possible meningitis (via initial inclusion in each study). The summary prevalence of these studies is 10%. This summary

prevalence could be viewed as the posttest probability of the overall clinical examination, because all of the children were judged sick enough to undergo definite testing for meningitis. Assuming a prevalence of disease of 1%, the LR for the clinical impression of meningitis as its own independent “test” would be 11.00. Thus, the clinical suspicion of disease that a health care provider derives from clinical history and examination may, in itself, be a useful test that warrants follow-through to further diagnostic testing. However, although necessary for rapid comprehensive synthesis of complex clinical information, much is unknown about the process of clinical judgment and decision-making. Clinical impressions are prone to error, and efforts to minimize error by maxi-

mizing pretest probability through accurate clinical prediction or decision rules will offer improved patient care.<sup>21–25</sup>

It seems clinically sensible that the combinations of some findings listed in Table 3 would have a greater impact on the probability of meningitis than the individual findings. Only 2 studies examined combinations of findings. It is unfortunate that original subject data from statistical models used in these studies were unavailable; thus, LRs could not be calculated. Nonetheless, Weber et al<sup>17</sup> and Berkley et al<sup>11,12</sup> had constructed logistic regression models of varying combinations of features in an attempt to obtain sets of predictor variables with an optimal balance of sensitivities and specificities. The best combination model in the

**TABLE 4** Unsupported Features of Pediatric Meningitis: Clinical Features From Prospective Studies With Statistically Insignificant Results

Symptoms (13)	Signs (28)
Lethargic or drowsy <sup>10</sup>	Simple seizures <sup>10</sup>
Cough <sup>13,17</sup>	Focal seizures <sup>8,9</sup>
Cyanosis <sup>12,13</sup>	Fever not otherwise specified <sup>13,15,17</sup>
Family history of seizure <sup>9</sup>	Tachypnea <sup>13,17</sup>
History of seizures outside febrile-convulsion age range <sup>12</sup>	Chest indrawing <sup>13,17</sup>
History of difficulty breathing <sup>17</sup>	Low oxygen <sup>17</sup>
History of vomiting <sup>12,17</sup>	Shock <sup>10,16</sup>
History of diarrhea <sup>10</sup>	Severe malnutrition <sup>10,15</sup>
Fever for <3 d <sup>10</sup>	Dehydration <sup>10</sup>
Fever for >3 d <sup>10</sup>	Age 1–6 mo <sup>10</sup>
Male gender <sup>10</sup>	Age <2 y <sup>10</sup>
Female gender <sup>10</sup>	Age 6–10 y <sup>10</sup>
Chest pain <sup>17</sup>	Age 10–14 y <sup>10</sup>
	Palpable spleen <sup>15</sup>
	Palmar pallor <sup>11,17</sup>
	Tachypnea <sup>13,17</sup>
	Cyanosis <sup>13</sup>
	Crepitations <sup>17</sup>
	Delayed capillary refill <sup>12</sup>
	Hypothermia <sup>12</sup>
	Respiratory distress <sup>13,17</sup>
	Palpable temperature gradient <sup>12</sup>
	Severe wasting <sup>12</sup>
	Malaria parasite on slide <sup>12,17</sup>
	Extracranial focal infection <sup>8,9</sup>
	Appears sick <sup>17</sup>
	Opisthotonos <sup>15</sup>

These features were examined in 1 or more articles of this review as referenced, but data were statistically insignificant (CIs for LRs crossed 1).

Weber et al study,<sup>17</sup> which combined a history of seizures, being lethargic or unconscious, or having a stiff neck, had a sensitivity of 98% and specificity of 70%.<sup>17</sup> This combination of features is a simplified Integrated Management of Childhood Illness referral criteria, a set of guidelines initially developed by the World Health Organization to identify sick children in need of referral.<sup>17,26</sup> However, Berkley et al<sup>11</sup> later tested this same model and found it to be only 85% sensitive and 59% specific. Further models from Berkley et al<sup>12</sup> included 1 with a high sensitivity of 97% but low specificity of 44% and combined nonmalarious fever with any 1 of the following: bulging fontanel; neck stiffness; cyanosis; seizures (outside of febrile seizure age range); partial seizures; and impaired consciousness. Another model combined impaired consciousness with any 1 of the following: bulging fontanel; neck stiffness;

partial seizure; cyanosis; seizure (outside of febrile-seizure age range); it was found to be less sensitive (79%) but more specific (80%).<sup>12</sup> With a life-threatening highly morbid condition, diagnostic models that maximize sensitivity are essential. However, population overassessment, resulting from application of low-specificity models, is also of concern, particularly for regions in which distance or resource restrictions limit access to further care. Thus, the ideal clinical model for pediatric meningitis is still unclear, and prospective evaluation and validation of known and new prediction models in varying populations are imperative. Although many of the symptoms and signs with available data demonstrated poor accuracy (Table 4), these findings have not been otherwise studied in combination. In addition, many other widely described features, otherwise reported in textbooks or review

articles, have not been examined for validity in prospective studies. These commonly described clinical features warrant further prospective examination to confirm soundness of continued use in the context of meningitis. When considering the results of this systematic review, clinicians should remain prudent regarding decision-making for young infants and particularly should not rely on the absence of archetypal features as reassurance of absence of disease. Several investigators from the included studies noted infants with meningitis who displayed few or no classic features of the disease. It is well accepted clinically that young infants with nonspecific yet concerning features such as fever, lethargy, poor feeding, or irritability, among others, must be approached with a high index of suspicion regardless of how well they appear, because the incidence of serious bacterial infection in this age group is much higher than that in older infants.

**LIMITATIONS**

This review was limited by heterogeneity in study settings, patient age, comorbidities, inclusion criteria, gold standard, and index-test definitions. However, the weight of each of these features on clinical heterogeneity is variable and uncertain. All studies were similar in that they examined unwell children initially encountered as outpatients in hospital emergency departments or hospital acute care clinics. All children had a spectrum of illness that raised the suspicion of meningitis, none were pretreated with antibiotics, and all had LPs performed. Nevertheless, the degree of tolerance for increasing heterogeneity must be balanced with potential diminution of accuracy in overall summary measures. Results of this meta-analysis should be applied with prudent consideration of its limitations and to patient

populations that resemble those of the included studies (Table 1).

Ideally, meta-analyses of clinical features in pediatrics would provide accurate summary reports of the usefulness of clinical features in clinically relevant age groups reflective of changing pediatric physiology. It is unfortunate that this meta-analysis can only provide single summary data for the child (age not defined), because precise age categorization of findings were either absent or dissimilar. This leaves uncertainty, for example, as to when the examination of an older child begins to reflect that of an adult or how the examination of a neonate differs from that of an older infant.

Other notable limitations are the insufficient a priori definitions of the individual clinical findings. When viewed as separate diagnostic “tests” each clinical feature, as in any diagnostic-accuracy study, requires precise definitions to ensure reproducibility and a standardized interpretability. For example, neck stiffness may have varied from slightly stiff or tender for 1 set of researchers to rigid for other researchers. Tone up may mean increased muscle tone or hypertonicity,

but it was not specifically defined in the original article. Even fever had variable descriptions, and the finding showed no utility when it was not quantified by actual temperature. For future research, careful attention must be paid to clear definitions and precision ratings of clinical findings to standardize performance of the physical examination and ensure reproducibility.

## CONCLUSIONS

Several useful clinical features that are more likely to be present in children with meningitis compared with those without disease have been identified and are supported, with limitations, by prospectively collected data. Many other described features of meningitis are currently unsupported by available data and warrant further definitive examination. No clinical feature is diagnostic in isolation, and the most accurate combination of clinical features to raise or lower suspicion of meningitis is still unclear.

## ACKNOWLEDGMENTS

There was no external funding obtained for the design or conduct of the study; collection, management, analysis, or interpretation of the data; or

preparation, review, or approval of the manuscript. Dr Curtis had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

We thank Lisa Tjosvold, BA, MLIS (Alberta Research Centre for Health Evidence) for assistance with the literature search; Belinda Allan and Lisa Chambers (Division of Pediatric Emergency Medicine, Department of Pediatrics, University of Alberta) for help with retrieval of relevant articles; and Clay Bordley, MD, MPH (Division of Hospital and Emergency Medicine, Department of Pediatrics, School of Medicine, Duke University Medical Center, Durham, NC), Dennis A. Clements, MD, PhD, MPH (Duke Children's Primary Care, Duke Global Health Institute, Center for Latin American and Caribbean Studies, Duke University), and Rose Hatala, MD (Department of Medicine, St Paul's Hospital, University of British Columbia, Vancouver, British Columbia, Canada) for valuable advice on earlier versions of the manuscript. None of the acknowledged individuals received compensation for their contributions.

## REFERENCES

1. Straus SE, Thorpe KE, Holroyd-Leduc J. How do I perform a lumbar puncture and analyze the results to diagnose bacterial meningitis? *JAMA*. 2006;296(16):2012–2022
2. Selbst SM, Friedman MJ, Singh SB. Epidemiology and etiology of malpractice lawsuits involving children in US emergency departments and urgent care centers. *Pediatr Emerg Care*. 2005;21(3):165–169
3. Carroll AE, Buddenbaum JL. Malpractice claims involving pediatricians: epidemiology and etiology [published correction appears in *Pediatrics*. 2007;120(4):935]. *Pediatrics*. 2007;120(1):10–17
4. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol*. 2003;3:25
5. Holleman DR, Simel DL. Does the clinical examination predict airflow limitation [published correction appears in *JAMA*. 1995;273(17):1334]? *JAMA*. 1995;273(4):313–319
6. *Review Manager (RevMan)*. Copenhagen, Denmark: Nordic Cochrane Centre; 2008
7. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557–560
8. Akpede GO, Sykes RM, Abiodun PO. Indications for lumbar puncture in children presenting with convulsions and fever of acute onset: experience in the Children's Emergency Room of the University of Benin Teaching Hospital, Nigeria. *Ann Trop Paediatr*. 1992;12(4):385–389
9. Akpede GO, Sykes RM. Convulsions with fever of acute onset in school-age children in Benin City, Nigeria. *J Trop Pediatr*. 1993;39(5):309–311
10. Akpede GO. Presentation and outcome of sporadic acute bacterial meningitis in children in the African meningitis belt: recent experience from northern Nigeria highlighting emergent factors in outcome. *West Afr J Med*. 1995;14(4):217–226
11. Berkley JA, Mwangi I, Ngetsa CJ, et al. Diagnosis of acute bacterial meningitis in children at a district hospital in sub-Saharan Africa. *Lancet*. 2001;357(9270):1753–1757
12. Berkley JA, Versteeg AC, Mwangi I, Lowe BS, Newton CR. Indicators of acute bacterial meningitis in children at a rural Kenyan district hospital. *Pediatrics*. 2004;114(6). Available at: [www.pediatrics.org/cgi/content/full/114/6/e713](http://www.pediatrics.org/cgi/content/full/114/6/e713)
13. Lehmann D, Yeka W, Rongap T, et al. Aetiology and clinical signs of bacterial meningitis in children admitted to Goroka Base Hospital, Papua New Guinea, 1989–1992. *Ann Trop Paediatr*. 1999;19(1):21–32



14. Lembo RM, Marchant CD. Acute phase reactants and risk of bacterial meningitis among febrile infants and children. *Ann Emerg Med.* 1991;20(1):36–40
15. Sigaúque B, Roca A, Sanz S, et al. Acute bacterial meningitis among children, in Manhica, a rural area in southern Mozambique. *Acta Trop.* 2008;105(1):21–27
16. Walsh-Kelly C, Nelson DB, Smith DS, et al. Clinical predictors of bacterial versus aseptic meningitis in childhood. *Ann Emerg Med.* 1992;21(8):910–914
17. Weber MW, Herman J, Jaffar S, et al. Clinical predictors of bacterial meningitis in infants and young children in the Gambia. *Trop Med Int Health.* 2002;7(9):722–731
18. Grimes DA, Schulz KF. Refining clinical diagnosis with likelihood ratios. *Lancet.* 2005;365(9469):1500–1505
19. McGee S. Simplifying likelihood ratios. *J Gen Intern Med.* 2002;17(8):646–649
20. Fagan TJ. Nomogram for Bayes's theorem. *N Engl J Med.* 1975;293(5):257
21. Berner ES, Graber ML. Overconfidence as a cause of diagnostic error in medicine. *Am J Med.* 2008;121(5 suppl):S2–S23
22. Cook C. Is clinical gestalt good enough? *J Man Manip Ther.* 2009;17(1):6–7
23. Croskerry P, Norman G. Overconfidence in clinical decision making. *Am J Med.* 2008;121(5 suppl):S24–S29
24. Kempainen RR, Migeon MB, Wolf FM. Understanding our mistakes: a primer on errors in clinical reasoning. *Med Teach.* 2003;25(2):177–181
25. Klein JG. Five pitfalls in decisions about diagnosis and prescribing. *BMJ.* 2005;330(7494):781–783
26. Tulloch J. Integrated approach to child health in developing countries. *Lancet.* 1999;354(suppl 2):S116–S120

**Would You Read Here or There?:** Like the protagonist in *Green Eggs and Ham*, Americans are being offered many choices as to how and where they can read. And just like in Dr Seuss's book, they are discovering that they have been missing something good. As reported in *The Wall Street Journal* (Fowler G, August 25, 2010), Americans using e-readers are reading more than ever. In a survey of 1200 e-reader owners, 40% reported reading more after purchasing an e-reader while only 2% reported reading less. Though only about 11 million Americans own one of the 3 common e-readers, Amazon's Kindle, Apple's iPad, or Sony's Reader, the news is a welcome departure from a 2007 study which reported that Americans were spending less time reading books. Famously, almost half of young adults 18-24 reported having not read any books for pleasure. E-reader owners not only increased their purchases of e-books over the past year but also hardcover books. Overall, owners of e-readers read 2.6 books a month compared to 1.9 for print book readers. The increased popularity of e-books is reflected in national sales. In 2009, print book sales in the US fell 51% compared to a 1.9% increase in e-book sales. While print books don't have to be put away during takeoff or landing of an airplane, e-readers come with sample chapters to try before purchase, back-lighting that allow for reading in the dark, and importantly for many, text size selection. While I still would not recommend reading with a mouse, reading most anywhere can be fun.

Noted by JFL, MD and WVR, MD

## Clinical Features Suggestive of Meningitis in Children: A Systematic Review of Prospective Data

Sarah Curtis, Kent Stobart, Ben Vandermeer, David L. Simel and Terry Klassen  
*Pediatrics* 2010;126;952; originally published online October 25, 2010;  
DOI: 10.1542/peds.2010-0277

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/126/5/952.full.html>

### References

This article cites 24 articles, 4 of which can be accessed free at:  
<http://pediatrics.aappublications.org/content/126/5/952.full.html#ref-list-1>

### Citations

This article has been cited by 7 HighWire-hosted articles:  
<http://pediatrics.aappublications.org/content/126/5/952.full.html#related-urls>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Infectious Diseases**  
[http://pediatrics.aappublications.org/cgi/collection/infectious\\_diseases\\_sub](http://pediatrics.aappublications.org/cgi/collection/infectious_diseases_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://pediatrics.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://pediatrics.aappublications.org/site/misc/reprints.xhtml>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Clinical Features Suggestive of Meningitis in Children: A Systematic Review of Prospective Data**

Sarah Curtis, Kent Stobart, Ben Vandermeer, David L. Simel and Terry Klassen

*Pediatrics* 2010;126;952; originally published online October 25, 2010;

DOI: 10.1542/peds.2010-0277

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/126/5/952.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

