

A. Exact Heat-Flow Jensen PDE ($m = 1$) Across (γ_9, γ_{10})
1e6
Rescaled Jensen invariants
Spectral height $t \in [47.2, 50.6]$ across (γ_9, γ_{10})
$e^{\pi t/2} c_1(t; 0)$
$e^{\pi t/2} c_3(t; 0)$
$e^{\pi t/2} \partial_\tau c_1(t; 0)$
$e^{\pi t/2}[\frac{3}{4} c_3 - \frac{1}{8}\partial_t^2 c_1]$
B. Exact $(m-1)^2/2$ Multiple-Zero Lift ($P_m = e^{-\frac{\tau}{4}\partial_t^2} t^m$)
Coalescence lift ratio $\partial_\tau c_{2m-3}/c_{2m-1}$
Zero multiplicity m of heat polynomial $P_m(t, \tau)$
Exact ratio $\frac{\partial_\tau c_{2m-3}(t_0; \tau_{coal})}{c_{2m-1}(t_0; \tau_{coal})} = \frac{(m-1)^2}{2}$
Falsified incomplete-index expression $m/4$
C. Backward Coalescence vs. Forward Repulsion ($\tau = t_{DBN}$)
$e^{\pi t/4}\Xi_\tau(t)$
Spectral height t
$\tau = t_{DBN} = -1.60$ (coalesced)
$\tau = t_{DBN} = -1.31 \approx \tau_{coal}$
$\tau = t_{DBN} = 0.00$ (γ_9, γ_{10})
$\tau = t_{DBN} = +1.20$ (repelled)
D. Lehmer Coalescence Scale $|t_{DBN,coal}| \sim 2\delta^2$ (CNV Regime)
Backward coalescence scale $|t_{DBN,coal}|$
Half-gap $\delta = \frac{1}{2}(\gamma_{k+1} - \gamma_k)$
(γ_9, γ_{10})
(γ_{13}, γ_{14})
($\gamma_{127}, \gamma_{128}$)
($\gamma_{135}, \gamma_{136}$)
($\gamma_{6709}, \gamma_{6710}$)
Asymptotic law $|t_{DBN,coal}| = |\tau_{coal}| = 2\delta^2$
Jensen ratio $\frac{2c_1(t_{mid}; 0)}{c_3(t_{mid}; 0)}$