DE BRANGES REPRODUCING KERNELS, THE UNIVERSAL TRANSVERSE SQUARE-CASCADE FOR LEHMER PAIRS, AND THE LEIBNIZ–WRONSKIAN IDENTITY FOR RIEMANN'S Ξ-FUNCTION

THIJS MISSIAEN

Abstract. We establish three exact structural identities for the parity-reduced transverse Wronskian $\widetilde{W}(d,t) = \frac{1}{2d}\partial_d|\Xi(t-\mathrm{i}d)|^2$ of Riemann's Ξ-function. First, for the linear Hermite–Biehler family $E_\alpha(z) = \Xi(z) - \mathrm{i}\alpha\Xi'(z)$ $(\alpha > 0)$, we prove that the diagonal De Branges reproducing kernel satisfies $K_\alpha(t-\mathrm{i}d, t-\mathrm{i}d) \equiv \frac{\alpha}{\pi}\widetilde{W}(d,t)$ globally, decoupling local off-critical zero exclusion from the global Sonine positivity obstructions of Conrey and Li (2000). Second, at any critical-line zero of multiplicity $m \geq 1$, all lower transverse dipole moments $c_1(t_0), \dots, c_{2m-3}(t_0)$ vanish while $c_{2m-1}(t_0) = \frac{m}{(m!)^2}(\Xi^{(m)}(t_0))^2 > 0$, forming an isotropic three-square bowl $\widetilde{W}(d, t_{\mathrm{mid}}+\tau) = \frac{1}{2}(\Xi''(t_{\mathrm{mid}}))^2(\delta^2+\tau^2+d^2) + \mathcal{O}((\delta+|\tau|+d)^3) > 0$ across any Lehmer pair of half-gap $\delta > 0$. Third, for any real-analytic multiplier $\mathcal{M}(z)$, we prove the cross-term-free Leibniz–Wronskian identity $\widetilde{W}_{\mathcal{M}}(d,t) \equiv |\mathcal{M}(t-\mathrm{i}d)|^2\widetilde{W}(d,t) + |\Xi(t-\mathrm{i}d)|^2\widetilde{W}_{1,\mathcal{M}}(d,t)$ and combine it with a 0%-overshoot exponential prime-gauge sieve to close the Lehmer uncertainty zone under the explicit Gabcke–Olver remainder bound.

1. Introduction

The non-vanishing of Riemann's completed zeta function $\xi(s)$ off the critical line $\mathrm{Re}(s) = \frac{1}{2}$ is governed by the transverse radial monotonicity of the modulus squared $|\Xi(t-\mathrm{i}d)|^2$, where $s = \frac{1}{2}+d+\mathrm{i}t$ and $z = -\mathrm{i}(s-\frac{1}{2}) = t-\mathrm{i}d$. In Papers I–III of this series [10, 11, 12], the parity-reduced transverse Wronskian $\widetilde{W}(d,t) := d^{-1}W(d,t) = \frac{1}{2d}\partial_d|\Xi(t-\mathrm{i}d)|^2$ was introduced and analyzed via complex Wigner saddle-point coalescence and harmonic nodal-chamber geometry. In this paper, we establish three exact structural identities linking $\widetilde{W}(d,t)$ to De Branges reproducing kernels [5, 6, 3], critical-line zero coalescence across Lehmer pairs [15, 9, 4], and multiplicative prime-sieve gauges under explicit Riemann–Siegel remainder bounds [8, 2]:

Theorem 1.1 (De Branges kernel identity, square-cascade, and Leibniz–Wronskian framework)**.** *Let $\Xi(z) = \xi(\frac{1}{2}+\mathrm{i}z)$ and let $\widetilde{W}(d,t) = \frac{1}{2d}\partial_d|\Xi(t-\mathrm{i}d)|^2 = \sum_{n=1}^{\infty} c_{2n-1}(t)\, d^{2n-2}$. The following three structural properties hold:*

(1) ***Exact De Branges reproducing kernel identity:*** *For every $\alpha > 0$, let $E_\alpha(z) := \Xi(z) - \mathrm{i}\alpha\Xi'(z)$ and $E_\alpha^{\#}(z) := \overline{E_\alpha(\bar{z})} = \Xi(z) + \mathrm{i}\alpha\Xi'(z)$. At every point $z = t-\mathrm{i}d \in \mathbb{C}\setminus\mathbb{R}$ $(d \neq 0)$, the diagonal reproducing kernel satisfies the exact global identity*

$$K_\alpha(t-\mathrm{i}d, t-\mathrm{i}d) := \frac{|E_\alpha(t-\mathrm{i}d)|^2 - |E_\alpha^{\#}(t-\mathrm{i}d)|^2}{4\pi d} \equiv \frac{\alpha}{\pi}\,\widetilde{W}(d,t), \tag{1.1}$$

while any finite M-point Cauchy–Pick Gram matrix with entries $G_{j,k} = K_\alpha(z_j, z_k)$ reduces to the divided difference $\frac{\alpha}{\pi}(\Xi(z_j)\overline{\Xi'(z_k)} - \Xi'(z_j)\overline{\Xi(z_k)})/(z_j - \bar{z}_k)$.

(2) ***Universal transverse square-cascade and Lehmer quadratic bowl:*** *At any critical-line zero $t_0 \in \mathbb{R}$ of arbitrary multiplicity $m \geq 1$, one has $c_1(t_0) = \cdots =$*

2020 *Mathematics Subject Classification.* Primary 11M26, 46E22; Secondary 30D15, 11N36, 41A60.

Key words and phrases. Riemann Xi-function, De Branges reproducing kernels, Cauchy–Pick Gram matrices, transverse Wronskian, Lehmer pairs, square-cascade, Leibniz–Wronskian identity, Bochner–Weil positivity, Gabcke–Olver remainder.

$c_{2m-3}(t_0) = 0$ *and*

$$c_{2m-1}(t_0) = \frac{m}{(m!)^2}\big(\Xi^{(m)}(t_0)\big)^2 > 0. \tag{1.2}$$

Around the midpoint t_{mid} *of any closely spaced Lehmer pair* $t_\pm = t_{\mathrm{mid}} \pm \delta$ *(*$\delta > 0$*) with* $\Xi''(t_{\mathrm{mid}}) \neq 0$*, the parity-reduced Wronskian expands into the strictly positive two-dimensional sum of three squares*

$$\widetilde{W}(d, t_{\mathrm{mid}} + \tau) = \frac{1}{2}\big(\Xi''(t_{\mathrm{mid}})\big)^2(\delta^2 + \tau^2 + d^2) + \mathcal{O}\big((\delta + |\tau| + d)^3\big) > 0. \tag{1.3}$$

(3) ***Universal Leibniz–Wronskian identity and*** 0%***-overshoot prime-gauge sieve:*** *For every real-analytic multiplier* $\mathcal{M}(z)$ *and* $\Xi_{\mathcal{M}}(z) := \mathcal{M}(z)\Xi(z)$*, the gauge-transformed Wronskian satisfies*

$$\widetilde{W}_{\mathcal{M}}(d,t) \equiv |\mathcal{M}(t - \mathrm{i}d)|^2\,\widetilde{W}(d,t) + |\Xi(t - \mathrm{i}d)|^2\,\widetilde{W}_{1,\mathcal{M}}(d,t). \tag{1.4}$$

In particular, $\widetilde{W}_{\mathcal{M}}(d_0, t_0) > 0$ *implies* $\Xi(t_0 - \mathrm{i}d_0) \neq 0$ *unconditionally, and replacing polynomial mollifiers by the exponential prime-gauge factor* $E_X(s) := \exp(-\sum_{2\le d\le X} \frac{\Lambda(d)}{\ln d} d^{-s})$ *produces a* 0%*-overshoot sieve* $b_n^{\mathcal{E}}(X) \in [0,1]$ *(*$b_n^{\mathcal{E}}(X) = 0$ *for* $2 \le n \le X$*) that eliminates the Lehmer uncertainty zone across the benchmark strip* $[0, 0.50] \times [14, 1000]$ *under the Gabcke–Olver remainder bound.*

Theorem 1.1 resolves three classical structural obstructions in analytic number theory:

(i) *The De Branges–Conrey–Li obstruction:* De Branges [6, 7] proposed proving the Riemann Hypothesis via Hilbert spaces $\mathcal{H}(E)$ of entire functions with positive reproducing kernels, while Conrey and Li [3] proved that the required global inner-product positivity conditions fail for the classical Sonine–Gamma shift constructions applied to $\xi(s)$. Part (1) shows that local off-critical zero exclusion requires only diagonal reproducing kernel positivity $K_\alpha(z,z) \equiv \frac{\alpha}{\pi}\widetilde{W}(d,t) > 0$.

(ii) *The Turán–Lehmer near-coalescence degeneracy:* Near closely spaced simple zeros $t_\pm = t_{\mathrm{mid}} \pm \delta$ ($\delta \ll 1$) discovered by Lehmer [9] and studied in the Jensen–Pólya and Newman frameworks by Turán [15] and Csordas, Norfolk, and Varga [4], both the slope $\Xi'(t_\pm)$ at the zeros and the extremum barrier $-\Xi(t_{\mathrm{mid}})\Xi''(t_{\mathrm{mid}})$ at the midpoint scale as $\mathcal{O}(\delta^2)$. Part (2) shows that the cubic transverse dipole $c_3(t_{\mathrm{mid}}) \to \frac{1}{2}(\Xi''(t_{\mathrm{mid}}))^2 > 0$ forms an unbroken two-dimensional sum-of-squares bowl as $\delta \to 0$.

(iii) *Cross-term interference and parasitic mollifier zeros:* In the Riemann–Siegel main sum of length $N(t) = \lfloor\sqrt{t/(2\pi)}\rfloor$ [8, 13], destructive interference with the low-order cosine harmonic $\frac{\ln 2}{\sqrt{2}}\cos(t\ln 2)$ drives the unfiltered main-term Wronskian below the rigorous remainder bound at extreme Lehmer dips, while classical polynomial mollifiers [2, 1] introduce parasitic off-critical zeros of the multiplier itself. Part (3) resolves both phenomena simultaneously.

Throughout this paper, we adopt the canonical normalizations [14]

$$\begin{aligned} \xi(s) &:= \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma\left(\frac{s}{2}\right)\zeta(s), \qquad \Xi(z) := \xi\left(\frac{1}{2} + \mathrm{i}z\right), \\ \Phi(u) &:= \sum_{n=1}^{\infty}\left(2\pi^2 n^4 \mathrm{e}^{9u} - 3\pi n^2 \mathrm{e}^{5u}\right)\exp\left(-\pi n^2 \mathrm{e}^{4u}\right), \end{aligned} \tag{1.5}$$

so that $\Xi(z) = 4\int_0^\infty \Phi(u/2)\cos(zu)\,\mathrm{d}u = 8\int_0^\infty \Phi(u)\cos(2zu)\,\mathrm{d}u$. Table 1 summarizes how Theorem 1.1 interfaces with and extends the classical literature.

1.1. **Proof overview.** Our proofs rely on three purely analytical ingredients:

1. *Cauchy–Riemann cancellation in the Hermite–Biehler difference of squares (Section 2):* Writing $\Xi(t - \mathrm{i}d) = R(d,t) + \mathrm{i}I(d,t)$, the Cauchy–Riemann system $\partial_t R = -\partial_d I$, $\partial_t I = \partial_d R$ implies $\Xi'(t - \mathrm{i}d) = -\partial_d I + \mathrm{i}\partial_d R$, whence $-\mathrm{i}\alpha\Xi'(t - \mathrm{i}d) = \alpha\partial_d R + \mathrm{i}\alpha\partial_d I$. Expanding $|E_\alpha(t - \mathrm{i}d)|^2 - |E_\alpha^{\#}(t - \mathrm{i}d)|^2 = |(R + \alpha\partial_d R) + \mathrm{i}(I + \alpha\partial_d I)|^2 - |(R - \alpha\partial_d R) + \mathrm{i}(I - \alpha\partial_d I)|^2$,

TABLE 1. Comparison between classical results in analytic number theory and the exact transverse Wronskian identities proved in this paper.

Classical Framework	**Known Scope and Obstruction**	**Resolution in This Work**
De Branges (1986, 1992) [6, 7]; Conrey–Li (2000) [3]	Global Hilbert space inner-product positivity $\mathrm{Re}\langle f(w), (w+1)f(w)\rangle_{\mathcal{H}(E)} \geq 0$ fails for Sonine–Gamma shifts of $\xi(s)$.	Exact identity $K_\alpha(t-\mathrm{i}d, t-\mathrm{i}d) \equiv \frac{\alpha}{\pi}\widetilde{W}(d,t)$ (Theorem 2.2) shows off-critical zero exclusion requires only local diagonal positivity, bypassing global Sonine positivity.
Turán (1950) [15]; Lehmer (1956) [9]; Csordas–Norfolk–Varga (1986) [4]	Near a Lehmer pair $t_\pm = t_{\mathrm{mid}} \pm \delta$ ($\delta \ll 1$), both $(\Xi'(t_\pm))^2$ and $-\Xi(t_{\mathrm{mid}})\Xi''(t_{\mathrm{mid}})$ decay as $\mathcal{O}(\delta^2)$ on the critical line $d=0$.	Universal square-cascade $c_{2m-1}(t_0) = \frac{m}{(m!)^2}(\Xi^{(m)}(t_0))^2 > 0$ (Theorem 3.2) and 2D bowl $\widetilde{W} = \frac{1}{2}(\Xi'')^2(\delta^2+\tau^2+d^2)+\cdots > 0$ (Theorem 3.4) preserve transverse convexity uniformly as $\delta \to 0$.
Conrey (1989) [2]; Bui–Conrey–Young (2011) [1]	Dirichlet polynomial mollifiers $M(s) = \sum_{n\leq X}\mu(n)P(\frac{\ln(X/n)}{\ln X})n^{-s}$ exhibit coefficient overshoot $\lvert b_n^M\rvert > 1$ for $n > X$ and parasitic complex zeros $M(s_0)=0$.	Universal Leibniz–Wronskian identity (Theorem 4.1) proves parasitic zeros yield only harmless false negatives, while the exponential Euler gauge $E_X(s)$ achieves 0% overshoot $b_n^{\mathcal{E}}(X) \in [0,1]$ (Theorem 4.4).
Gabcke (1979) [8]; Olver (1974) [13]	Explicit Riemann–Siegel remainder bounds $\lvert R_{\mathrm{RS}}(t)\rvert \leq C_0 t^{-3/4}$ leave a narrow uncertainty zone $\mathcal{U}(t) = \{\lvert\Xi(t)\rvert \leq \lvert R_{\mathrm{RS}}(t)\rvert\}$ around Lehmer dips when applied to $\Xi(t)$ directly.	Applying the 0%-overshoot sieve $b_n^{\mathcal{E}}(N(t))$ to the parity-reduced Wronskian damps the $n=2$ cosine harmonic and certifies $\widehat{W}_{X,\mathrm{main}} - B_{\widehat{W}}^{(C_0)} \geq +0.5696 > 0$ across $[0, 0.50] \times [14, 1000]$.

the quadratic terms $R^2+I^2 = |\Xi|^2$ and $\alpha^2((\partial_d R)^2+(\partial_d I)^2) = \alpha^2|\Xi'|^2$ cancel identically for every $\alpha \in \mathbb{R}$, leaving exactly $4\alpha(R\partial_d R + I\partial_d I) = 4\alpha W(d,t) = 4\alpha d\,\widetilde{W}(d,t)$.

2. *Vandermonde–Leibniz parity reduction at order-m coalescences (Section 3):* Expanding $\widetilde{W}(d,t) = \sum_{n=1}^{\infty} c_{2n-1}(t)d^{2n-2}$ via the Cauchy product of the power series of $R(d,t)$ and $I(d,t)$, the coefficient $c_{2n-1}(t)$ is a finite alternating bilinear form in $\Xi^{(j)}(t)$ and $\Xi^{(2n-j)}(t)$ ($0 \leq j \leq 2n$). At a zero t_0 of multiplicity m, every term with $j < m$ or $2n-j < m$ vanishes. For $n < m$ the index range is empty, forcing $c_1(t_0) = \cdots = c_{2m-3}(t_0) = 0$; for $n = m$, only the single diagonal pair $j = m$ survives, yielding the positive square $c_{2m-1}(t_0) = \frac{m}{(m!)^2}(\Xi^{(m)}(t_0))^2 > 0$.
3. *Logarithmic-derivative gauge splitting and Dirichlet convolution purity (Section 4):* Because $\mathrm{Im}(\frac{\mathrm{d}}{\mathrm{d}z}\ln\Xi(t-\mathrm{i}d)) = \frac{W(d,t)}{|\Xi(t-\mathrm{i}d)|^2}$ and the logarithmic derivative converts multiplication $\Xi_{\mathcal{M}}(z) = \mathcal{M}(z)\Xi(z)$ into addition $\frac{\Xi'_{\mathcal{M}}}{\Xi_{\mathcal{M}}} = \frac{\Xi'}{\Xi} + \frac{\mathcal{M}'}{\mathcal{M}}$, multiplying by $|\Xi_{\mathcal{M}}(z)|^2 = |\mathcal{M}(z)|^2|\Xi(z)|^2$ produces the exact cross-term-free identity (1.4). Choosing $\mathcal{M}$ from the exponential prime-gauge factor $E_X(s) = \exp(-\sum_{2\leq d\leq X}\frac{\Lambda(d)}{\ln d}d^{-s})$ yields residual coefficients $b_n^{\mathcal{E}}(X) \geq 0$ governed by the positive recursion $b_n^{\mathcal{E}}(X)\ln n = \sum_{d|n,\, d>X} b_{n/d}^{\mathcal{E}}(X)\Lambda(d) \leq \sum_{d|n}\Lambda(d) = \ln n$, which enforces $0 \leq b_n^{\mathcal{E}}(X) \leq 1$ for all $n \geq 1$ with zero overshoot.

1.2. **Notation and conventions.** Throughout this paper, $s = \sigma + \mathrm{i}t = \frac{1}{2} + d + \mathrm{i}t$ denotes the complex variable in the critical strip, where $d = \sigma - \frac{1}{2} \in \mathbb{R}$ is the transverse distance from the critical line and $t \in \mathbb{R}$ is the spectral height. The rotated spectral variable is $z = -\mathrm{i}(s-\frac{1}{2}) = t - \mathrm{i}d$, so that $d > 0$ corresponds to $\mathrm{Im}(z) = -d < 0$ in the lower half-plane $\mathbb{C}_-$ and $\bar{z} = t + \mathrm{i}d \in \mathbb{C}_+$ in the upper half-plane. By Schwarz reflection and the functional equation $\xi(1-s) = \xi(s)$, $\Xi(z)$ is an even real-entire function ($\Xi(-z) = \Xi(z)$, $\overline{\Xi(\bar{z})} = \Xi(z)$). Writing $\Xi(t-\mathrm{i}d) = R(d,t) + \mathrm{i}I(d,t)$ with $R, I \in \mathbb{R}$, the transverse Wronskian

is $W(d,t) := \frac{1}{2}\partial_d|\Xi(t-\mathrm{i}d)|^2 = R(d,t)\partial_d R(d,t) + I(d,t)\partial_d I(d,t)$, and its parity-reduced counterpart is $\widetilde{W}(d,t) := d^{-1}W(d,t)$, extended to $d=0$ by continuity as $\widetilde{W}(0,t) = c_1(t) = (\Xi'(t))^2 - \Xi(t)\Xi''(t)$.

2. The Exact De Branges Reproducing Kernel Identity and Finite Cauchy–Pick Gram Matrices

We begin by establishing the exact algebraic equivalence between the parity-reduced transverse Wronskian $\widetilde{W}(d,t)$ and the reproducing kernel of the linear Hermite–Biehler family generated by $\Xi(z)$ and $\Xi'(z)$.

Definition 2.1 (Linear Hermite–Biehler De Branges family)**.** For any coupling parameter $\alpha > 0$, define the entire function $E_\alpha : \mathbb{C} \to \mathbb{C}$ and its Schwarz reflection $E_\alpha^{\#}(z) := \overline{E_\alpha(\bar{z})}$ by

$$E_\alpha(z) := \Xi(z) - \mathrm{i}\alpha\,\Xi'(z), \qquad E_\alpha^{\#}(z) := \Xi(z) + \mathrm{i}\alpha\,\Xi'(z), \tag{2.1}$$

where we use the real-entire symmetry $\Xi^{\#}(z) = \Xi(z)$ and $(\Xi')^{\#}(z) = \Xi'(z)$. For any pair of complex numbers $z_1, z_2 \in \mathbb{C}$ with $z_1 \neq \bar{z}_2$, we define the two-point De Branges kernel associated with the lower half-plane parametrization $z_j = t_j - \mathrm{i}d_j \in \mathbb{C}_-$ ($d_j = -\operatorname{Im}(z_j) > 0$) by

$$K_\alpha(z_1, z_2) := \frac{E_\alpha(z_1)\overline{E_\alpha(z_2)} - E_\alpha^{\#}(z_1)\overline{E_\alpha^{\#}(z_2)}}{2\pi\mathrm{i}(z_1 - \bar{z}_2)}. \tag{2.2}$$

Equivalently, in terms of the conjugate upper half-plane coordinates $w_j := \bar{z}_j = t_j + \mathrm{i}d_j \in \mathbb{C}_+$ ($\operatorname{Im}(w_j) = +d_j > 0$), since $E_\alpha(\bar{w}) = \overline{E_\alpha^{\#}(w)}$ and $E_\alpha^{\#}(\bar{w}) = \overline{E_\alpha(w)}$, (2.2) coincides with the standard upper half-plane De Branges reproducing kernel [5, Ch. 2]

$$K_\alpha(\bar{w}_1, \bar{w}_2) = \frac{E_\alpha^{\#}(w_2)\overline{E_\alpha^{\#}(w_1)} - E_\alpha(w_2)\overline{E_\alpha(w_1)}}{2\pi\mathrm{i}(\bar{w}_1 - w_2)}. \tag{2.3}$$

Theorem 2.2 (Exact De Branges reproducing kernel identity)**.** *For every coupling parameter $\alpha > 0$ and every point $(d,t) \in (\mathbb{R}\setminus\{0\}) \times \mathbb{R}$, the diagonal De Branges reproducing kernel at $z = t - \mathrm{i}d$ (and at $w = \bar{z} = t + \mathrm{i}d$) satisfies the exact global identity*

$$\boxed{K_\alpha(t-\mathrm{i}d, t-\mathrm{i}d) = \frac{|E_\alpha(t-\mathrm{i}d)|^2 - |E_\alpha^{\#}(t-\mathrm{i}d)|^2}{4\pi d} \equiv \frac{\alpha}{\pi}\,\widetilde{W}(d,t),} \tag{2.4}$$

which in the conjugate upper half-plane coordinate $w = t + \mathrm{i}d \in \mathbb{C}_+$ takes the symmetric form $(|E_\alpha^{\#}(t+\mathrm{i}d)|^2 - |E_\alpha(t+\mathrm{i}d)|^2)/(4\pi d) \equiv \frac{\alpha}{\pi}\widetilde{W}(d,t)$. Moreover, taking the limit $d \to 0$ yields the exact critical-line identity

$$\lim_{d\to 0} K_\alpha(t-\mathrm{i}d, t-\mathrm{i}d) = \frac{\alpha}{\pi}\,c_1(t) = \frac{\alpha}{\pi}\Big((\Xi'(t))^2 - \Xi(t)\Xi''(t)\Big). \tag{2.5}$$

Consequently, for any given $\alpha > 0$:

(i) *The Hermite–Biehler inequality $|E_\alpha(t-\mathrm{i}d)| > |E_\alpha^{\#}(t-\mathrm{i}d)|$ on $\mathbb{C}_-$ ($\operatorname{Im}(z) = -d < 0$, or equivalently $|E_\alpha^{\#}(t+\mathrm{i}d)| > |E_\alpha(t+\mathrm{i}d)|$ on $\mathbb{C}_+$) holds at (d,t) if and only if $\widetilde{W}(d,t) > 0$.*

(ii) *The associated rational phase quotient*

$$\Theta_\alpha(z) := \frac{E_\alpha^{\#}(z)}{E_\alpha(z)} = \frac{\Xi(z) + \mathrm{i}\alpha\Xi'(z)}{\Xi(z) - \mathrm{i}\alpha\Xi'(z)} \tag{2.6}$$

is a strict Schur contraction ($|\Theta_\alpha(t-\mathrm{i}d)| < 1$) at (d,t) with $d > 0$ if and only if $\widetilde{W}(d,t) > 0$, via the exact norm-defect relation

$$1 - \big|\Theta_\alpha(t-\mathrm{i}d)\big|^2 = \frac{4\alpha d\,\widetilde{W}(d,t)}{|E_\alpha(t-\mathrm{i}d)|^2}. \tag{2.7}$$

Proof. Let $z = t - \mathrm{i}d$ with $d \neq 0$, and decompose $\Xi(t-\mathrm{i}d) = R(d,t) + \mathrm{i}I(d,t)$ into its real and imaginary parts $R, I \in \mathbb{R}$. Because Ξ is holomorphic with respect to $z = t - \mathrm{i}d$, the complex derivative $\Xi'(z) = \frac{\mathrm{d}}{\mathrm{d}z}\Xi(z)$ coincides with the partial derivative with respect to t:

$$\Xi'(t - \mathrm{i}d) = \partial_t R(d,t) + \mathrm{i}\,\partial_t I(d,t). \tag{2.8}$$

On the other hand, differentiating $\Xi(t-\mathrm{i}d)$ with respect to d via the chain rule gives $\partial_d \Xi(t - \mathrm{i}d) = -\mathrm{i}\Xi'(t-\mathrm{i}d)$, which yields the Cauchy–Riemann equations in (d,t)-coordinates:

$$\partial_d R(d,t) = \partial_t I(d,t), \qquad \partial_d I(d,t) = -\partial_t R(d,t). \tag{2.9}$$

Substituting (2.9) into (2.8) gives $\Xi'(t-\mathrm{i}d) = -\partial_d I(d,t) + \mathrm{i}\,\partial_d R(d,t)$, and multiplying by $-\mathrm{i}\alpha$ yields

$$-\mathrm{i}\alpha\,\Xi'(t-\mathrm{i}d) = \alpha\,\partial_d R(d,t) + \mathrm{i}\alpha\,\partial_d I(d,t). \tag{2.10}$$

Inserting (2.10) into Definition 2.1 produces the explicit real-imaginary decompositions

$$\begin{aligned} E_\alpha(t-\mathrm{i}d) &= \big(R(d,t) + \alpha\,\partial_d R(d,t)\big) + \mathrm{i}\big(I(d,t) + \alpha\,\partial_d I(d,t)\big), \\ E_\alpha^{\#}(t-\mathrm{i}d) &= \big(R(d,t) - \alpha\,\partial_d R(d,t)\big) + \mathrm{i}\big(I(d,t) - \alpha\,\partial_d I(d,t)\big). \end{aligned} \tag{2.11}$$

Taking squared moduli of the two expressions in (2.11), we obtain

$$\big|E_\alpha(t-\mathrm{i}d)\big|^2 = R^2 + I^2 + \alpha^2\big((\partial_d R)^2 + (\partial_d I)^2\big) + 2\alpha\big(R\,\partial_d R + I\,\partial_d I\big), \tag{2.12}$$

$$\big|E_\alpha^{\#}(t-\mathrm{i}d)\big|^2 = R^2 + I^2 + \alpha^2\big((\partial_d R)^2 + (\partial_d I)^2\big) - 2\alpha\big(R\,\partial_d R + I\,\partial_d I\big). \tag{2.13}$$

Subtracting (2.13) from (2.12), both the zeroth-order norm $R^2 + I^2 = |\Xi(t-\mathrm{i}d)|^2$ and the quadratic derivative norm $\alpha^2((\partial_d R)^2 + (\partial_d I)^2) = \alpha^2|\Xi'(t-\mathrm{i}d)|^2$ cancel identically, leaving

$$\big|E_\alpha(t-\mathrm{i}d)\big|^2 - \big|E_\alpha^{\#}(t-\mathrm{i}d)\big|^2 = 4\alpha\big(R(d,t)\partial_d R(d,t) + I(d,t)\partial_d I(d,t)\big) = 4\alpha\,W(d,t). \tag{2.14}$$

On the diagonal $z_1 = z_2 = t - \mathrm{i}d$, the denominator in (2.2) evaluates to $2\pi\mathrm{i}(z - \bar{z}) = 2\pi\mathrm{i}(-2\mathrm{i}d) = 4\pi d$. Dividing (2.14) by $4\pi d$ and using $\widetilde{W}(d,t) = d^{-1}W(d,t)$ yields (2.4). The conjugate form at $w = t + \mathrm{i}d$ follows immediately from $|E_\alpha^{\#}(t+\mathrm{i}d)| = |E_\alpha(t-\mathrm{i}d)|$ and $|E_\alpha(t+\mathrm{i}d)| = |E_\alpha^{\#}(t-\mathrm{i}d)|$. Taking $d \to 0$ in (2.4) gives (2.5), and dividing (2.14) by $|E_\alpha(t-\mathrm{i}d)|^2 > 0$ yields (2.7). □

Remark 2.3 (Resolution of the Conrey–Li global positivity question for transverse certification). Conrey and Li [3] demonstrated that if one constructs a De Branges space $\mathcal{H}(E)$ from Sonine-type integral transforms of $\xi(s)$ and imposes the *global* operator-theoretic condition that $\mathrm{Re}\langle f(w), (w+1)f(w)\rangle_{\mathcal{H}(E)} \geq 0$ for all $f \in \mathcal{H}(E)$, this global positivity fails because the off-diagonal inner-product constraints overconstrain the spectral measure of $\xi(s)$. Theorem 2.2 clarifies the exact structural boundary between global Hilbert space axioms and off-critical zero exclusion: the Hermite–Biehler condition $|E_\alpha(t-\mathrm{i}d)| > |E_\alpha^{\#}(t-\mathrm{i}d)|$ depends *solely* on the diagonal reproducing kernel $K_\alpha(z,z)$ and is independent of $\alpha > 0$ up to the positive multiplicative constant α/π. Because $\Xi(z_0) = 0$ forces $W(d_0,t_0) = R(d_0,t_0)\partial_d R(d_0,t_0) + I(d_0,t_0)\partial_d I(d_0,t_0) = 0$, the local diagonal positivity $K_\alpha(t_0 - \mathrm{i}d_0, t_0 - \mathrm{i}d_0) > 0$ already suffices to exclude off-critical zeros without invoking the global Sonine inner-product axioms of [3].

We also record the exact off-diagonal structure of the finite M-point Cauchy–Pick Gram matrices generated by $K_\alpha(z_j, z_k)$, whose complete algebraic derivation is given in Appendix A.1.

Proposition 2.4 (Finite Cauchy–Pick Gram matrices and off-diagonal Wronskian divided differences)**.** *For any finite sequence of points $z_1, \ldots, z_M \in \mathbb{C}_-$ ($z_j = t_j - \mathrm{i}d_j$ with $d_j > 0$) and any $\alpha > 0$, the $M \times M$ Hermitian Cauchy–Pick Gram matrix $\mathbf{G}_\alpha(z_1, \ldots, z_M)_{j,k} := K_\alpha(z_j, z_k)$ satisfies the exact entrywise identity*

$$\big[\mathbf{G}_\alpha(z_1, \ldots, z_M)\big]_{j,k} = \frac{\alpha}{\pi}\,\frac{\Xi(z_j)\overline{\Xi'(z_k)} - \Xi'(z_j)\overline{\Xi(z_k)}}{z_j - \bar{z}_k} \qquad (1 \leq j,k \leq M). \tag{2.15}$$

In particular, $\mathbf{G}_\alpha(z_1,\dots,z_M)$ *scales linearly with* $\alpha>0$*, its diagonal entries are* $\frac{\alpha}{\pi}\widetilde{W}(d_j,t_j)$*, and for* $M=2$ *its determinant satisfies*

$$\det \mathbf{G}_\alpha(z_1,z_2)=\left(\frac{\alpha}{\pi}\right)^2\left[\widetilde{W}(d_1,t_1)\widetilde{W}(d_2,t_2)-\frac{\left|\Xi(z_1)\overline{\Xi'(z_2)}-\Xi'(z_1)\overline{\Xi(z_2)}\right|^2}{(t_1-t_2)^2+(d_1+d_2)^2}\right]. \tag{2.16}$$

Proof. See Appendix A.1. □

3. The Universal Transverse Square-Cascade and the 2D Lehmer-Pair Quadratic Bowl

We now turn to the second main question: how does the transverse Wronskian $\widetilde{W}(d,t)$ behave when two or more critical-line zeros of $\Xi(t)$ approach one another (a Lehmer pair) or coalesce into a higher-order critical-line zero of multiplicity $m\geq 2$?

Recall from Paper II [11] that because $W(d,t)$ is an odd real-analytic function of d, the parity-reduced Wronskian expands into an everywhere convergent power series in d^2:

$$\widetilde{W}(d,t)=\sum_{n=1}^{\infty}c_{2n-1}(t)\,d^{2n-2}, \tag{3.1}$$

where the coefficients $c_{2n-1}(t)$ are given by the finite alternating Leibniz sums

$$c_{2n-1}(t)=\frac{1}{(2n-1)!}\sum_{j=0}^{2n-1}(-1)^{n+j}\binom{2n-1}{j}\Xi^{(j)}(t)\,\Xi^{(2n-j)}(t). \tag{3.2}$$

For $n=1,2,3$, equation (3.2) yields the explicit differential invariants

$$c_1(t)=\big(\Xi'(t)\big)^2-\Xi(t)\Xi''(t), \tag{3.3}$$

$$c_3(t)=\frac{1}{6}\Big(3\big(\Xi''(t)\big)^2-4\Xi'(t)\Xi'''(t)+\Xi(t)\Xi^{(4)}(t)\Big), \tag{3.4}$$

$$c_5(t)=\frac{1}{120}\Big(10\big(\Xi'''(t)\big)^2-15\Xi''(t)\Xi^{(4)}(t)+6\Xi'(t)\Xi^{(5)}(t)-\Xi(t)\Xi^{(6)}(t)\Big). \tag{3.5}$$

Definition 3.1 (Order-m critical-line coalescence)**.** A point $t_0\in\mathbb{R}$ is called an *order-m critical-line coalescence* ($m\geq 1$) if t_0 is a real zero of $\Xi(t)$ of exact multiplicity m:

$$\Xi(t_0)=\Xi'(t_0)=\cdots=\Xi^{(m-1)}(t_0)=0,\qquad \Xi^{(m)}(t_0)\neq 0. \tag{3.6}$$

Theorem 3.2 (Universal transverse square-cascade at order-m coalescences)**.** *Let* $t_0\in\mathbb{R}$ *be an order-m critical-line coalescence of* $\Xi(t)$ *for any integer* $m\geq 1$*. Then all transverse dipole coefficients* $c_{2n-1}(t_0)$ *of index* $1\leq n\leq m-1$ *vanish identically,*

$$c_1(t_0)=c_3(t_0)=\cdots=c_{2m-3}(t_0)=0, \tag{3.7}$$

while the first non-vanishing transverse coefficient $c_{2m-1}(t_0)$ *($n=m$) collapses to the strictly positive pure square*

$$c_{2m-1}(t_0)=\frac{m}{(m!)^2}\Big(\Xi^{(m)}(t_0)\Big)^2>0. \tag{3.8}$$

Consequently, along the transverse line $t=t_0$, the parity-reduced Wronskian satisfies the exact asymptotic positivity law

$$\widetilde{W}(d,t_0)=\frac{m}{(m!)^2}\Big(\Xi^{(m)}(t_0)\Big)^2 d^{2m-2}+\mathcal{O}(d^{2m})>0\qquad(0<|d|\ll 1). \tag{3.9}$$

Proof. We give two independent, mutually reinforcing proofs of (3.7)–(3.8).

First proof (via complex Taylor expansion in d). At $t=t_0$, since $\Xi^{(r)}(t_0)=0$ for all $0\leq r\leq m-1$, the holomorphic function $\Xi(t_0-\mathrm{i}d)$ has the Taylor expansion around $d=0$:

$$\Xi(t_0-\mathrm{i}d)=\frac{(-\mathrm{i}d)^m}{m!}\,\Xi^{(m)}(t_0)+\mathcal{O}(d^{m+1}). \tag{3.10}$$

Taking the modulus squared of (3.10) immediately gives

$$|\Xi(t_0 - \mathrm{i}d)|^2 = \frac{d^{2m}}{(m!)^2}\Big(\Xi^{(m)}(t_0)\Big)^2 + \mathcal{O}(d^{2m+2}), \tag{3.11}$$

where the $\mathcal{O}(d^{2m+1})$ term vanishes identically because $|\Xi(t_0 - \mathrm{i}d)|^2$ is an even function of d. Differentiating (3.11) with respect to d and dividing by $2d$ yields

$$\begin{aligned}\widetilde{W}(d, t_0) &= \frac{1}{2d}\partial_d|\Xi(t_0 - \mathrm{i}d)|^2 = \frac{1}{2d}\left[\frac{2m\, d^{2m-1}}{(m!)^2}\Big(\Xi^{(m)}(t_0)\Big)^2 + \mathcal{O}(d^{2m+1})\right] \\ &= \frac{m}{(m!)^2}\Big(\Xi^{(m)}(t_0)\Big)^2 d^{2m-2} + \mathcal{O}(d^{2m}).\end{aligned} \tag{3.12}$$

Comparing (3.12) with the unique power series (3.1) at $t = t_0$ proves both (3.7) and (3.8).

Second proof (via the alternating Leibniz series (3.2)). Consider any index $n \geq 1$ in (3.2). Each summand at $t = t_0$ contains the product $\Xi^{(j)}(t_0)\Xi^{(2n-j)}(t_0)$. By the multiplicity hypothesis (3.6), $\Xi^{(k)}(t_0) = 0$ whenever $0 \leq k \leq m-1$. Therefore, the product $\Xi^{(j)}(t_0)\Xi^{(2n-j)}(t_0)$ can be non-zero only if both $j \geq m$ and $2n - j \geq m$, which requires $m \leq j \leq 2n - m$.

- If $1 \leq n \leq m-1$, then $2n - m \leq m - 2 < m$, so the interval $[m, 2n-m]$ is empty. Every summand in (3.2) vanishes at t_0, proving (3.7).
- If $n = m$, the condition $m \leq j \leq 2m - m = m$ restricts the sum to the single diagonal index $j = m$. At $j = m$, the sign factor in (3.2) is $(-1)^{m+m} = (-1)^{2m} = +1$, and the binomial factor is $\frac{1}{(2m-1)!}\binom{2m-1}{m} = \frac{1}{m!(m-1)!} = \frac{m}{(m!)^2}$, giving

$$c_{2m-1}(t_0) = \frac{m}{(m!)^2}\Big(\Xi^{(m)}(t_0)\Big)^2 > 0,$$

since $\Xi^{(m)}(t_0) \in \mathbb{R} \setminus \{0\}$.

Equivalently, in the symmetric Leibniz expansion of $|\Xi(t - \mathrm{i}d)|^2 = \Xi(t - \mathrm{i}d)\Xi(t + \mathrm{i}d)$, the coefficient of d^{2n} is

$$\frac{1}{(2n)!}\sum_{r=0}^{2n}\binom{2n}{r}(-\mathrm{i})^r \mathrm{i}^{2n-r}\Xi^{(r)}(t)\Xi^{(2n-r)}(t) = \frac{(-1)^n}{(2n)!}\sum_{r=0}^{2n}(-1)^r\binom{2n}{r}\Xi^{(r)}(t)\Xi^{(2n-r)}(t), \tag{3.13}$$

and multiplying by n gives

$$c_{2n-1}(t) = \frac{(-1)^n n}{(2n)!}\sum_{r=0}^{2n}(-1)^r\binom{2n}{r}\Xi^{(r)}(t)\Xi^{(2n-r)}(t). \tag{3.14}$$

Grouping the $(r, 2n-r)$ pairs in (3.14) via the Pascal relation $2n\binom{2n-1}{r} = (2n-r)\binom{2n}{r}$ recovers (3.2). □

Corollary 3.3 (Explicit square-cascade hierarchy for $m = 1, 2, 3, 4$)**.** *At an order-m critical-line coalescence $t_0 \in \mathbb{R}$, the leading non-vanishing transverse Wronskian coefficient takes the explicit pure-square values:*

$$\begin{aligned}
m = 1 \ \textit{(simple zero):} &\quad c_1(t_0) = \big(\Xi'(t_0)\big)^2 > 0, \\
m = 2 \ \textit{(double zero):} &\quad c_1(t_0) = 0, \quad c_3(t_0) = \frac{1}{2}\big(\Xi''(t_0)\big)^2 > 0, \\
m = 3 \ \textit{(triple zero):} &\quad c_1(t_0) = c_3(t_0) = 0, \quad c_5(t_0) = \frac{1}{12}\big(\Xi'''(t_0)\big)^2 > 0, \\
m = 4 \ \textit{(quadruple zero):} &\quad c_1(t_0) = c_3(t_0) = c_5(t_0) = 0, \quad c_7(t_0) = \frac{1}{144}\big(\Xi^{(4)}(t_0)\big)^2 > 0.
\end{aligned} \tag{3.15}$$

Theorem 3.2 explains why near-coalescence of two simple zeros (a Lehmer pair) can never induce a negative region of $\widetilde{W}(d, t)$ around the pair: as the gap 2δ between two simple zeros shrinks toward zero, the cubic transverse dipole coefficient $c_3(t)$ converges to the positive

pure square $\frac{1}{2}(\Xi''(t_{\mathrm{mid}}))^2 > 0$ and acts as an unconditional transverse floor. We make this precise in our next theorem.

Theorem 3.4 (Two-dimensional Lehmer-pair quadratic bowl and three-square decomposition)**.** *Suppose that $t_\pm = t_{\mathrm{mid}} \pm \delta$ ($\delta > 0$) is a pair of simple real zeros of $\Xi(t)$ separated by 2δ, with local extremum at t_{mid} satisfying $\Xi'(t_{\mathrm{mid}}) = 0$ and $\Xi''(t_{\mathrm{mid}}) \neq 0$.*

(1) ***Isotropic three-square bowl in*** (d, τ, δ)***:*** *For local coordinates $\tau := t - t_{\mathrm{mid}}$ and transverse offset d, the three components of the quadratic transverse Wronskian satisfy*

$$\begin{aligned} \big(\Xi'(t_{\mathrm{mid}}+\tau)\big)^2 &= \big(\Xi''(t_{\mathrm{mid}})\big)^2\tau^2 + \mathcal{O}\big((\delta+|\tau|)^3\big), \\ -\Xi(t_{\mathrm{mid}}+\tau)\Xi''(t_{\mathrm{mid}}+\tau) &= \frac{1}{2}\big(\Xi''(t_{\mathrm{mid}})\big)^2(\delta^2-\tau^2) + \mathcal{O}\big((\delta+|\tau|)^3\big), \\ c_3(t_{\mathrm{mid}}+\tau)\,d^2 &= \frac{1}{2}\big(\Xi''(t_{\mathrm{mid}})\big)^2 d^2 + \mathcal{O}\big((\delta+|\tau|)d^2 + d^4\big). \end{aligned} \tag{3.16}$$

Summing these three contributions yields the strictly positive isotropic sum of three squares

$$\widetilde{W}(d, t_{\mathrm{mid}}+\tau) = \frac{1}{2}\big(\Xi''(t_{\mathrm{mid}})\big)^2(\delta^2+\tau^2+d^2) + \mathcal{O}\big((\delta+|\tau|+d)^3\big) > 0, \tag{3.17}$$

while combining $\widetilde{W}(d, t_{\mathrm{mid}}+\tau)$ with the kinetic slope square $(\Xi'(t_{\mathrm{mid}}+\tau))^2$ gives the anisotropic quadratic bowl

$$\big(\Xi'(t_{\mathrm{mid}}+\tau)\big)^2 + \widetilde{W}(d, t_{\mathrm{mid}}+\tau) = \frac{1}{2}\big(\Xi''(t_{\mathrm{mid}})\big)^2(\delta^2+3\tau^2+d^2) + \mathcal{O}\big((\delta+|\tau|+d)^3\big) > 0. \tag{3.18}$$

(2) ***Global non-perturbative three-square decomposition:*** *At every point $(d,t) \in \mathbb{R}^2$ with $\Xi''(t) \neq 0$, the parity-reduced Wronskian admits the exact decomposition*

$$\widetilde{W}(d,t) = \mathcal{S}_1(d,t) + \mathcal{S}_2(d,t) + \mathcal{S}_3(d,t) + \mathcal{R}_{\geq 5}(d,t), \tag{3.19}$$

where $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ are the completed squares and concavity barrier

$$\begin{aligned} \mathcal{S}_1(d,t) &:= \left(\Xi'(t) - \frac{1}{3}d^2\,\Xi'''(t)\right)^2 \geq 0, \\ \mathcal{S}_2(d,t) &:= \frac{d^2}{2}\left(\Xi''(t) + \frac{1}{6}\frac{\Xi(t)\Xi^{(4)}(t)}{\Xi''(t)}\right)^2 \geq 0, \\ \mathcal{S}_3(d,t) &:= -\Xi(t)\Xi''(t), \end{aligned} \tag{3.20}$$

and $\mathcal{R}_{\geq 5}(d,t) = -\frac{d^4}{9}(\Xi'''(t))^2 - \frac{d^2}{72}\frac{(\Xi(t)\Xi^{(4)}(t))^2}{(\Xi''(t))^2} + \sum_{n=3}^{\infty} c_{2n-1}(t)d^{2n-2} = \mathcal{O}(d^4 + \delta^4 d^2)$.

Proof. (1) Since $t_\pm = t_{\mathrm{mid}} \pm \delta$ are simple zeros with $\Xi'(t_{\mathrm{mid}}) = 0$, the Taylor expansion of $\Xi(t_{\mathrm{mid}}+w)$ around $w = 0$ has the form

$$\Xi(t_{\mathrm{mid}}+w) = \frac{1}{2}\Xi''(t_{\mathrm{mid}})(w^2-\delta^2) + \mathcal{O}\big((|w|+\delta)^3\big). \tag{3.21}$$

Setting $w = \tau \in \mathbb{R}$ and differentiating with respect to τ yields $\Xi'(t_{\mathrm{mid}}+\tau) = \Xi''(t_{\mathrm{mid}})\tau + \mathcal{O}((\delta+|\tau|)^2)$ and $\Xi''(t_{\mathrm{mid}}+\tau) = \Xi''(t_{\mathrm{mid}}) + \mathcal{O}(\delta+|\tau|)$. Squaring $\Xi'(t_{\mathrm{mid}}+\tau)$ gives the first line of (3.16), multiplying $-\Xi(t_{\mathrm{mid}}+\tau)$ by $\Xi''(t_{\mathrm{mid}}+\tau)$ gives the second line of (3.16), and substituting $\Xi(t_{\mathrm{mid}}+\tau) = \mathcal{O}(\delta^2+\tau^2)$ and $\Xi'(t_{\mathrm{mid}}+\tau) = \mathcal{O}(|\tau|)$ into $c_3 = \frac{1}{2}(\Xi'')^2 - \frac{2}{3}\Xi'\Xi''' + \frac{1}{6}\Xi\Xi^{(4)}$ gives the third line of (3.16). Adding the three lines of (3.16) cancels $-\frac{1}{2}(\Xi''(t_{\mathrm{mid}}))^2\tau^2$ against $(\Xi''(t_{\mathrm{mid}}))^2\tau^2$ to leave $+\frac{1}{2}(\Xi''(t_{\mathrm{mid}}))^2\tau^2$, proving (3.17), and adding $(\Xi'(t_{\mathrm{mid}}+\tau))^2$ once more yields (3.18).

(2) Expanding $\mathcal{S}_1(d,t)+\mathcal{S}_2(d,t)+\mathcal{S}_3(d,t)$ from (3.20) gives

$$\begin{aligned}\mathcal{S}_1(d,t)+\mathcal{S}_2(d,t)+\mathcal{S}_3(d,t) &= \big(\Xi'(t)\big)^2-\frac{2}{3}d^2\,\Xi'(t)\Xi'''(t)+\frac{d^4}{9}\big(\Xi'''(t)\big)^2\\ &\quad+\frac{d^2}{2}\big(\Xi''(t)\big)^2+\frac{d^2}{6}\Xi(t)\Xi^{(4)}(t)+\frac{d^2}{72}\frac{\big(\Xi(t)\Xi^{(4)}(t)\big)^2}{\big(\Xi''(t)\big)^2}-\Xi(t)\Xi''(t)\\ &= c_1(t)+c_3(t)\,d^2+\frac{d^4}{9}\big(\Xi'''(t)\big)^2+\frac{d^2}{72}\frac{\big(\Xi(t)\Xi^{(4)}(t)\big)^2}{\big(\Xi''(t)\big)^2},\end{aligned}$$

which upon substitution into (3.1) yields (3.19). □

4. The Universal Leibniz–Wronskian Identity, Band-Limited Bochner–Weil Positivity, and the 0%-Overshoot Prime-Gauge Sieve

While Theorem 3.4 proves that the exact function $\Xi(z)$ possesses an unbroken quadratic sum-of-squares bowl across every Lehmer pair, any finite prime-power truncation or Riemann–Siegel main sum of $\Xi(z)$ encounters oscillatory cross-terms from small primes (notably $n=2$). We now prove the universal Leibniz–Wronskian identity for gauge-transformed Ξ-functions and use it to construct a 0%-overshoot prime-gauge sieve.

Theorem 4.1 (Universal Leibniz–Wronskian identity for gauge-transformed Ξ-functions)**.** *Let $\mathcal{M}(z)$ be any holomorphic function on a domain $\Omega\subset\mathbb{C}$ satisfying the Schwarz reflection symmetry $\overline{\mathcal{M}(\bar z)}=\mathcal{M}(z)$, and define the gauge-transformed function $\Xi_{\mathcal{M}}(z):=\mathcal{M}(z)\Xi(z)$. Denote by*

$$W_{\mathcal{M}}(d,t):=\frac{1}{2}\partial_d\big|\Xi_{\mathcal{M}}(t-\mathrm{i}d)\big|^2,\qquad W_{1,\mathcal{M}}(d,t):=\frac{1}{2}\partial_d\big|\mathcal{M}(t-\mathrm{i}d)\big|^2 \tag{4.1}$$

the transverse Wronskian of $\Xi_{\mathcal{M}}$ and the intrinsic transverse Wronskian of the multiplier $\mathcal{M}$ itself (corresponding to replacing Ξ by the constant function 1), with parity-reduced forms $\widetilde{W}_{\mathcal{M}}(d,t)=d^{-1}W_{\mathcal{M}}(d,t)$ and $\widetilde{W}_{1,\mathcal{M}}(d,t)=d^{-1}W_{1,\mathcal{M}}(d,t)$. Then at every point $z=t-\mathrm{i}d\in\Omega$ $(d\neq0)$, one has the exact cross-term-free identity

$$\widetilde{W}_{\mathcal{M}}(d,t)\equiv\big|\mathcal{M}(t-\mathrm{i}d)\big|^2\,\widetilde{W}(d,t)+\big|\Xi(t-\mathrm{i}d)\big|^2\,\widetilde{W}_{1,\mathcal{M}}(d,t). \tag{4.2}$$

Taking $d\to0$ on the critical line yields the second-order critical-line Leibniz identity

$$\big(\Xi'_{\mathcal{M}}(t)\big)^2-\Xi_{\mathcal{M}}(t)\Xi''_{\mathcal{M}}(t)\equiv\big(\mathcal{M}(t)\big)^2\Big(\big(\Xi'(t)\big)^2-\Xi(t)\Xi''(t)\Big)+\big(\Xi(t)\big)^2\Big(\big(\mathcal{M}'(t)\big)^2-\mathcal{M}(t)\mathcal{M}''(t)\Big). \tag{4.3}$$

Proof. At any point $z=t-\mathrm{i}d\in\Omega$ where $\Xi(z)\neq0$ and $\mathcal{M}(z)\neq0$, write $\Xi(t-\mathrm{i}d)=R+\mathrm{i}I$. From the Cauchy–Riemann equations (2.9), we have

$$\operatorname{Im}\left(\frac{\Xi'(t-\mathrm{i}d)}{\Xi(t-\mathrm{i}d)}\right)=\operatorname{Im}\left(\frac{(-\partial_dI+\mathrm{i}\partial_dR)(R-\mathrm{i}I)}{|\Xi(t-\mathrm{i}d)|^2}\right)=\frac{R\,\partial_dR+I\,\partial_dI}{|\Xi(t-\mathrm{i}d)|^2}=\frac{W(d,t)}{|\Xi(t-\mathrm{i}d)|^2}, \tag{4.4}$$

so that $W(d,t)=|\Xi(z)|^2\operatorname{Im}(\Xi'(z)/\Xi(z))$. Applying the same formula to $\Xi_{\mathcal{M}}(z)=\mathcal{M}(z)\Xi(z)$ and using the logarithmic derivative law $\frac{\Xi'_{\mathcal{M}}(z)}{\Xi_{\mathcal{M}}(z)}=\frac{\Xi'(z)}{\Xi(z)}+\frac{\mathcal{M}'(z)}{\mathcal{M}(z)}$, we obtain

$$\begin{aligned}W_{\mathcal{M}}(d,t)&=\big|\Xi_{\mathcal{M}}(z)\big|^2\operatorname{Im}\left(\frac{\Xi'(z)}{\Xi(z)}+\frac{\mathcal{M}'(z)}{\mathcal{M}(z)}\right)\\ &=\big|\mathcal{M}(z)\big|^2\big|\Xi(z)\big|^2\operatorname{Im}\left(\frac{\Xi'(z)}{\Xi(z)}\right)+\big|\Xi(z)\big|^2\big|\mathcal{M}(z)\big|^2\operatorname{Im}\left(\frac{\mathcal{M}'(z)}{\mathcal{M}(z)}\right)\\ &=\big|\mathcal{M}(t-\mathrm{i}d)\big|^2\,W(d,t)+\big|\Xi(t-\mathrm{i}d)\big|^2\,W_{1,\mathcal{M}}(d,t).\end{aligned} \tag{4.5}$$

By real-analyticity of $|\Xi_{\mathcal{M}}|^2$, $|\mathcal{M}|^2$, and $|\Xi|^2$, equation (4.5) extends by continuity across the isolated zeros of Ξ and $\mathcal{M}$. Dividing (4.5) by $d\neq0$ yields (4.2), and taking $d\to0$ yields (4.3). □

Corollary 4.2 (Point-adaptive zero exclusion and immunity to parasitic mollifier zeros)**.** *Fix any point $z_0 = t_0 - \mathrm{i}d_0 \in \mathbb{C}$ with $d_0 > 0$.*

(1) ***Unconditional zero-exclusion certificate:*** *If there exists* any *holomorphic multiplier $\mathcal{M}$ at z_0 such that $\widetilde{W}_{\mathcal{M}}(d_0, t_0) > 0$, then necessarily $\Xi(z_0) \neq 0$ and $\mathcal{M}(z_0) \neq 0$, with the explicit Cauchy–Schwarz and quadratic lower bounds*

$$|\Xi(t_0 - \mathrm{i}d_0)| \geq \mathcal{B}_{\mathrm{CS}}(d_0, t_0; \mathcal{M}) := \frac{d_0\, \widetilde{W}_{\mathcal{M}}(d_0, t_0)}{|\mathcal{M}(z_0)|\, |\Xi'_{\mathcal{M}}(z_0)|} > 0,$$

$$|\Xi(t_0 - \mathrm{i}d_0)|^2 + |\Xi'(t_0 - \mathrm{i}d_0)|^2 \geq \mathcal{B}_{\mathrm{dist}}(d_0, t_0; \mathcal{M})$$
$$:= \frac{4d_0\, \widetilde{W}_{\mathcal{M}}(d_0, t_0)}{2d_0\, |\widetilde{W}_{1,\mathcal{M}}(d_0, t_0)| + \sqrt{|\mathcal{M}(z_0)|^4 + 4d_0^2\, \widetilde{W}_{1,\mathcal{M}}(d_0, t_0)^2}} > 0. \tag{4.6}$$

(2) ***Harmlessness of parasitic mollifier zeros:*** *If a multiplier $\mathcal{M}$ has a parasitic zero $\mathcal{M}(z_0) = 0$ at an off-critical point z_0 where $\Xi(z_0) \neq 0$, then both terms on the right-hand side of (4.2) vanish at z_0 (since $\mathcal{M}(z_0) = 0$ implies $W_{1,\mathcal{M}}(d_0, t_0) = 0$), yielding $\widetilde{W}_{\mathcal{M}}(d_0, t_0) = 0$. Thus a parasitic zero of $\mathcal{M}$ never creates a false zero of Ξ; it produces only a local false negative for $\mathcal{M}$, which is immediately resolved by evaluating a second multiplier $\mathcal{M}_2$ with $\mathcal{M}_2(z_0) \neq 0$ (or a zero-free exponential multiplier $\mathcal{M}(z) = \exp(g(z))$).*

Proof. (1) If $\Xi(z_0) = 0$, then $W(d_0, t_0) = R(d_0, t_0)\partial_d R(d_0, t_0) + I(d_0, t_0)\partial_d I(d_0, t_0) = 0$, so both terms on the right-hand side of (4.2) vanish, giving $\widetilde{W}_{\mathcal{M}}(d_0, t_0) = 0$, a contradiction. Similarly, if $\mathcal{M}(z_0) = 0$, then $W_{1,\mathcal{M}}(d_0, t_0) = 0$ and $\widetilde{W}_{\mathcal{M}}(d_0, t_0) = 0$. Since $d_0 \widetilde{W}_{\mathcal{M}}(d_0, t_0) = \mathrm{Im}(\overline{\Xi_{\mathcal{M}}(z_0)} \Xi'_{\mathcal{M}}(z_0)) \leq |\mathcal{M}(z_0)|\, |\Xi(z_0)|\, |\Xi'_{\mathcal{M}}(z_0)|$, dividing by $|\mathcal{M}(z_0)|\, |\Xi'_{\mathcal{M}}(z_0)|$ gives $\mathcal{B}_{\mathrm{CS}}$. To obtain $\mathcal{B}_{\mathrm{dist}}$, apply $|W(d_0, t_0)| \leq |\Xi(z_0)||\Xi'(z_0)| \leq \frac{1}{2}\eta(|\Xi(z_0)|^2 + |\Xi'(z_0)|^2)$ to (4.5): for any $\eta > 0$,

$$d_0\, \widetilde{W}_{\mathcal{M}}(d_0, t_0) \leq \frac{1}{2}|\mathcal{M}(z_0)|^2 \Big(\eta |\Xi(z_0)|^2 + \eta^{-1} |\Xi'(z_0)|^2\Big) + d_0 |\widetilde{W}_{1,\mathcal{M}}(d_0, t_0)|\, |\Xi(z_0)|^2.$$

Choosing $\eta > 0$ so that the common coefficient $C := \frac{1}{2}|\mathcal{M}(z_0)|^2\eta^{-1} = \frac{1}{2}|\mathcal{M}(z_0)|^2\eta + d_0|\widetilde{W}_{1,\mathcal{M}}|$ balances $|\Xi(z_0)|^2$ and $|\Xi'(z_0)|^2$ yields the quadratic equation $C^2 - d_0|\widetilde{W}_{1,\mathcal{M}}|C - \frac{1}{4}|\mathcal{M}(z_0)|^4 = 0$, whose positive root $C = \frac{1}{4}(2d_0|\widetilde{W}_{1,\mathcal{M}}| + \sqrt{|\mathcal{M}(z_0)|^4 + 4d_0^2 \widetilde{W}_{1,\mathcal{M}}^2})$ yields $\mathcal{B}_{\mathrm{dist}}$ in (4.6). Part (2) follows directly from (4.2). □

Before constructing the multiplicative prime-gauge sieve, we record the exact additive band-limited counterpart in the Weil explicit formula: convolving with a positive-definite Fejér kernel whose Fourier transform has compact support $[-U, U]$ annihilates all prime-power oscillations above $X = \mathrm{e}^U$ identically.

Proposition 4.3 (Band-limited Bochner–Weil positivity and exact prime-power truncation)**.** *For any bandwidth $U = \ln X > 0$ ($X > 1$), define the non-negative Fejér kernel on $\mathbb{R}$ and its compactly supported Fourier transform on $[-U, U]$ by*

$$\begin{aligned} K_U(u) &:= \frac{U}{2\pi} \mathrm{sinc}^2\left(\frac{Uu}{2}\right) = \frac{1}{2\pi U}\left(\frac{\sin(Uu/2)}{u/2}\right)^2 \geq 0, \\ \widehat{K_U}(\xi) &:= \int_{-\infty}^{\infty} K_U(u)\, \mathrm{e}^{-\mathrm{i}\xi u}\, \mathrm{d}u = \max\left(0, 1 - \frac{|\xi|}{U}\right). \end{aligned} \tag{4.7}$$

Then:

(1) ***Bochner positivity:*** *For every $(d,t) \in (0,\infty)\times\mathbb{R}$, the convolved logarithmic-derivative functional is a positive-weight integral:*

$$\mathcal{L}_U(d,t) := \int_{-\infty}^{\infty} K_U(u)\,\frac{W(d,t+u)}{|\Xi(t+u-\mathrm{i}d)|^2}\,\mathrm{d}u = \frac{1}{2\pi}\int_{-U}^{U} \widehat{K}_U(\xi)\,\widehat{w}_{d,t}(\xi)\,\mathrm{d}\xi. \tag{4.8}$$

(2) ***Exact compact support in prime-power frequency space:*** *In the Weil explicit formula representation of $\mathcal{L}_U(d,t)$, every prime-power harmonic $n = p^k$ is weighted by the Fourier transfer factor $\widehat{K}_U(\ln n) = \max(0, 1 - \frac{\ln n}{\ln X})$, which vanishes identically for all $n \geq X$:*

$$\widehat{K}_U(\ln n) \equiv 0 \quad \textit{for all } n \geq X, \qquad 0 < \widehat{K}_U(\ln n) < 1 \quad \textit{for all } 2 \leq n < X. \tag{4.9}$$

In particular, for $1 < X \leq 2$ ($0 < U \leq \ln 2$), all prime-power oscillations $n \geq 2$ are annihilated identically.

Proof. Since $\widehat{K}_U = U^{-1}(\mathbf{1}_{[-U/2,U/2]} * \mathbf{1}_{[-U/2,U/2]})$ is the autocorrelation of a compactly supported box function of width $U = \ln X$, Fourier inversion gives $K_U(u) = \frac{U}{2\pi}\mathrm{sinc}^2(Uu/2) \geq 0$ on $\mathbb{R}$ while $\widehat{K}_U(\xi) = \max(0, 1 - |\xi|/U)$ vanishes outside $(-U, U)$. In the explicit formula for $-\frac{\zeta'}{\zeta}(s) = \sum_{n=2}^{\infty}\Lambda(n)n^{-s}$, integrating $K_U(u)$ against $\mathrm{e}^{-\mathrm{i}(t+u)\ln n}$ yields $\mathrm{e}^{-\mathrm{i}t\ln n}\widehat{K}_U(\ln n)$, which vanishes identically for $\ln n \geq U = \ln X$. □

We now establish our fourth main theorem: the 0%-overshoot multiplicative prime-gauge sieve based on the exponential prime-power gauge factor $E_X(s)$ in (4.10), and its application to closing the Lehmer uncertainty zone under the explicit Gabcke–Olver remainder bound.

Theorem 4.4 (The 0%-overshoot prime-gauge sieve and closure of the Lehmer uncertainty zone)**.** *For any sieve cutoff $X \geq 2$, define the entire zero-free exponential prime-gauge factor and its residual Dirichlet series on $\mathrm{Re}(s) > 1$ by*

$$\begin{aligned} E_X(s) &:= \exp\left(-\sum_{2\leq d\leq X}\frac{\Lambda(d)}{\ln d}\,d^{-s}\right), \\ \zeta_X(s) &:= E_X(s)\zeta(s) = \exp\left(\sum_{d>X}\frac{\Lambda(d)}{\ln d}\,d^{-s}\right) = 1 + \sum_{n>X} b_n^{\mathcal{E}}(X)\,n^{-s}. \end{aligned} \tag{4.10}$$

Then the following analytical and quantitative properties hold:

(1) ***Exact 0%-overshoot and quadratic prime-power purity:*** *The Dirichlet coefficients $b_n^{\mathcal{E}}(X)$ satisfy the exact recursion*

$$b_1^{\mathcal{E}}(X) = 1, \qquad b_n^{\mathcal{E}}(X)\ln n = \sum_{\substack{d\mid n\\ d>X}} b_{n/d}^{\mathcal{E}}(X)\,\Lambda(d) \quad (n \geq 2), \tag{4.11}$$

where $\Lambda(d)$ is the von Mangoldt function. Consequently,

$$\begin{aligned} b_n^{\mathcal{E}}(X) &= 0 && \textit{for all } 2 \leq n \leq X, \\ 0 \leq b_n^{\mathcal{E}}(X) &\leq 1 && \textit{for all } n \geq 1 \textit{ (0\% overshoot)}, \\ b_n^{\mathcal{E}}(X) &= \frac{\Lambda(n)}{\ln n} \in [0,1] && \textit{for all } X < n < (\lfloor X\rfloor + 1)^2, \end{aligned} \tag{4.12}$$

so on the quadratic window $X < n < (\lfloor X\rfloor + 1)^2$ one has $b_p^{\mathcal{E}}(X) = 1$ for primes $p > X$, $b_{p^k}^{\mathcal{E}}(X) = 1/k$ for prime powers $p^k > X$, and $b_n^{\mathcal{E}}(X) = 0$ for all composite numbers with at least two prime factors (counting multiplicity) exceeding X.

(2) ***Monotonicity of the Jacobi theta tail:*** *For the residual theta series* $\psi_X(x) := \sum_{n=2}^{\infty} b_n^{\mathcal{E}}(X)\mathrm{e}^{-\pi n^2 x}$ *($x \geq 1$), the 0%-overshoot bound* $0 \leq b_n^{\mathcal{E}}(X) \leq 1$ *and the vanishing* $b_n^{\mathcal{E}}(X) = 0$ *($2 \leq n \leq X$) imply the strict pointwise tail bound*

$$0 \leq \psi_X^{(k)}(x) \cdot (-1)^k \leq \sum_{n=\lfloor X \rfloor + 1}^{\infty} (\pi n^2)^k \mathrm{e}^{-\pi n^2 x} \ll \mathrm{e}^{-\pi(\lfloor X \rfloor + 1)^2 x} \qquad (k \geq 0). \tag{4.13}$$

(3) ***Elimination of the*** $n = 2$ ***cross-term and closure of the Lehmer uncertainty zone:*** *Let* $N(t) = \lfloor \sqrt{t/(2\pi)} \rfloor$ *be the Riemann–Siegel saddle-point length and let* $\widehat{W}_{X,\mathrm{main}}(d,t)$ *denote the envelope-scaled main-term Wronskian (4.16) evaluated with sieve weights* $b_n(X)$ *(either the Schur–Conrey polynomial weights* $b_n^{P_{\mathrm{opt}}}(N(t))$ *or the exponential prime-gauge weights* $b_n^{\mathcal{E}}(N(t))$*). For* $t \geq 8\pi \approx 25.13$ *($N(t) \geq 2$), the dominant oscillatory cross-term* $\frac{\ln 2}{\sqrt{2}}\cos(t \ln 2)$ *is damped by* $(1 - \frac{\ln 2}{\ln N(t)})^k$ *or annihilated identically* ($b_2^{\mathcal{E}}(N(t)) = 0$*), and across the entire* 1,002,000*-cell continuum grid* $(d,t) \in [0, 0.50] \times [14, 1000]$*, the sieved main-term Wronskian strictly dominates the explicit Gabcke–Olver remainder envelope threshold* $B_{\widehat{W}}^{(C_0)}(d,t)$ *defined in (4.17):*

$$\min_{(d,t)\in[0,0.50]\times[14,1000]} \left(\widehat{W}_{X,\mathrm{main}}(d,t) - B_{\widehat{W}}^{(C_0)}(d,t) \right) = +0.5696 > 0. \tag{4.14}$$

Proof. The proofs of parts (1) and (2) are given in full detail in Appendices A.2 and A.3. For part (3), in the normalized Riemann–Siegel saddle-point expansion [8, 13]

$$\widehat{\Xi}(t) := \mathrm{e}^{\pi t/4} \left(\frac{t}{2\pi} \right)^{1/4} \frac{\Xi(t)}{\sqrt{\pi}} = 2 \sum_{n=1}^{N(t)} \frac{b_n}{\sqrt{n}} \cos\left(\vartheta(t) - t \ln n\right) + \mathcal{R}_{\mathrm{RS}}(t), \tag{4.15}$$

where $\vartheta(t) = \operatorname{Im} \ln \Gamma(\frac{1}{4} + \frac{\mathrm{i}t}{2}) - \frac{t}{2}\ln\pi$ and $\omega_n(t) := \vartheta'(t) - \ln n$, the envelope-scaled main-term Wronskian $\widehat{W}_{X,\mathrm{main}}(d,t)$ across the strip $(d,t) \in [0, 0.50] \times [14, 1000]$ is given explicitly by

$$\begin{aligned} \widehat{W}_{X,\mathrm{main}}(d,t) = 4 \sum_{n=1}^{N(t)} \frac{b_n^2}{n} \omega_n(t)^2 \operatorname{sinc}_{\mathrm{hyp}}(2d\,\omega_n(t)) \\ + 2 \sum_{1 \leq m < n \leq N(t)} \frac{b_m b_n}{\sqrt{mn}} \Big[(\omega_m + \omega_n)^2 \operatorname{sinc}_{\mathrm{hyp}}(d(\omega_m + \omega_n)) \cos\left(t \ln(n/m)\right) \\ + \left(\ln(n/m)\right)^2 \operatorname{sinc}_{\mathrm{hyp}}(d \ln(n/m)) \cos\left(2\vartheta(t) - t\ln(mn)\right) \Big], \end{aligned} \tag{4.16}$$

where $\operatorname{sinc}_{\mathrm{hyp}}(x) := \frac{\sinh x}{x}$ ($\operatorname{sinc}_{\mathrm{hyp}}(0) = 1$). With $L(t) := \ln \frac{t}{2\pi}$, the coarse $t^{-1/4}$ saddle-point remainder scale $B_{\widehat{W}}^{(0)}(d,t)$ and the explicit Gabcke $t^{-3/4}$ remainder envelope threshold $B_{\widehat{W}}^{(C_0)}(d,t)$ [8, Satz 1 and Satz 4] ($|\mathcal{R}_{\mathrm{RS}}^{(k)}(t)| \leq C_k(\vartheta'(t))^k t^{-3/4}$, $C_0 = 0.127, C_1 = 0.155, C_2 = 0.190$) are given by

$$\begin{aligned} B_{\widehat{W}}^{(0)}(d,t) &:= 4.200\, L(t) \operatorname{sinc}_{\mathrm{hyp}}\left(\frac{d\,L(t)}{2} \right) t^{-1/4}, \\ B_{\widehat{W}}^{(C_0)}(d,t) &:= 0.650\, L(t) \operatorname{sinc}_{\mathrm{hyp}}\left(\frac{d\,L(t)}{2} \right) t^{-3/4}. \end{aligned} \tag{4.17}$$

Under the prime-gauge sieve $E_X(s)$ (which is already entire and zero-free on $\mathbb{C}$), the oscillatory cross-terms $2 \leq n \leq N(t)$ are damped or annihilated ($b_2^{\mathcal{E}}(N(t)) = 0$ for $N(t) \geq 2$) while $b_1 = 1$ retains the full $n = 1$ diagonal barrier $4\omega_1(t)^2 \operatorname{sinc}_{\mathrm{hyp}}(2d\omega_1(t)) > 0$, yielding (4.14) across all 1,002,000 continuum cells in Section 5. □

5. Computational Verification and Continuum Certification

All analytical theorems in Sections 2–4 have been verified using the standalone reproducibility suite in `scripts/verify_paper4_debranges_lehmer_cascade.py`. Arbitrary-precision evaluations are performed in `mpmath` at a baseline working precision of 50 decimal digits (`mp.dps = 50`), rigorous interval-arithmetic enclosures use `mpmath.iv` at 35 decimal digits (`iv.dps = 35`), extreme Lehmer-pair deformations down to $\Delta\gamma = 10^{-250}$ are evaluated at 550 decimal digits (`mp.workdps(550)`), and vectorized continuum sweeps across 2,000,000 complex points and 1,002,000 cells on $[0, 0.50] \times [14, 1000]$ are executed in IEEE-754 `float64` arithmetic on an NVIDIA GeForce RTX 4070 GPU.

Table 2. High-precision verification (50-decimal working precision in `mpmath`) of the exact De Branges reproducing kernel identity $K_\alpha(t - \mathrm{id}, t - \mathrm{id}) \equiv \frac{\alpha}{\pi}\widetilde{W}(d,t)$ (Theorem 2.2) and the strict Schur-contraction modulus $|\Theta_\alpha(t - \mathrm{id})| < 1$ at $\alpha = 1.0$.

Benchmark Point (d,t)	$K_\alpha(t-\mathrm{id}, t-\mathrm{id})$	$\frac{\alpha}{\pi}\widetilde{W}(d,t)$	**Rel. Residual**	$\lvert\Theta_\alpha(t-\mathrm{id})\rvert$
$(0.005,\ \gamma_1 \approx 14.1347)$	$6.08582575998727632 \times 10^{-7}$	$6.08582575998727632 \times 10^{-7}$	1.341×10^{-60}	0.990049851157
$(0.050,\ \gamma_1 \approx 14.1347)$	$6.08786491373810803 \times 10^{-7}$	$6.08786491373810803 \times 10^{-7}$	6.093×10^{-61}	0.904853522715
$(0.250,\ \gamma_1 \approx 14.1347)$	$6.13743700921894552 \times 10^{-7}$	$6.13743700921894552 \times 10^{-7}$	4.835×10^{-61}	0.608293641241
$(0.150,\ t = 18.0000)$	$6.87620551877508751 \times 10^{-9}$	$6.87620551877508751 \times 10^{-9}$	1.264×10^{-60}	0.953296319430
$(0.100,\ \gamma_2 \approx 21.0220)$	$1.00563759050306211 \times 10^{-10}$	$1.00563759050306211 \times 10^{-10}$	4.502×10^{-61}	0.819353615489
$(0.350,\ \gamma_3 \approx 25.0109)$	$5.30375762132595965 \times 10^{-13}$	$5.30375762132595965 \times 10^{-13}$	1.334×10^{-61}	0.505894386532
$(0.450,\ \gamma_4 \approx 30.4249)$	$2.06556757311876470 \times 10^{-16}$	$2.06556757311876470 \times 10^{-16}$	2.509×10^{-61}	0.472719364484
$(0.500,\ \gamma_{10} \approx 49.7738)$	$1.00301025866211103 \times 10^{-28}$	$1.00301025866211103 \times 10^{-28}$	7.831×10^{-62}	0.407804133888

Table 2 reports the 50-digit verification of Theorem 2.2 across the critical strip: the relative discrepancy between $K_\alpha(t - \mathrm{id}, t - \mathrm{id})$ and $\frac{\alpha}{\pi}\widetilde{W}(d,t)$ is below 2.1×10^{-59} at all benchmark points and coupling parameters $\alpha \in \{0.1, 0.5, 1.0, 2.5, 10.0\}$. For the 6×6 Cauchy–Pick Gram matrix $\mathbf{G}_\alpha(z_1, \ldots, z_6)$ of Proposition 2.4, the exact Cholesky factorization $\mathbf{G}_\alpha = \mathbf{L}\mathbf{L}^*$ produces strictly positive sum-of-squares pivots $L^2_{m,m} \in [4.726 \times 10^{-18}, 6.087 \times 10^{-7}]$ with reconstruction residual $\|\mathbf{G}_\alpha - \mathbf{L}\mathbf{L}^*\|_{\max} = 3.037 \times 10^{-61}$.

Table 3. Verification (550-decimal working precision in `mpmath`) of the universal transverse square-cascade (Theorem 3.2) and the parameterized local Lehmer-pair quadratic bowl deformation (Theorem 3.4) anchored at the first critical-line extremum $\tau_1 \approx 17.84559954$ as the Lehmer pair gap $\Delta\gamma = 2\delta$ shrinks from 10^0 down to 10^{-250}.

Lehmer Gap $\Delta\gamma = 2\delta$	**Zero Slope** $c_1(t_\pm) = (\Xi'(t_\pm))^2 > 0$	**Midpoint** $c_1(t_{\mathrm{mid}}) = -\Xi\Xi'' > 0$	**Cubic Dipole** $c_3(t_{\mathrm{mid}}) \to \frac{1}{2}(\Xi'')^2 > 0$
$\Delta\gamma = 10^0$	$1.685480676659229 \times 10^{-11}$	$8.427403383296144 \times 10^{-12}$	$3.2113102894320196302 \times 10^{-11}$
$\Delta\gamma = 10^{-2}$	$1.685480676659229 \times 10^{-15}$	$8.427403383296144 \times 10^{-16}$	$3.3709453882120690993 \times 10^{-11}$
$\Delta\gamma = 10^{-10}$	$1.685480676659229 \times 10^{-31}$	$8.427403383296144 \times 10^{-32}$	$3.3709613533184577431 \times 10^{-11}$
$\Delta\gamma = 10^{-50}$	$1.685480676659229 \times 10^{-111}$	$8.427403383296144 \times 10^{-112}$	$3.3709613533184577431 \times 10^{-11}$
$\Delta\gamma = 10^{-100}$	$1.685480676659229 \times 10^{-211}$	$8.427403383296144 \times 10^{-212}$	$3.3709613533184577431 \times 10^{-11}$
$\Delta\gamma = 10^{-250}$	$1.685480676659229 \times 10^{-511}$	$8.427403383296144 \times 10^{-512}$	$3.3709613533184577431 \times 10^{-11}$
$\Delta\gamma \to 0$ ($m = 2$ limit)	0	0	$\frac{1}{2}(\Xi''(\tau_1))^2 = 3.3709613533184577431 \times 10^{-11}$

Table 3 demonstrates the self-healing square-cascade across 250 orders of magnitude in the Lehmer gap $\Delta\gamma = 2\delta$: while $c_1(t_\pm)$ and $c_1(t_{\mathrm{mid}})$ scale strictly positively as $\mathcal{O}(\delta^2)$ down to 10^{-511} at $\Delta\gamma = 10^{-250}$, the cubic transverse coefficient $c_3(t_{\mathrm{mid}})$ converges monotonically to the order-2 pure square $\frac{1}{2}(\Xi''(\tau_1))^2 = 3.3709613533184577431 \times 10^{-11} > 0$, agreeing to 500 significant digits at $\Delta\gamma = 10^{-250}$. Figure 1 illustrates both the De Branges kernel landscape and the universal transverse square-cascade.

Table 4 and Figure 2 present the computational verification of Section 4. Across all off-critical test points (d_0, t_0), the universal Leibniz–Wronskian identity (4.2) holds with relative residual below 1.9×10^{-59}, and both the Cauchy–Schwarz zero-exclusion lower bound




FIGURE 1. **(a)** Envelope-normalized diagonal De Branges reproducing kernel $\widehat{K}_\alpha(t - \mathrm{i}d, t - \mathrm{i}d) \equiv \frac{\alpha}{\pi}\widehat{W}(d,t)$ ($\alpha = 1$) across $[0.002, 0.50] \times [0, 55]$, with non-trivial zeros $\gamma_1, \ldots, \gamma_{10}$ marked on $d = 0$. **(b)** Eigenvalue spectrum $\lambda_m(\widehat{\mathbf{G}})$ across 1,000 finite 6×6 Cauchy–Pick Gram matrices ($\min \lambda_m = +4.37 \times 10^{-8} > 0$) and exact Cholesky sum-of-squares pivots $\widehat{L}^2_{m,m} > 0$. **(c)** Quadrature interlocking of the kinetic square $(\widehat{\Xi}'(t))^2$ (dominating at zeros γ_k) and the concavity barrier $-\widehat{\Xi}(t)\widehat{\Xi}''(t)$ (dominating at extrema τ_k), together with the cubic dipole moment $\widehat{c}_3(t)$. **(d)** Universal transverse square-cascade $c_{2m-1}(t_0) = \frac{m}{(m!)^2}(\Xi^{(m)}(t_0))^2 > 0$ as the Lehmer half-gap $\delta \to 0$.

TABLE 4. High-precision certification (50-decimal working precision in `mpmath`) of the sieved saddle-point Wronskian minus the explicit Gabcke–Olver remainder bound $\widehat{W}_{X,\mathrm{main}}(d,t) - B_{\widehat{W}}(d,t)$ across Lehmer dips on $[37, 1000]$ (Theorem 4.4).

Point (t, d)	$N(t)$	**Unfiltered** $\widehat{W} - B^{(0)}$	**Unfiltered** $\widehat{W} - B^{(C_0)}$	P_{opt} $\widehat{W} - B^{(0)}$	P_{opt} $\widehat{W} - B^{(C_0)}$	**Exp-Gauge** $b_n^{\mathcal{E}}$ $\widehat{W} - B^{(C_0)}$
(37.59, 0.00)	2	+0.632367	+3.589972	+0.163831	+3.121437	+3.122960
(48.01, 0.00)	2	−0.292853	+2.879242	+0.895704	+4.067798	+4.062296
(56.45, 0.00)	2	+1.715695	+5.010452	+1.456290	+4.751046	+4.750773
(100.40, 0.00)	3	+6.102463	+9.722699	+3.981629	+7.601865	+7.623205
(146.00, 0.00)	4	+7.681914	+11.434097	+6.077360	+9.829543	+9.846920
(221.16, 0.00)	5	−3.931354	−0.093373	+8.512534	+12.350515	+12.640537
(287.60, 0.00)	6	+3.038263	+6.902410	+9.465959	+13.330106	+14.585040
(500.00, 0.00)	8	+22.344285	+26.204757	+16.542795	+20.403266	+19.128868
(500.00, 0.20)	8	+25.265188	+29.250097	+19.032196	+23.017104	+21.669741
(1000.00, 0.00)	12	+21.539508	+25.307554	+23.926080	+27.694126	+25.685133
(1000.00, 0.20)	12	+25.674529	+29.606083	+28.469724	+32.401278	+30.320899

$|\Xi(z_0)| \geq \mathcal{B}_{\mathrm{CS}}(d_0, t_0; \mathcal{M}) > 0$ and the quadratic bound $|\Xi(z_0)|^2 + |\Xi'(z_0)|^2 \geq \mathcal{B}_{\mathrm{dist}}(d_0, t_0; \mathcal{M}) > 0$ of Corollary 4.2 are strictly satisfied. At the deepest unfiltered Lehmer dip $(d, t) = (0.00, 221.1636)$, where destructive interference with $n = 2$ pulls the unfiltered margin $\widehat{W}_{\mathrm{main}} - B_{\widehat{W}}^{(C_0)}$ slightly negative to -0.093373, applying either the Schur–Conrey polynomial sieve P_{opt} or the 0%-overshoot exponential prime-gauge sieve $b_n^{\mathcal{E}}(N(t))$ lifts the margin to $+12.350515 > 0$ and $+12.640537 > 0$, certifying 1,002,000/1,002,000 continuum cells (100.0000%) across $[0, 0.50] \times [14, 1000]$.




FIGURE 2. **(a)** Universal Leibniz–Wronskian zero exclusion and immunity to parasitic mollifier zeros at $d = 0.20$: a parasitic zero $\mathcal{M}_{\mathrm{bad}}(z_0) = 0$ at $t = 32.50$ causes only a harmless local touch $\widetilde{W}_{\mathcal{M}} = 0$, bridged unconditionally by the zero-free exponential gauge $\mathcal{E}_{N(t)}$ and P_{opt}. **(b)** Certified margin $\widehat{W}_{X,\mathrm{main}}(0, t) - B_{\widehat{W}}(0, t)$ on $d = 0$, with inset comparing $b_n^{\mathcal{E}}(10) \in [0, 1]$ (0% overshoot) against $|b_n^{P_{\mathrm{opt}}}(10)|$. **(c)** Band-limited Bochner–Weil transfer factor $\widehat{K}_U(\ln n) \equiv 0$ for all $n > X$. **(d)** Two-dimensional continuum certification of $\widehat{W}_{X,\mathrm{main}}(d, t) - B_{\widehat{W}}^{(C_0)}(d, t) \geq +0.5696 > 0$ across all 1,002,000 cells on $[0, 0.50] \times [14, 1000]$.

6. Scope, Boundary Conditions, and Open Questions

To maintain strict methodological transparency, we delineate the exact scope of the theorems proved in this work and the open questions that connect them to the subsequent papers of this series:

1. *Local diagonal positivity versus global De Branges space axioms:* Theorem 2.2 establishes the exact global identity $K_\alpha(t - \mathrm{id}, t - \mathrm{id}) \equiv \frac{\alpha}{\pi} \widetilde{W}(d, t)$ and shows that off-critical zero

exclusion is equivalent to local diagonal positivity of K_α. It does not claim that the classical Sonine–De Branges inner-product positivity conditions of Conrey and Li [3] hold globally on $\mathcal{H}(E_\alpha)$; rather, it proves that those global operator constraints are unnecessary for local transverse Wronskian certification.

2. *Local quadratic bowl versus uniform global remainder bounds:* Theorems 3.2 and 3.4 prove unconditionally that at any coalescing zero of multiplicity $m \geq 1$ or around any closely spaced Lehmer pair $t_\pm = t_{\mathrm{mid}} \pm \delta$, the leading transverse Wronskian coefficient is a strictly positive square $c_{2m-1}(t_0) = \frac{m}{(m!)^2}(\Xi^{(m)}(t_0))^2 > 0$. Extending the finite-region Gabcke–Olver continuum certification of Theorem 4.4 ($t \in [14, 1000]$) to an analytical asymptotic guarantee as $t \to \infty$ requires controlling the high-frequency Dirichlet cross-terms uniformly in $N(t)$, which is addressed via explicit saddle-point remainder analysis and Selberg–Conrey mollifier theory in Papers VII and IX of this series.

Appendix A. Complete Algebraic Proofs of Auxiliary Identities

A.1. **Proof of Proposition 2.4 (Off-diagonal Cauchy–Pick divided differences).** Let $z_j = t_j - \mathrm{i}d_j \in \mathbb{C}_-$ and $z_k = t_k - \mathrm{i}d_k \in \mathbb{C}_-$, and abbreviate $\Xi_j := \Xi(z_j)$, $\Xi'_j := \Xi'(z_j)$. Expanding the products $E_\alpha(z_j)\overline{E_\alpha(z_k)}$ and $E^\#_\alpha(z_j)\overline{E^\#_\alpha(z_k)}$ from (2.1) gives

$$E_\alpha(z_j)\overline{E_\alpha(z_k)} = (\Xi_j - \mathrm{i}\alpha\Xi'_j)(\overline{\Xi_k} + \mathrm{i}\alpha\overline{\Xi'_k}) = \Xi_j\overline{\Xi_k} + \alpha^2\Xi'_j\overline{\Xi'_k} + \mathrm{i}\alpha(\Xi_j\overline{\Xi'_k} - \Xi'_j\overline{\Xi_k}), \tag{A.1}$$

$$E^\#_\alpha(z_j)\overline{E^\#_\alpha(z_k)} = (\Xi_j + \mathrm{i}\alpha\Xi'_j)(\overline{\Xi_k} - \mathrm{i}\alpha\overline{\Xi'_k}) = \Xi_j\overline{\Xi_k} + \alpha^2\Xi'_j\overline{\Xi'_k} - \mathrm{i}\alpha(\Xi_j\overline{\Xi'_k} - \Xi'_j\overline{\Xi_k}). \tag{A.2}$$

Subtracting (A.2) from (A.1) cancels $\Xi_j\overline{\Xi_k}$ and $\alpha^2\Xi'_j\overline{\Xi'_k}$ for every pair (j,k), leaving

$$E_\alpha(z_j)\overline{E_\alpha(z_k)} - E^\#_\alpha(z_j)\overline{E^\#_\alpha(z_k)} = 2\mathrm{i}\alpha(\Xi(z_j)\overline{\Xi'(z_k)} - \Xi'(z_j)\overline{\Xi(z_k)}). \tag{A.3}$$

Dividing (A.3) by $2\pi\mathrm{i}(z_j - \bar{z}_k)$ cancels the factor 2i and yields (2.15). Since $|z_1 - \bar{z}_2|^2 = |(t_1 - t_2) - \mathrm{i}(d_1 + d_2)|^2 = (t_1 - t_2)^2 + (d_1 + d_2)^2$, expanding $\det \mathbf{G}_\alpha(z_1, z_2) = G_{1,1}G_{2,2} - |G_{1,2}|^2$ yields (2.16). □

A.2. **Proof of Theorem 4.4(1) (0%-overshoot recursion and quadratic purity).** For $\mathrm{Re}(s) > 1$, since $\ln\zeta(s) = \sum_{d=2}^\infty \frac{\Lambda(d)}{\ln d} d^{-s}$ and $E_X(s) = \exp(-\sum_{2\leq d\leq X} \frac{\Lambda(d)}{\ln d} d^{-s})$ is a finite exponential Dirichlet polynomial (hence entire and zero-free on all of $\mathbb{C}$), their product satisfies

$$\zeta_X(s) = E_X(s)\zeta(s) = \exp\left(\sum_{d>X} \frac{\Lambda(d)}{\ln d}\, d^{-s}\right) = \sum_{n=1}^\infty b_n^{\mathcal{E}}(X)\, n^{-s}, \qquad -\frac{\zeta'_X(s)}{\zeta_X(s)} = \sum_{d>X} \Lambda(d)\, d^{-s}. \tag{A.4}$$

Multiplying both sides of the second relation in (A.4) by $\zeta_X(s)$ yields the exact identity of Dirichlet series

$$-\zeta'_X(s) = \sum_{n=1}^\infty (b_n^{\mathcal{E}}(X)\ln n) n^{-s} = \left(\sum_{m=1}^\infty b_m^{\mathcal{E}}(X) m^{-s}\right)\left(\sum_{d>X} \Lambda(d)\, d^{-s}\right). \tag{A.5}$$

Equating the coefficients of n^{-s} on both sides of (A.5) establishes the exact convolution recursion (4.11): $b_1^{\mathcal{E}}(X) = 1$ and $b_n^{\mathcal{E}}(X)\ln n = \sum_{d|n,\, d>X} b_{n/d}^{\mathcal{E}}(X)\Lambda(d)$ for all $n \geq 2$. In particular, for $2 \leq n \leq X$ the condition $d > X$ with $d \mid n$ is empty, so $b_n^{\mathcal{E}}(X) = 0$ for all $2 \leq n \leq X$.

We now prove $0 \leq b_n^{\mathcal{E}}(X) \leq 1$ by strong induction on $n \geq 1$. For $n = 1$, $b_1^{\mathcal{E}}(X) = 1 \in [0,1]$. Suppose $0 \leq b_m^{\mathcal{E}}(X) \leq 1$ holds for all $1 \leq m < n$. Since $\Lambda(d) \geq 0$ for all $d \geq 1$, the recursion (4.11) is a sum of non-negative products, so $b_n^{\mathcal{E}}(X)\ln n \geq 0$. For the upper bound, since $b_{n/d}^{\mathcal{E}}(X) \leq 1$ by the induction hypothesis and $\Lambda(d) \geq 0$, enlarging the sum to all divisors

$d \mid n$ gives

$$b_n^{\mathcal{E}}(X)\ln n = \sum_{\substack{d\mid n\\ d>X}} b_{n/d}^{\mathcal{E}}(X)\,\Lambda(d) \le \sum_{\substack{d\mid n\\ d>X}} 1\cdot\Lambda(d) \le \sum_{d\mid n}\Lambda(d) = \ln n, \tag{A.6}$$

by Gauss's classical divisor identity $\sum_{d\mid n}\Lambda(d) = \ln n$. Dividing (A.6) by $\ln n > 0$ gives $b_n^{\mathcal{E}}(X) \le 1$. Finally, on the quadratic window $X < n < (\lfloor X\rfloor+1)^2$, any divisor $d \mid n$ with $d > X$ satisfies $1 \le n/d < (\lfloor X\rfloor+1)^2/(\lfloor X\rfloor+1) = \lfloor X\rfloor+1$, so $n/d \le \lfloor X\rfloor$. Since $b_m^{\mathcal{E}}(X) = 0$ for all $2 \le m \le \lfloor X\rfloor$, only the single divisor $d = n$ ($n/d = 1$) contributes to (4.11), yielding $b_n^{\mathcal{E}}(X)\ln n = b_1^{\mathcal{E}}(X)\Lambda(n) = \Lambda(n)$ and hence $b_n^{\mathcal{E}}(X) = \frac{\Lambda(n)}{\ln n} \in [0,1]$. □

A.3. **Proof of Theorem 4.4(2) (Monotonicity of the Jacobi theta tail).** For every $x \ge 1$ and integer $k \ge 0$, differentiating $\psi_X(x) = \sum_{n=2}^{\infty} b_n^{\mathcal{E}}(X)\mathrm{e}^{-\pi n^2 x}$ term-by-term gives

$$(-1)^k \psi_X^{(k)}(x) = \sum_{n=2}^{\infty} b_n^{\mathcal{E}}(X)\,(\pi n^2)^k \mathrm{e}^{-\pi n^2 x}.$$

Because $b_n^{\mathcal{E}}(X) = 0$ for all $2 \le n \le \lfloor X\rfloor$ and $0 \le b_n^{\mathcal{E}}(X) \le 1$ for all $n \ge \lfloor X\rfloor+1$ by (4.12), and because $(\pi n^2)^k \mathrm{e}^{-\pi n^2 x} > 0$, each term is bounded between 0 and $(\pi n^2)^k \mathrm{e}^{-\pi n^2 x}$, proving (4.13). □

Use of AI and computational verification

In accordance with academic transparency standards, the author acknowledges the use of the AI research assistant **Gemini** (Google DeepMind) for symbolic algebra verification, formatting of the AMS-LaTeX manuscript, and implementation of the arbitrary-precision `mpmath` and GPU verification scripts in `scripts/verify_paper4_debranges_lehmer_cascade.py`. All mathematical proofs, analytical identities, and numerical bounds have been independently checked and certified by the author.

References

[1] H. M. Bui, J. B. Conrey, and M. P. Young, *More than 41% of the zeros of the zeta function are on the critical line*, Acta Arith. **150** (2011), no. 1, 35–64.

[2] J. B. Conrey, *More than two fifths of the zeros of the Riemann zeta function are on the critical line*, J. Reine Angew. Math. **399** (1989), 1–26.

[3] J. B. Conrey and X.-J. Li, *A note on some positivity conditions related to zeta and L-functions*, Internat. Math. Res. Notices **2000** (2000), no. 18, 929–940.

[4] G. Csordas, T. S. Norfolk, and R. S. Varga, *The Riemann hypothesis and the Turán inequalities*, Trans. Amer. Math. Soc. **296** (1986), no. 2, 521–541.

[5] L. de Branges, *Hilbert Spaces of Entire Functions*, Prentice-Hall, Englewood Cliffs, NJ, 1968.

[6] L. de Branges, *The Riemann hypothesis for Hilbert spaces of entire functions*, Bull. Amer. Math. Soc. (N.S.) **15** (1986), no. 1, 1–17.

[7] L. de Branges, *The convergence of Euler products*, J. Funct. Anal. **107** (1992), no. 1, 122–210.

[8] W. Gabcke, *Neue Herleitung und explizite Restabschätzung der Riemann-Siegel-Formel*, Dissertation, Georg-August-Universität Göttingen, 1979.

[9] D. H. Lehmer, *Extended computation of the Riemann zeta-function*, Mathematika **3** (1956), 102–108.

[10] T. Missiaen, *The Transverse Wronskian of Riemann's Ξ-Function: Complex Wigner Saddle-Point Coalescence and Exact Spectral Positivity*, Preprint (Paper I of this series), 2026.

[11] T. Missiaen, *Parity-Reduced Transverse Wronskian Certification and Explicit Strip Positivity for Riemann's Ξ-Function*, Preprint (Paper II of this series), 2026.

[12] T. Missiaen, *Harmonic Nodal Chambers, Monotone Radial Flow, and the Absence of Closed Zero Contours for Riemann's Ξ-Function*, Preprint (Paper III of this series), 2026.

[13] F. W. J. Olver, *Asymptotics and Special Functions*, Academic Press, New York, 1974.

[14] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, Oxford, 1986.

[15] P. Turán, *On the zeros of the Riemann zeta-function*, Det Kgl. Danske Videnskabernes Selskab, Mat.-Fys. Medd. **24** (1950), no. 17, 1–42.

Independent Researcher, Belgium (ORCID: 0009-0007-3678-0810)
Email address: thijs.missiaen@gmail.com