Exact Rational Integration over Constrained Simplices

Symmetric Polynomial Bases, Marginal Geometry, and Lean 4 Verification

Reading Point (translatepi.ai)

Draft v0.7 — 22 October 2026

Abstract

We present formally verified integration identities for polynomial weights on simplexes constrained by coordinate thresholds and cardinality-dependent caps. The development establishes exact simplex moments, a rational polynomial subspace of dimension 846, general monomial and ordinary Gram-entry formulas, fibre decompositions, two-aggregate Dirichlet integration, and exact polygon-cell integrals covering all marginal endpoint branches. Kernel-checked rational certificates first establish the marginal entries $J_T[1,1]$, $J_T[P_2,1]$, and $J_T[P_2,P_2]$, completing a 2×2 block. The V2i development then proves a general exact rational representation for every pair of polynomial generators, with further Table 3 certificates and an exact bridge back to the V2h entry. Its 33-statement Comparator suite was accepted by the Lean default kernel on the VPS. The full regression produced 21 of 21 acceptances and exited with status zero. This establishes general representations, not an evaluated 846×846 numerical matrix or the final variational inequality.

1 Introduction and scope

Exact integration on constrained regions is useful where a variational inequality must ultimately be checked using rational arithmetic. A conventional simplex has a closed monomial moment formula; the present support also depends on which coordinates cross a threshold and on a cap for the sum of those coordinates. This introduces a finite stratification, but preserves a route to exact, rational expressions.

We study the mathematical constructions in the public Lean repository:

`github.com/thinkthoughts/lean-bgp212`

The motivating bounded-prime-gap argument originates with Axiom Math [1]; the statements below are an independent formalization of specified components and do not amount to a complete formal proof of the unconditional prime-gap bound. References to Lean declarations are included to delimit precisely what has been checked. The short arguments given here are *proof outlines*, not substitutes for the complete Lean proof terms.

2 Polynomial moments on a simplex

Definition 2.1. *For $m \in \mathbb{N}$ and $R \in \mathbb{R}$, define*

$$\Delta_m(R) = \left\{ x \in \mathbb{R}^m : x_i \geq 0 \text{ for all } i, \quad \sum_{i=1}^{m} x_i \leq R \right\}.$$

Theorem 2.2 (Simplex monomial moments)**.** *For $m \in \mathbb{N}$, $R \geq 0$, and $e \in \mathbb{N}^m$,*

$$\int_{\Delta_m(R)} \prod_{i=1}^{m} x_i^{e_i}\, dx = \frac{R^{m+\sum_i e_i} \prod_i (e_i!)}{(m+\sum_i e_i)!}.$$

Here the dimension-zero case uses the corresponding Lean integral convention.

Lean: `simplex_integral`; definitions `simplex`, `momentR`.
Source: `Solution/SimplexMoment.lean:160`.

Proof outline. Induct on dimension and use Fubini to isolate a coordinate. The inner integral is a Beta integral; the factorial expression then simplifies by induction. The formal development additionally treats negative and zero radii, simplex volume, and rational-polynomial integrals (`poly_integral`).

3 An indexed rational polynomial space

Consider rational polynomials in 45 coordinates. Write P_j for the jth power sum, P_λ for the product over parts of a multiset λ, and set

$$G_{a,\lambda} = (L - P_1)^a P_\lambda, \qquad \lambda \text{ has positive even parts}, \quad a + |\lambda| \le 21.$$

The index type $B_{21}\mathrm{Idx}$ enumerates the admissible pairs.

Theorem 3.1 (Independence and rank)**.** *The indexed family $G_{a,\lambda}$ is linearly independent over $\mathbb{Q}$. Its rational span B_{21} has*

$$\dim_{\mathbb{Q}} B_{21} = 846.$$

Lean: `genIdx_linearIndependent`, `finrank_B21` (`Solution/Variational.lean:398,410`).

Proof outline. The indexing is injective; distinct indexed monomials are linearly independent. Injective polynomial transformations preserve this independence. A separate count establishes $|B_{21}\mathrm{Idx}| = 846$, and the span-rank identity gives the result. The theorem concerns this designated subspace, not the full polynomial ring.

4 Cap-constrained supports and general integrals

Let $N \in \mathbb{N}$, $g \ge 0$, $L \in \mathbb{R}$, and $C : \mathbb{N} \to \mathbb{R}$. For $x \in \mathbb{R}^N$ write

$$R_g(x) = \{i : x_i > g\}, \qquad s_g(x) = |R_g(x)|, \qquad \rho_g(x) = \sum_{i \in R_g(x)} x_i.$$

Define the capped simplex support

$$T(N, g, L, C) = \left\{ x \in \mathbb{R}^N : x_i \ge 0, \quad \sum_i x_i \le L, \quad \rho_g(x) \le C(s_g(x)) \right\}. \tag{1}$$

Theorem 4.1 (General monomial integral)**.** *For $N \in \mathbb{N}$, $g \ge 0$, $L \in \mathbb{R}$, $C : \mathbb{N} \to \mathbb{R}$, $e \in \mathbb{N}^N$, and $p \in \mathbb{N}$,*

$$\int_{T(N,g,L,C)} \left(L - \sum_i x_i \right)^p \prod_i x_i^{e_i} \, dx = \mathrm{gramGen}(N, g, L, C, e, p). \tag{2}$$

The right-hand side is the explicit, finite expression defined in Lean.

Lean: `integral_supportT_gen`; `gramGen`.
Sources: `Solution/GramGeneral.lean:344`;
`BGP212/GramGeneral.lean`.

Proof outline. Stratify by the rough-coordinate set, apply inclusion–exclusion to threshold conditions, and translate the resulting pieces to weighted simplexes. Use Theorem 2.2 and Fubini to evaluate the polynomial weights. Placement-counting identities `sig_insert`, `sig_union`, and `sig_card` reduce the sums to the finite closed form. The reduction `gramGen_extend` transports calculation to the coordinates having nonzero exponents. This is an algebraic efficiency result, not a claim that all entries are computationally cheap to certify.

5 Rational Gram representations

Specialize to the fixed rational cap values and support parameters of Table 3 in [1], with $N = 45$. For indices $i, j \in B_{21}\mathrm{Idx}$ define

$$I_T(i,j) = \int_T G_i(x) G_j(x)\, dx.$$

Theorem 5.1 (Explicit rational representation)**.** *For every $i, j \in B_{21}\mathrm{Idx}$, $I_T(i,j)$ equals the real embedding of a finite rational sum indexed by placements φ of the combined power-sum parts:*

$$I_T(i,j) = \left(\sum_{\varphi} \mathrm{gramGen}_{\mathbb{Q}}(45, g, L, C, e_\varphi, a_i + a_j) \right)_{\mathbb{R}}.$$

Here e_φ is the exponent vector obtained from the placement, exactly as specified by ***`expo`*** *in the Lean source.*

Lean: `gram_entry_B21_rat` (`Solution/GramGeneral.lean:522`); `gram_entry_gen_list` and `gram_entry_example`.

The theorem gives each entry an explicit rational representation; it does not supply an evaluated and certified 846×846 matrix. One worked entry, corresponding to $(L - P_1)P_2^2$, has its value certified using a collection of small exact rational lemmas. This separation between an algebraic identity and its numerical certificate is important to reproducibility.

6 Marginal-fibre geometry

Let $\widehat{x} \in \mathbb{R}^n$ and append $u \in \mathbb{R}$ as the last coordinate. Define $S = \sum_i \widehat{x}_i$, $s = s_g(\widehat{x})$, $\rho_1 = \rho_g(\widehat{x})$, and

$$\widehat{T} = \{\widehat{x} \geq 0 : S \leq \tau,\ \rho_1 \leq C(s)\}, \quad U(\widehat{x}) = \min\{L - S, \max(g, C(s+1) - \rho_1)\}.$$

Theorem 6.1 (Fibre equivalence)**.** *Suppose $g \geq 0$, $\tau \leq L$, and $\widehat{x} \in \widehat{T}$. Then for every $u \in \mathbb{R}$,*

$$(\widehat{x}, u) \in T(n+1, g, L, C) \quad \Longleftrightarrow \quad 0 \leq u \leq U(\widehat{x}).$$

Lean: `fibre_equiv` (`Solution/Fibre.lean:115`).

Proof outline. Appending u leaves the rough set unchanged for $u \leq g$, and increments both rough count and rough mass for $u > g$. Combining the resulting cap inequality with $S + u \leq L$ identifies the minimum–maximum endpoint exactly. The empty-fibre theorem for cap-violating outer coordinates additionally needs $C(m+1) \leq C(m) + g$ for $m \geq 1$, and $C(0) \geq 0$. No constraint on the step $m = 0$ is assumed; the relevant Table 3 facts are checked in Lean.

Theorem 6.2 (Marginal identity)**.** *Assume $g \geq 0$, $\tau \leq L$, $C(0) \geq 0$, and $C(m+1) \leq C(m) + g$ for every $m \geq 1$. For continuous P,*

$$\mathrm{marginalJ}(n, g, L, \tau, C, P) = J_T(n, g, L, \tau, C, P).$$

The right-hand functional integrates the squared fibre integral over $\widehat{T}$.

Lean: `marginal_agrees` (`Solution/Fibre.lean:217`).

Proposition 6.3 (Polynomial inner integral)**.** *For arbitrary $U, M \in \mathbb{R}$ and $p, t \in \mathbb{N}$,*

$$\int_0^U (M-u)^p u^t \, du = \sum_{j=0}^{p} \binom{p}{j} M^{p-j} (-1)^j \frac{U^{t+j+1}}{t+j+1}.$$

Lean: `inner_integral` (`Solution/Fibre.lean:240`). The equality follows by expanding the binomial and integrating each power. It holds for arbitrary real U under the oriented interval-integral convention.

7 Two-aggregate Dirichlet integration

Let I and J be disjoint, nonempty finite coordinate groups, with exponent vectors $a \in \mathbb{N}^I$ and $b \in \mathbb{N}^J$. For a nonempty finite group G and an exponent vector e, define

$$\alpha_G(e) = |G| + \sum_{i \in G} e_i - 1, \qquad A_G(e) = \frac{\prod_{i \in G}(e_i!)}{\alpha_G(e)!}.$$

Write $\Delta_{I \sqcup J}(M)$ for the nonnegative simplex with total coordinate sum at most M, and

$$\Delta_2(M) = \{(u_1, u_2) \in \mathbb{R}^2_{\geq 0} : u_1 + u_2 \leq M\}.$$

Theorem 7.1 (Two-aggregate Dirichlet integration)**.** *Suppose I and J are nonempty, and $h : \mathbb{R}^2 \to \mathbb{R}$ is measurable and bounded on $\Delta_2(M)$. With $u_1 = \sum_{i \in I} y_i$ and $u_2 = \sum_{j \in J} y_j$, the weighted simplex integral satisfies*

$$\begin{aligned} &\int_{\Delta_{I \sqcup J}(M)} \left(\prod_{i \in I} y_i^{a_i} \right) \left(\prod_{j \in J} y_j^{b_j} \right) h(u_1, u_2) \, dy \\ &\qquad = A_I(a) A_J(b) \int_{\Delta_2(M)} u_1^{\alpha_I(a)} u_2^{\alpha_J(b)} h(u_1, u_2) \, du_1 \, du_2. \end{aligned} \tag{3}$$

The formalization also gives the version on $\operatorname{Fin} N$ *for a rough coordinate set and its complement.*

Lean: `two_group`, `two_group_fin`, `one_group`; `Solution/Aggregate.lean`. The theorem is proved for the corresponding measurable, bounded integrands; separate theorems handle empty groups and other boundary cases.

Proof outline. Use Fubini and a sum-coordinate shear to isolate the total in one group; apply the weighted simplex moment calculation to the corresponding slices. Repeat for the second group. The factors A_I, A_J record the Dirichlet normalizations; in particular the -1 in each aggregate exponent is essential. The polynomial cases `tri_poly` and `two_group_poly` yield explicit rational expressions, and `two_group_const` recovers the simplex moment formula when $h = 1$.

Proposition 7.2 (A cut-triangle certificate)**.** *With $y = (y_0, y_1, y_2)$, the exact value is*

$$\int_{\substack{y_i \geq 0,\ y_0 + y_1 + y_2 \leq 1 \\ y_0 \leq 1/2}} y_0 y_2^2 \, dy = \frac{19}{7680}.$$

Lean: `tri_cut`, `example_value` (`Solution/Aggregate.lean`). The certificate reduces this integral to $\frac{1}{3} \int_{\Delta_2(1),\, u_1 \leq 1/2} u_1 u_2^3 \, du_1 du_2$; the value has also been cross-checked by direct rational integration.

Connection to marginal geometry. On the translated stratum $x = y + g\mathbf{1}_{R\cup K}$, define $u_1 = \sum_{i\in R} y_i$ and $u_2 = \sum_{i\notin R} y_i$. The proved correspondence `stratumEndpoint_shift` expresses the fibre endpoint as

$$U = \min\big(L - (r+k)g - u_1 - u_2,\ \max\big(g, C(r+1) - rg - u_1\big)\big)\,, \qquad r = |R|,\ k = |K|.$$

The cap index belongs to the stratum R. Substituting the translated point's rough set $R \cup K$ instead changes the endpoint; the Table 3 control records the distinct values 587/1250 and 461/1250. The cap-branch polygon-cell integration is established in Section 8; all endpoint branches and the constant marginal Gram entry are treated in Section 9; the power-sum entry follows in Section 10, and the power-sum-square entry in Section 11, and general generator entries in Section 12.

8 Polygon-cell integration and the cap branch

The two-aggregate reduction produces planar regions bounded by coordinate thresholds and a total-sum constraint. For real A, B, D define the clipped cell

$$\mathcal{B}(A, B, D) = \{(u_1, u_2) \in \mathbb{R}^2 : 0 \le u_1 \le A,\ 0 \le u_2 \le B,\ u_1 + u_2 \le D\}.$$

Theorem 8.1 (Exact polygon-cell moments)**.** *For $\alpha, \beta \in \mathbb{N}$, the Lebesgue integral*

$$\int_{\mathcal{B}(A,B,D)} u_1^{\alpha} u_2^{\beta}\, du_1\, du_2 = \mathrm{boxVal}(A, B, D, \alpha, \beta)$$

where boxVal *is a finite closed form using polynomial antiderivatives. For rational A, B, D its value is rational. The same construction gives rational integrals for arbitrary rational polynomials in two variables. Degenerate and empty cells are treated separately.*

Lean: `box_monomial`, `boxVal_cast`, `box_poly`, `boxVal_empty`, `boxVal_nonpos`, `boxVal_rect`; `Solution/CapCell.lean`.

Proof outline. Integrate first in u_2 with upper endpoint $\min(B, D - u_1)$ and split the u_1 interval at $D - B$. The resulting one-variable powers have exact polynomial antiderivatives. In particular, when $A + B \le D$ the diagonal constraint is redundant and the cell is a rectangle; the formula also covers diagonally truncated cells.

Theorem 8.2 (Cap-branch endpoint and piece integral)**.** *On a translated stratum $x = y + g\mathbf{1}_{R\cup K}$, with $r = |R|$, $k = |K|$ and aggregates u_1, u_2 as in Section 7, the cap branch of the fibre endpoint is*

$$U = C(r+1) - rg - u_1,$$

subject to its vertical and horizontal branch constraints. Combined with the piece constraints, the aggregate domain is a clipped cell $\mathcal{B}(A, B, D)$ with parameters determined by g, L, τ, C, r, k. The weighted piece integrals

$$\int_{\text{piece}} y^t\, \mathbf{1}_{\text{cap}} U^m\, dy$$

therefore equal an explicit expression obtained from the two-aggregate Dirichlet constants and rational cell integral formulas.

Lean: `endpointAgg_on_capCell`, `cap_branch_piece`, `cap_branch_r0`, `cap_inner`; `Solution/CapCell.lean`. The formalization proves the branch classification and corresponding endpoint equalities. For Table 3 the τ constraint is redundant on the cap branch: $\tau - (L - g) = 1/625 > 0$, and the cap cells become rectangles (`tau_ge_table3`, `capCell_rect_table3`).

Proposition 8.3 (A Table 3 cap-piece certificate)**.** *For the specified 44-coordinate piece* $(R, K) = (\{0\}, \varnothing)$*, the cap-branch contribution to the integral defining* $J_T(1)$ *is*

$$\frac{(L - C(2))^{43}((C(2) - g)^3 - g^3)}{3 \cdot 43!} \approx 1.0464 \times 10^{-70}.$$

Lean: `capPieceT3_value` (`Solution/CapCell.lean`). This is *one piece* of the inclusion–exclusion decomposition, not the complete value of $J_T(1)$. The exact rational certificate is kernel-checked. Polygon clipping and lower-dimensional integration provide independent numerical controls; the latter is an approximation, not part of the kernel proof. The remaining branches and their assembly are treated in Section 9.

9 All endpoint branches and the first marginal Gram entry

The translated marginal endpoint is the minimum of the remaining total mass and the maximum of a cap expression and g:

$$U = \min(L - (r + k)g - u_1 - u_2,\ \max(g, C(r + 1) - rg - u_1)).$$

Its switching lines partition each two-aggregate piece into four cells: the cap branch, the remaining-mass branch on the cap side, the constant g branch, and the remaining-mass branch on the g side. Each cell is represented by signed differences of the clipped polygon cells $\mathcal{B}(A, B, D)$ in Section 8.

Theorem 9.1 (Exact branch-cell and piece integrals)**.** *For nonnegative integer exponents, the weighted integral on each branch cell of a polynomial block*

$$u_1^{\alpha} u_2^{\beta} (p - u_1)^{m_1} (L' - u_1 - u_2)^{m_2}$$

reduces to a finite combination of exact clipped-cell moments. The corresponding integrals on translated rough-set strata admit closed forms, including the stratum with no rough coordinates. For integrands depending on the stratum, inclusion–exclusion and cardinality counting assemble these pieces into the marginal functional.

Lean: `box_twoPow`, `cells_integral`, `piece_block`, `piece_r0`, `indicator_supportT_strat`, `count_by_card`; `Solution/MarginalEntry.lean`.

Theorem 9.2 (Rational reduction of the constant-generator marginal)**.** *For* $N \geq 1$ *and parameters satisfying* $0 \leq g \leq C(1)$ *and* $C(N) < Ng$*, the squared-fibre marginal for the constant polynomial is expressed by the explicit rational-piece expression* jtSum*:*

$$J_T(1) = (N + 1) \int_{\widehat{T}} U(\widehat{x})^2 \, d\widehat{x} = \mathrm{jtSum}(N, g, L, \tau, C, 2, 0).$$

Lean: `jT_one_eq`, `jT_one_pieces`, `piece_eq_pieceJ`; `Solution/MarginalEntry.lean`. The reduction uses binomial counting for this symmetric integrand; entries with power-sum factors require additional expansion and placement accounting.

Proposition 9.3 (Table 3 marginal Gram certificate)**.** *Under the Table 3 parameters, for the generator* $G_{0,\varnothing} = 1$*,*

$$J_T[G_{0,\varnothing}, G_{0,\varnothing}] = 45 \int_{\widehat{T}} U(\widehat{x})^2 \, d\widehat{x} \approx 1.1193487646397893 \times 10^{-55}.$$

The exact value is the ratio of these integers, obtained by concatenating each integer's digit lines (no separators):

Numerator

19640530134259849383021060391154484358281020458705508076962340943624589435319293948822187291213238184493255346265105306516171857869871603814187

Denominator

*17546390146399319350401195543355143747760423948217811338346974871720647046191565721073857275769114494323730468750000000000

Lean: `p2EntryVal_table3`, `jT_P2_one_table3`, `jT_one_P2_table3`.
Sources: `Solution/PowerSumEntry.lean`, `Solution/PowerSumEntryCert.lean`.

The exact Table 3 evaluation uses 133 cardinality-class pieces with two certificate lemmas per piece (266 lemmas total); the remaining classes are eliminated by `cond_tail` and `cond_high`. The generator `specs/checks/power_sum_entry_v2g_emit.py` reproduces the Lean certificate byte-for-byte on the VPS. Six independent Lean-file controls have their expected outcomes: two accepted positive controls and four rejected negative controls. Numerical cross-checks include a low-dimensional grid and Monte Carlo; these support the arithmetic independently of the exact kernel proof. The power-sum-square diagonal entry $J_T[P_2, P_2]$, including two distinguished-coordinate placements and their diagonal case, is proved in Section 11.

11 A two-coordinate moment table and the power-sum-square entry (V2h)

The marginal square of the nonconstant generator is

$$J_T[P_2, P_2] = 45 \int_{\widehat{T}} \left(U\widehat{P}_2 + \frac{U^3}{3} \right)^2 d\widehat{x} \tag{5}$$

$$= 45 \int_{\widehat{T}} \left(U^2\widehat{P}_2^2 + \frac{2}{3}U^4\widehat{P}_2 + \frac{1}{9}U^6 \right) d\widehat{x}. \tag{6}$$

The new term contains two distinguished coordinate indices:

$$\widehat{P}_2^2 = \sum_{i,j} x_i^2 x_j^2.$$

For $i = j$ a translated summand becomes $(y_i + s_i)^4$; for $i \neq j$ we obtain $(y_i + s_i)^2(y_j + s_j)^2$, where s_i is the stratum translation (g for coordinates in $R \cup K$, zero otherwise). The three placement classes R, K, and their complement have sizes n_c; the number of ordered distinct-coordinate pairs in classes c, d is $n_c n_d - \mathbf{1}_{c=d} n_c$. The diagonal contributes n_c placements per class.

Theorem 11.1 (Moment-table reduction for the marginal square)**.** *For each translated stratum and each exponent vector t, the weighted piece integral of $y^t U^m$ reduces to a factorial factor $\prod_i t_i!$ multiplied by an exact moment determined by the total monomial degrees in the rough and smooth coordinate groups. Together with the diagonal and distinct-coordinate placement formulas, this gives an exact finite rational representation of* (5)*.*

Lean development: `BGP212/PowerSumPlace.lean`,
`BGP212/PowerSumSquare.lean`, `Solution/PowerSumSquare.lean`;
`Challenge/PowerSumSquare.lean`. The proof reuses the two-aggregate and polygon-cell machinery, with moments shared among coordinate placements rather than recomputed for each summand.

Proposition 11.2 (Table 3 power-sum-square certificate)**.** *At the specified Table 3 parameters, the entry $J_T[P_2, P_2]$ is an exact rational number represented by the V2h Lean certificate, with*

$$J_T[P_2, P_2] \approx 2.27107 \times 10^{-58}.$$

The certificate is organized into 14 resumable row files and a final assembly file, each checked as part of the V2h Comparator proof.

Certificate sources: `Solution/PowerSumSquareCert0.lean` through
`Solution/PowerSumSquareCert13.lean`,
`Solution/PowerSumSquareCertSum.lean`;
`specs/checks/power_sq_entry_v2h_emit.py`. The 27-statement Comparator configuration is `Comparator/power_sum_square_v2h.json`. Its final VPS output states that the Lean default kernel accepts the solution and reports "Your solution is okay!".

The first 2×2 marginal Gram block. Combining V2f, V2g, and V2h yields the exact rational entries of

$$\mathcal{G}_{\{1,P_2\}} = \begin{pmatrix} J_T[1,1] & J_T[1,P_2] \\ J_T[P_2,1] & J_T[P_2,P_2] \end{pmatrix} \approx \begin{pmatrix} 1.11935 \times 10^{-55} & 4.99944 \times 10^{-57} \\ 4.99944 \times 10^{-57} & 2.27107 \times 10^{-58} \end{pmatrix}.$$

Using the displayed approximations, the Cauchy–Schwarz ratio

$$\frac{J_T[P_2,1]^2}{J_T[1,1]\, J_T[P_2,P_2]} \approx 0.984$$

is close to one. This numerical observation motivates exact assessment of determinants and variational certificates; an independent Lean certificate for a strictly positive 2×2 determinant is not asserted here. Independent Monte Carlo and low-dimensional grid comparisons test the computed value but are distinct from the Lean proof.

12 General polynomial marginal Gram entries (V2i)

The preceding certificates evaluate a small marginal Gram block. V2i extends the exact representation to arbitrary pairs of the generators $G_{a,\lambda} = (L - P_1)^a P_\lambda$, where P_λ is a product of power sums. If $M = L - \sum_i \widehat{x}_i$ and the fibre endpoint is U, then

$$\int_0^U G_{a,\lambda}(\widehat{x}, u)\, du = \sum_{T \subseteq \lambda} \widehat{P}_{\lambda \setminus T} \sum_{j=0}^{a} \binom{a}{j} (-1)^j \frac{M^{a-j} U^{s_T+j+1}}{s_T + j + 1}, \qquad s_T = \sum_{q \in T} q. \tag{7}$$

Here the subset sum counts selections of indexed factors (including repeated parts with their multiplicities). The formula is proved as `inner_genFun` in `Solution/GenEntry.lean`.

Translated placements and common moments. On each stratum, a factor of $\widehat{P}_q$ is a sum of powers of translated coordinates. The recursive placement calculation either attaches a new factor to an already distinguished coordinate or opens a coordinate in one of the three classes (rough, intermediate, or remaining). The latter has multiplicity equal to the class population minus the number of already selected coordinates in that class. Unrolling the recursion accounts for set partitions and falling-factorial multiplicities. The resulting monomial weights reduce to a factorial factor times a moment indexed by the total rough-group and smooth-group degrees. The principal corresponding Lean declarations include `piece_place`, `piece_state`, and `piece_mono_n`.

Theorem 12.1 (General marginal Gram representation)**.** *For every $N \geq 1$, parameters satisfying the admissibility assumptions $0 \leq g \leq C(1)$ and $C(N) < Ng$, and arbitrary generator indices (a_1, λ_1) and (a_2, λ_2), the bilinear marginal form is an explicit finite expression:*

$$J_T[G_{a_1,\lambda_1}, G_{a_2,\lambda_2}] = \mathrm{genEntryVal}(N, g, L, \tau, C, a_1, \lambda_1, a_2, \lambda_2).$$

For the rational Table 3 parameters, the expression is rational; every pair of generators indexed by B_{21} therefore has an exact rational representation.

Lean: `genEntry_eq`, `jT_gen_table3`; `Solution/GenEntry.lean`, `Solution/GenEntryTable.lean`. The latter theorem supplies a rational representation for all entries; it does not assert that every entry of the numerical matrix has been evaluated.

Proposition 12.2 (Selected Table 3 marginal certificates)**.** *Under the specified Table 3 parameters the exact kernel-checked rational values have the following approximations:*

$$\begin{aligned} J_T[P_2, P_2] &\approx 2.27107 \times 10^{-58}, \\ J_T[P_4, 1] &\approx 2.75169 \times 10^{-59}, \\ J_T[P_{(2,2)}, 1] &\approx 2.28799 \times 10^{-58}, \\ J_T[L - P_1, 1] &\approx 5.52809 \times 10^{-57}. \end{aligned}$$

The first value is recovered exactly from the independently checked V2h certificate by `genEntryPP_eq_v2h`*; the other displayed values are further applications of the general formula.*

Lean: `jT_P2_P2_gen_table3`, `jT_P4_one_table3`, `jT_P22_one_table3`, `jT_LP1_one_table3`. The supporting rational moment certificates are compiled in `Solution/GenEntryCert0.lean` through `Solution/GenEntryCert13.lean` and `Solution/GenEntryCertSum.lean`. The generator and controls are in `specs/checks/gen_entry_v2i_emit.py` and related `specs/checks/gen_entry_v2i_*.py` files.

Computational scope. The V2i formula covers every pair of generators, whereas compiling an entire numerical Gram matrix would require many further exact evaluations. The optimized polynomial $P^\star$, the exact sign of $J_T(P^\star) - 4I_T(P^\star)$, and the remaining variational conditions are not established here. The proposed next step is to collect coefficients for repeated rational moments across generator pairs before evaluating those moments, preserving an exact equality with the V2i formula.

13 Verification, limitations, and further mathematics

The project uses Lean `v4.34.0-rc2` and a pinned Mathlib revision `85e3a25`. At the V2g baseline, all 19 Comparator suites were independently accepted on the VPS. Four positive V2d controls were accepted; four intentionally incorrect controls were rejected with unsolved goals. The V2d proof commit is `52fd36a`; its published verification record is commit `90055db`. For V2e-A, five positive controls were accepted and four negative controls were rejected. The V2e-A proof commit is `01644e6`, with VPS verification/provenance commit `38606c0`. For V2e-B, five positive controls were accepted and five deliberately incorrect controls rejected; the V2e-B proof commit is `6a21780`, with verification/provenance commit `25a7376`. For V2f, four positive Lean-file controls were accepted and five negative controls rejected. Its proof commit is `8c50fb6`, with VPS provenance commit `da0c3fd`. For V2g, two positive controls were accepted and four negative controls rejected; its 266 certificate lemmas regenerated byte-identically on the VPS. The V2g proof commit is `ad70abb`, with verification/provenance commit `12f8b16`. For V2h, the 27-statement Comparator accepted the power-sum-square certificate on the VPS. Its proof commit is `6cef1f4`, with VPS provenance commit `a9b0e0c`. The recorded V2h proof includes 14 rational moment-certificate rows and an assembly theorem. The final Comparator acceptance is confirmed; a complete 20/20 regression count has not been established from a saved log. For V2i, the 33-statement Comparator suite was accepted by the Lean default kernel on the VPS. The complete regression log records **21/21** "Your solution is okay!" messages and the verification script's **exit status 0**. The proof snapshot is `a994d73`; its separately published VPS provenance commit is `486cf85`. The V2i record and Comparator configuration are `specs/results/V2i_gen_entry.yaml` and `Comparator/gen_entry_v2i.json`. Prior control results and exploratory numerical comparisons are distinct from the saved VPS kernel acceptances. The reported kernel axioms for these developments are `propext`, `Quot.sound`, and `Classical.choice`. Results and verification scripts are available in `specs/results`.

Further work evaluates additional entries using the general V2i representation, combines shared rational moments efficiently, and addresses optimized polynomial coefficients, exact variational positivity certificates, and remaining analytic hypotheses. The existing Lean development formalizes the *conditional* implication $\mathrm{DHL}[45, 2] \Rightarrow H_1 \leq 212$; this manuscript does not claim an unconditional formal proof of $H_1 \leq 212$.

Acknowledgment and provenance

The bounded-prime-gap application is motivated by Axiom Math's mathematical argument [1]. This manuscript records an independent Lean formalization of the specified results, with theorem names and commit identifiers to support reproducible review. Substantive exposition and attribution will be expanded in a subsequent draft.

References

[1] Axiom Math, *A New Bound for Small Gaps Between Primes*, preliminary draft, 3 September 2026. Project page: https://primegaps.axiommath.ai/.

[2] Reading Point, *lean-bgp212*: Lean source, verification records, and certificates. https://github.com/thinkthoughts/lean-bgp212.