Exact Rational Integration over Constrained Simplices

Symmetric Polynomial Bases, Marginal Geometry, and Lean 4 Verification

Reading Point (translatepi.ai)

Draft v0.1 — 18 October 2026

Abstract

We present a formalized collection of integration identities for polynomial weights on simplexes constrained by coordinate thresholds and cardinality-dependent caps. The development includes exact simplex moments, a rational polynomial subspace of dimension 846, cap-parametric formulas for arbitrary monomial integrals, explicit rational representations of associated Gram entries, and a fibre decomposition for marginal functionals. These statements are formalized in Lean 4 and are supported by independently replayed Comparator suites and boundary controls. The construction arose in the analysis of a bounded-prime-gap argument, but this manuscript concerns the integration and polynomial mathematics in its own right. General marginal Gram evaluations and the remaining variational and analytic obligations lie outside the completed results.

1 Introduction and scope

Exact integration on constrained regions is useful where a variational inequality must ultimately be checked using rational arithmetic. A conventional simplex has a closed monomial moment formula; the present support also depends on which coordinates cross a threshold and on a cap for the sum of those coordinates. This introduces a finite stratification, but preserves a route to exact, rational expressions.

We study the mathematical constructions implemented in the public Lean repository `thinkthoughts/lean-bgp212`. The motivating bounded-prime-gap argument originates with Axiom Math [1]; the statements below are an independent formalization of specified components and do not amount to a complete formal proof of the unconditional prime-gap bound. References to Lean declarations are included to delimit precisely what has been checked. The short arguments given here are *proof outlines*, not substitutes for the complete Lean proof terms.

2 Polynomial moments on a simplex

Definition 2.1. *For $m \in \mathbb{N}$ and $R \in \mathbb{R}$, define*

$$\Delta_m(R) = \left\{ x \in \mathbb{R}^m : x_i \geq 0 \text{ for all } i, \quad \sum_{i=1}^{m} x_i \leq R \right\}.$$

Theorem 2.2 (Simplex monomial moments)**.** *For $m \in \mathbb{N}$, $R \geq 0$, and $e \in \mathbb{N}^m$,*

$$\int_{\Delta_m(R)} \prod_{i=1}^{m} x_i^{e_i}\, dx = \frac{R^{m+\sum_i e_i} \prod_i (e_i!)}{(m + \sum_i e_i)!}.$$

Here the dimension-zero case uses the corresponding Lean integral convention.

Lean: `simplex_integral`; definitions `simplex`, `momentR`.
Source: `Solution/SimplexMoment.lean:160`.

Proof outline. Induct on dimension and use Fubini to isolate a coordinate. The inner integral is a Beta integral; the factorial expression then simplifies by induction. The formal development additionally treats negative and zero radii, simplex volume, and rational-polynomial integrals (`poly_integral`).

3 An indexed rational polynomial space

Consider rational polynomials in 45 coordinates. Write P_j for the jth power sum, P_λ for the product over parts of a multiset λ, and set

$$G_{a,\lambda} = (L - P_1)^a P_\lambda, \qquad \lambda \text{ has positive even parts}, \quad a + |\lambda| \le 21.$$

The index type $B_{21}\mathrm{Idx}$ enumerates the admissible pairs.

Theorem 3.1 (Independence and rank)**.** *The indexed family $G_{a,\lambda}$ is linearly independent over $\mathbb{Q}$. Its rational span B_{21} has*

$$\dim_{\mathbb{Q}} B_{21} = 846.$$

Lean: `genIdx_linearIndependent`, `finrank_B21` (`Solution/Variational.lean:398,410`).

Proof outline. The indexing is injective; distinct indexed monomials are linearly independent. Injective polynomial transformations preserve this independence. A separate count establishes $|B_{21}\mathrm{Idx}| = 846$, and the span-rank identity gives the result. The theorem concerns this designated subspace, not the full polynomial ring.

4 Cap-constrained supports and general integrals

Let $N \in \mathbb{N}$, $g \ge 0$, $L \in \mathbb{R}$, and $C : \mathbb{N} \to \mathbb{R}$. For $x \in \mathbb{R}^N$ write

$$R_g(x) = \{i : x_i > g\}, \qquad s_g(x) = |R_g(x)|, \qquad \rho_g(x) = \sum_{i \in R_g(x)} x_i.$$

Define the capped simplex support

$$T(N, g, L, C) = \left\{ x \in \mathbb{R}^N : x_i \ge 0, \quad \sum_i x_i \le L, \quad \rho_g(x) \le C(s_g(x)) \right\}. \tag{1}$$

Theorem 4.1 (General monomial integral)**.** *For $N \in \mathbb{N}$, $g \ge 0$, $L \in \mathbb{R}$, $C : \mathbb{N} \to \mathbb{R}$, $e \in \mathbb{N}^N$, and $p \in \mathbb{N}$,*

$$\int_{T(N,g,L,C)} \left(L - \sum_i x_i \right)^p \prod_i x_i^{e_i} \, dx = \mathrm{gramGen}(N, g, L, C, e, p). \tag{2}$$

The right-hand side is the explicit, finite expression defined in Lean.

Lean: `integral_supportT_gen`; `gramGen`.
Sources: `Solution/GramGeneral.lean:344`;
`BGP212/GramGeneral.lean`.

Proof outline. Stratify by the rough-coordinate set, apply inclusion–exclusion to threshold conditions, and translate the resulting pieces to weighted simplexes. Use Theorem 2.2 and Fubini to evaluate the polynomial weights. Placement-counting identities `sig_insert`, `sig_union`, and `sig_card` reduce the sums to the finite closed form. The reduction `gramGen_extend` transports calculation to the coordinates having nonzero exponents. This is an algebraic efficiency result, not a claim that all entries are computationally cheap to certify.

5 Rational Gram representations

Specialize to the fixed rational cap values and support parameters of Table 3 in [1], with $N = 45$. For indices $i, j \in B_{21}\mathrm{Idx}$ define

$$I_T(i,j) = \int_T G_i(x) G_j(x)\, dx.$$

Theorem 5.1 (Explicit rational representation)**.** *For every $i, j \in B_{21}\mathrm{Idx}$, $I_T(i,j)$ equals the real embedding of a finite rational sum indexed by placements φ of the combined power-sum parts:*

$$I_T(i,j) = \left(\sum_{\varphi} \mathrm{gramGen}_{\mathbb{Q}}(45, g, L, C, e_\varphi, a_i + a_j) \right)_{\mathbb{R}}.$$

Here e_φ is the exponent vector obtained from the placement, exactly as specified by `expo` *in the Lean source.*

Lean: `gram_entry_B21_rat` (`Solution/GramGeneral.lean:522`); `gram_entry_gen_list` and `gram_entry_example`.

The theorem gives each entry an explicit rational representation; it does not supply an evaluated and certified 846×846 matrix. One worked entry, corresponding to $(L - P_1)P_2^2$, has its value certified using a collection of small exact rational lemmas. This separation between an algebraic identity and its numerical certificate is important to reproducibility.

6 Marginal-fibre geometry

Let $\widehat{x} \in \mathbb{R}^n$ and append $u \in \mathbb{R}$ as the last coordinate. Define $S = \sum_i \widehat{x}_i$, $s = s_g(\widehat{x})$, $\rho_1 = \rho_g(\widehat{x})$, and

$$\widehat{T} = \{\widehat{x} \geq 0 : S \leq \tau,\ \rho_1 \leq C(s)\}, \quad U(\widehat{x}) = \min\{L - S, \max(g, C(s+1) - \rho_1)\}.$$

Theorem 6.1 (Fibre equivalence)**.** *Suppose $g \geq 0$, $\tau \leq L$, and $\widehat{x} \in \widehat{T}$. Then for every $u \in \mathbb{R}$,*

$$(\widehat{x}, u) \in T(n+1, g, L, C) \quad \Longleftrightarrow \quad 0 \leq u \leq U(\widehat{x}).$$

Lean: `fibre_equiv` (`Solution/Fibre.lean:115`).

Proof outline. Appending u leaves the rough set unchanged for $u \leq g$, and increments both rough count and rough mass for $u > g$. Combining the resulting cap inequality with $S + u \leq L$ identifies the minimum–maximum endpoint exactly. The empty-fibre theorem for cap-violating outer coordinates additionally needs $C(m+1) \leq C(m) + g$ for $m \geq 1$, and $C(0) \geq 0$. No constraint on the step $m = 0$ is assumed; the relevant Table 3 facts are checked in Lean.

Theorem 6.2 (Marginal identity)**.** *Assume $g \geq 0$, $\tau \leq L$, $C(0) \geq 0$, and $C(m+1) \leq C(m) + g$ for every $m \geq 1$. For continuous P,*

$$\mathrm{marginalJ}(n, g, L, \tau, C, P) = J_T(n, g, L, \tau, C, P).$$

The right-hand functional integrates the squared fibre integral over $\widehat{T}$.

Lean: `marginal_agrees` (`Solution/Fibre.lean:217`).

Proposition 6.3 (Polynomial inner integral)**.** *For arbitrary $U, M \in \mathbb{R}$ and $p, t \in \mathbb{N}$,*

$$\int_0^U (M - u)^p u^t\, du = \sum_{j=0}^{p} \binom{p}{j} M^{p-j} (-1)^j \frac{U^{t+j+1}}{t+j+1}.$$

Lean: `inner_integral` (`Solution/Fibre.lean:240`). The equality follows by expanding the binomial and integrating each power. It holds for arbitrary real U under the oriented interval-integral convention.

7 Verification, limitations, and further mathematics

The project uses Lean `v4.34.0-rc2` and a pinned Mathlib revision `85e3a25`. At the V2d baseline, 15 Comparator suites were independently accepted on the VPS. Four positive V2d controls were accepted; four intentionally incorrect controls were rejected with unsolved goals. The V2d proof commit is `52fd36a`; its published verification record is commit `90055db`. The reported kernel axioms for the V2d development are `propext`, `Quot.sound`, and `Classical.choice`. Results and verification scripts are available in `specs/results`.

Further work is needed for aggregate Dirichlet integrals over functions of two group sums, all branches of the marginal endpoint, general J_T Gram evaluations, optimized polynomial coefficients, exact variational positivity certificates, and remaining analytic hypotheses. The existing Lean development formalizes the *conditional* implication $\mathrm{DHL}[45,2] \Rightarrow H_1 \leq 212$; this manuscript does not claim an unconditional formal proof of $H_1 \leq 212$.

Acknowledgment and provenance

The bounded-prime-gap application is motivated by Axiom Math's mathematical argument [1]. This manuscript records an independent Lean formalization of the specified results, with theorem names and commit identifiers to support reproducible review. Substantive exposition and attribution will be expanded in a subsequent draft.

References

[1] Axiom Math, *A New Bound for Small Gaps Between Primes*, preliminary draft, 3 September 2026. Project page: `https://primegaps.axiommath.ai/`.

[2] Reading Point, *lean-bgp212*: Lean source, verification records, and certificates. `https://github.com/thinkthoughts/lean-bgp212`.