

- (1) Show that there is no set  $V$  such that every set is a member of  $V$ .

**Solution:** Suppose not and let  $V$  be a set such that every set is a member of  $V$ . Define  $W = \{x \in V : x \notin x\}$ . Then  $W$  is a set by the axiom of comprehension. Since  $W$  is a set, either  $W \in W$  or  $W \notin W$ . If  $W \in W$ , then since  $W \in V$ , we must have  $W \notin W$ . Similarly, if  $W \notin W$ , then  $W \in W$ . In either case we get a contradiction. Hence  $V$  does not exist.  $\square$

- (2) Show that  $(x, y) = (a, b)$  iff  $x = a$  and  $y = b$ .

**Solution:** The right to left implication is obvious. So assume  $(x, y) = (a, b)$  and we'll show  $x = a$  and  $y = b$ . We consider two cases.

**Case  $x = y$ :** In this case,  $(x, y) = \{\{x\}, \{x, y\}\} = \{\{x\}\}$ . Hence  $(a, b) = \{\{a\}, \{a, b\}\} = \{\{x\}\}$ . It follows that  $\{x\} = \{a\} = \{a, b\}$ . So  $x = a = b$ . Hence  $x = y = a = b$ .

**Case  $x \neq y$ :** In this case,  $\{\{x\}, \{x, y\}\}$  is a set with two distinct members. It follows that  $\{\{a\}, \{a, b\}\}$  is also a set with two distinct members. So  $a \neq b$ . Now  $\{x\} \in \{\{a\}, \{a, b\}\}$  implies  $\{x\} = \{a\}$  since  $\{x\} \neq \{a, b\}$  (as the latter has two distinct members). Similarly,  $\{x, y\} = \{a, b\}$ . As  $x = a$ , and  $y \neq x$ , we get  $y = b$ .  $\square$

- (3) Suppose  $R$  is an equivalence relation on  $A$ . For each  $a \in A$ , define the  $R$ -equivalence class of  $a$  by  $[a] = \{b \in A : aRb\}$ . Show that  $\{[a] : a \in A\}$  is a partition of  $A$ . Furthermore, show that for every partition  $\mathcal{F}$  of  $A$ , there is an equivalence relation  $S$  on  $A$  such that  $\mathcal{F}$  is the set of all  $S$ -equivalence classes.

**Solution:** To show that  $\{[a] : a \in A\}$  is a partition of  $A$ , we need to show that  $\bigcup \{[a] : a \in A\} = A$  and for any two distinct  $R$ -equivalence classes  $[a], [b]$ , we must have  $[a] \cap [b] = \emptyset$ .

Since  $R$  is a reflexive relation on  $A$ , for every  $a \in A$ ,  $a \in [a]$ . Hence  $A \subseteq \bigcup \{[a] : a \in A\}$ . As  $[a] \subseteq A$  for every  $a \in A$ , we also have  $\bigcup \{[a] : a \in A\} \subseteq A$ . Thus  $\bigcup \{[a] : a \in A\} = A$ .

Next, towards a contradiction, suppose  $a, b \in A$ ,  $[a] \neq [b]$  and  $[a] \cap [b] \neq \emptyset$ . Fix  $c \in [a] \cap [b]$ . Since  $c \in [a]$ , we get  $aRc$ . Similarly,  $bRc$ . Since  $R$  is symmetric, it follows that  $cRb$ . Since  $aRc$  and  $cRb$ , using the fact that  $R$  is transitive, we get  $aRb$  and hence also  $bRa$  (as  $R$  is symmetric). We now claim the following.

$[a] \subseteq [b]$ : Fix  $x \in [a]$ . Then  $aRx$ . As  $bRa$ , by transitivity of  $R$ , we get  $bRx$ . Hence  $x \in [b]$ . So  $[a] \subseteq [b]$ .

$[b] \subseteq [a]$ : Fix  $y \in [b]$ . Then  $bRy$ . As  $aRb$ , by transitivity of  $R$ , we get  $aRy$ . Hence  $y \in [a]$ . So  $[b] \subseteq [a]$ .

It follows that  $[a] = [b]$  which contradicts our assumption that  $[a] \neq [b]$ . This finishes the proof that  $\{[a] : a \in A\}$  is a partition of  $A$ .

Now fix a partition  $\mathcal{F}$  of  $A$  and define a relation  $S$  on  $A$  as follows. For  $a, b \in A$ ,  $aSb$  iff there exists  $E \in \mathcal{F}$  such that both  $a$  and  $b$  are members of  $E$ .

Let us first check that  $S$  is an equivalence relation on  $A$ . It is clear that  $S$  is a symmetric relation on  $A$ . Since  $\bigcup \mathcal{F} = A$ , it follows that  $S$  is reflexive. Next suppose  $aSb$  and  $bSc$ . Fix  $E, F \in \mathcal{F}$  such that  $a, b \in E$  and  $b, c \in F$ . Since  $\mathcal{F}$  has pairwise disjoint members and since  $E \cap F \neq \emptyset$ , we must have  $E = F$ . Hence  $aSc$ . So  $S$  is transitive. It follows that  $S$  is an equivalence relation on  $A$ .

Finally, let us check that the set  $\{[a] : a \in A\}$  of  $S$ -equivalence classes is equal to  $\mathcal{F}$ . Let  $[a]$  be an  $S$ -equivalence class. Fix  $E \in \mathcal{F}$  such that  $a \in E$ . Then by the definition of  $S$ , it follows that  $[a] = \{b \in A : aSb\} = \{b \in A : b \in E\} = E$ . Conversely, if  $E \in \mathcal{F}$ , then for every  $a \in E$ ,  $[a] = E$ . Hence  $\{[a] : a \in A\} = \mathcal{F}$ .  $\square$

- (4) Suppose  $X \subseteq \omega$  satisfies  $0 \in X$  and  $(\forall y \in X)(y \cup \{y\} \in X)$ . Show that  $X = \omega$ .

**Solution.** By assumption,  $X$  is inductive and  $\omega$  is the intersection of all inductive sets, so  $\omega \subseteq X$ . As  $X \subseteq \omega$ ,  $X = \omega$ .  $\square$

- (5) Let  $(L, \prec)$  be a linear ordering. Prove the following.

(a)  $(L, \prec)$  is a well-ordering iff there is no sequence  $\langle x_n : n < \omega \rangle$  in  $L$  such that  $(\forall n < \omega)(x_{n+1} \prec x_n)$ .

(b)  $(L, \prec)$  is a well-ordering iff for every  $A \subseteq L$ ,  $(A, \prec)$  is isomorphic to an initial segment of  $(L, \prec)$ .

**Solution:** (a) First suppose that  $(L, \prec)$  is a well-ordering. We'll show that there is no  $\prec$ -decreasing sequence in  $L$ . Towards a contradiction, suppose there is a sequence  $\langle x_n : n < \omega \rangle$  in  $L$  such that for every  $n < \omega$ ,  $x_{n+1} \prec x_n$ . Let  $A = \{x_n : n < \omega\}$  be the range of this sequence. Then  $A$  has no  $\prec$ -least member which contradicts the fact that  $(L, \prec)$  is a well-ordering.

Now suppose  $(L, \prec)$  is not a well-ordering and fix a nonempty  $A \subseteq L$  such that  $A$  does not have a  $\prec$ -least member. We'll construct a  $\prec$ -decreasing sequence  $\langle x_n : n < \omega \rangle$  in  $L$ . Using the axiom of choice, fix a choice function  $F : \mathcal{P}(A) \setminus \{\emptyset\} \rightarrow A$ . So for every nonempty  $W \subseteq A$ ,  $F(W) \in W$ . By recursion on  $n < \omega$ , define  $\langle x_n : n < \omega \rangle$  as follows.  $x_0 = F(A)$  and for every  $n < \omega$ ,

$$x_{n+1} = F(\{x \in A : x \prec x_n\})$$

Note that this is well-defined since  $\{x \in A : x \prec x_n\}$  is nonempty (as  $A$  has no  $\prec$ -least member). It is clear that  $\langle x_n : n < \omega \rangle$  is as required.  $\square$

(b) First suppose  $(L, \prec)$  is a well-ordering. Fix  $A \subseteq L$ . We'll construct an isomorphism from  $(A, \prec)$  to an initial segment of  $(L, \prec)$ . Define

$$f = \{(x, a) \in L \times A : (\text{pred}(L, \prec, x), \prec) \cong (\text{pred}(A, \prec, a), \prec)\}$$

(i)  $f$  is a function: Clearly,  $f$  is a relation. To see that it is a function, fix  $(x, a), (x, b) \in f$  and we'll show that  $a = b$ . Towards a contradiction suppose  $a \neq b$ . Without loss of generality suppose  $a \prec b$ . Since  $(x, a)$  and  $(x, b)$  are both in  $f$ , we get  $(\text{pred}(L, \prec, x), \prec) \cong (\text{pred}(A, \prec, a), \prec)$  and  $(\text{pred}(L, \prec, x), \prec) \cong (\text{pred}(A, \prec, b), \prec)$ . Hence  $(\text{pred}(A, \prec, a), \prec) \cong (\text{pred}(A, \prec, b), \prec)$ . But this means that  $(\text{pred}(A, \prec, b), \prec)$  is a well-ordering that is isomorphic to a proper initial segment of itself. Contradiction. So  $f$  is a function.

(ii)  $f$  is injective: The proof is similar to (i) above.

(iii)  $\text{dom}(f)$  is an initial segment of  $(L, \prec)$ : Suppose  $x \in \text{dom}(f)$  and  $y \prec x$ . We need to show that  $y \in \text{dom}(f)$ . Let  $f(x) = a$ . Fix an isomorphism  $h : (\text{pred}(L, \prec, x), \prec) \rightarrow (\text{pred}(A, \prec, a), \prec)$ . Note that  $y \in \text{dom}(h)$ . Let  $h(y) = b$ . It is clear that  $h \upharpoonright \text{pred}(L, \prec, y)$  is an isomorphism from  $(\text{pred}(L, \prec, y), \prec)$  to  $(\text{pred}(A, \prec, b), \prec)$ . Hence  $(y, b) \in f$  and so  $y \in \text{dom}(f)$ .

(iv)  $\text{range}(f)$  is an initial segment of  $(A, \prec)$ : The proof is similar to (iii) above.

(v)  $f$  is an isomorphism from  $(\text{dom}(f), \prec)$  to  $(\text{range}(f), \prec)$ : Suppose  $x \prec y$  are in  $\text{dom}(f)$ . Put  $a = f(x)$  and  $b = f(y)$ . Using the definition of  $f$ , it follows that  $(\text{pred}(L, \prec, x), \prec) \cong (\text{pred}(A, \prec, a), \prec)$  and  $(\text{pred}(L, \prec, y), \prec) \cong (\text{pred}(A, \prec, b), \prec)$ . As  $x \prec y$ , it follows that  $(\text{pred}(A, \prec, a), \prec)$  is isomorphic to an initial segment of  $(\text{pred}(A, \prec, b), \prec)$ . Since no well-ordering can be isomorphic to a proper initial-segment of itself, it follows that  $a \prec b$ . So  $f$  is an isomorphism from  $(\text{dom}(f), \prec)$  to  $(\text{range}(f), \prec)$ .

(vi)  $\text{range}(f) = A$ : Suppose not. Let  $a = \min(A \setminus \text{range}(f))$ . Since  $\text{range}(f)$  is an initial segment of  $(A, \prec)$ , it follows that  $\text{range}(f) = \text{pred}(A, \prec, a)$ . We claim that  $\text{dom}(f) = L$ . For suppose not and let  $x = \min(L \setminus \text{dom}(f))$ . Then  $\text{dom}(f) = \text{pred}(L, \prec, x)$ . But this implies that  $(a, b) \in f$  using (i)-(v) above which is a contradiction. So  $\text{dom}(f) = L$ . Hence  $a \in \text{dom}(f)$ . Now observe that  $f(a) \prec a$  (since  $\text{range}(f) = \text{pred}(A, \prec, a)$ ) and iteratively applying  $f$ , we get  $a \succ f(a) \succ f(f(a)) \succ \dots$ . But this means that  $(L, \prec)$  has an infinite  $\prec$ -descending sequence which is impossible by part (a).

(i)-(vi) imply that  $(A, \prec)$  is isomorphic (via  $f^{-1}$ ) to an initial segment of  $(L, \prec)$  (namely  $\text{dom}(f)$ ).

Next we show the converse. Suppose for every  $A \subseteq L$ ,  $(A, \prec)$  is isomorphic to an initial segment of  $(L, \prec)$ . We'll show that  $(L, \prec)$  must be a well-ordering. We can assume that  $L \neq \emptyset$ . Let  $a \in L$ . Then  $(\{a\}, \prec)$  is isomorphic to an initial segment of  $(L, \prec)$ . This implies that  $L$  has a  $\prec$ -least element, say  $x$ . Now let  $A \subseteq L$  be nonempty and fix an isomorphism  $f : (A, \prec) \rightarrow (W, \prec)$  where  $W$  is an initial segment of  $(L, \prec)$ . Note that  $x \in W$ . Put  $a = f^{-1}(x)$ . Then  $a$  is the  $\prec$ -least element of  $A$ . It follows that  $(L, \prec)$  is a well-ordering.  $\square$

- (6) Suppose  $(X, \prec_1)$  and  $(Y, \prec_2)$  are well-orderings. Then exactly one of the following holds.

- (a)  $(X, \prec_1) \cong (Y, \prec_2)$ .
- (b) For some  $x \in X$ ,  $(\text{pred}(X, \prec_1, x), \prec_1) \cong (Y, \prec_2)$ .
- (c) For some  $y \in Y$ ,  $(\text{pred}(Y, \prec_2, y), \prec_2) \cong (X, \prec_1)$ .

Furthermore, in each of the three cases, the isomorphism is unique.

**Solution:** Define

$$f = \{(a, b) \in X \times Y : (\text{pred}(X, \prec_1, a), \prec_1) \cong (\text{pred}(Y, \prec_2, b), \prec_2)\}$$

(i)  $f$  is a function: Clearly,  $f$  is a relation. To see that it is a function, fix  $(a, b), (a, c) \in f$  and we'll show that  $b = c$ . Towards a contradiction suppose  $b \neq c$ . Without loss of generality suppose  $b \prec_2 c$ . Since  $(a, b)$  and  $(a, c)$  are both in  $f$ , we get  $(\text{pred}(X, \prec_1, a), \prec_1) \cong (\text{pred}(Y, \prec_2, b), \prec_2)$  and  $(\text{pred}(X, \prec_1, a), \prec_1) \cong (\text{pred}(Y, \prec_2, c), \prec_2)$ . Hence  $(\text{pred}(Y, \prec_2, c), \prec_2) \cong (\text{pred}(Y, \prec_2, b), \prec_2)$ . But this means that  $(\text{pred}(Y, \prec_2, c), \prec_2)$  is a well-ordering that is isomorphic to a proper initial segment of itself. Contradiction. So  $f$  is a function.

(ii)  $f$  is injective: The proof is similar to (i) above.

(iii)  $\text{dom}(f)$  is an initial segment of  $(X, \prec_1)$ : Suppose  $x \in \text{dom}(f)$  and  $y \prec_1 x$ . We need to show that  $y \in \text{dom}(f)$ . Let  $f(x) = a$ . Fix an isomorphism  $h : (\text{pred}(X, \prec_1, x), \prec_1) \rightarrow (\text{pred}(Y, \prec_2, a), \prec_2)$ . Note that  $y \in \text{dom}(h)$ . Let  $h(y) = b$ . It is clear that  $h \upharpoonright \text{pred}(X, \prec_1, y)$  is an isomorphism from  $(\text{pred}(X, \prec_1, y), \prec_1)$  to  $(\text{pred}(Y, \prec_2, b), \prec_2)$ . Hence  $(y, b) \in f$  and so  $y \in \text{dom}(f)$ .

(iv)  $\text{range}(f)$  is an initial segment of  $(A, \prec)$ : The proof is similar to (iii) above.

(v)  $f$  is an isomorphism from  $(\text{dom}(f), \prec_1)$  to  $(\text{range}(f), \prec_2)$ : Suppose  $x \prec_1 y$  are in  $\text{dom}(f)$ . Put  $a = f(x)$  and  $b = f(y)$ . Using the definition of  $f$ , it follows that  $(\text{pred}(X, \prec_1, x), \prec_1) \cong (\text{pred}(Y, \prec_2, a), \prec_2)$  and  $(\text{pred}(X, \prec_1, y), \prec_1) \cong (\text{pred}(Y, \prec_2, b), \prec_2)$ . As  $x \prec_1 y$ , it follows that  $(\text{pred}(Y, \prec_2, a), \prec_2)$  is isomorphic to an initial segment of  $(\text{pred}(Y, \prec_2, b), \prec_2)$ . Since no well-ordering can be isomorphic to a proper initial-segment of itself, we must have  $a \prec_2 b$ . So  $f$  is an isomorphism from  $(\text{dom}(f), \prec_1)$  to  $(\text{range}(f), \prec_2)$ .

(vi) Either  $\text{dom}(f) = X$  or  $\text{range}(f) = A$ : Suppose not. Let  $x$  be the  $\prec_1$ -least member of  $X \setminus \text{dom}(f)$  and let  $a$  be the  $\prec_2$ -least member of  $Y \setminus \text{range}(f)$ . Since  $\text{range}(f)$  is an initial segment of  $(Y, \prec_2)$ , it follows that  $\text{range}(f) = \text{pred}(Y, \prec_2, a)$ . Similarly,  $\text{dom}(f) = \text{pred}(X, \prec_1, x)$ . But now  $(x, a) \in f$  using (i)-(v) above which is a contradiction.

If  $\text{dom}(f) = X$  and  $\text{range}(f) = Y$ , we get clause (a). If  $\text{dom}(f) \neq X$  and  $\text{range}(f) = Y$ , we get clause (b). If  $\text{dom}(f) = X$  and  $\text{rng}(f) \neq Y$ , we get clause (c).

The uniqueness part follows from the fact that the only isomorphism from a well-ordering to itself is the identity function. □

- (7) Suppose  $X$  is a nonempty set and  $F$  is a choice function on  $\mathcal{P}(X) \setminus \{\emptyset\}$ . Let  $(X, \prec_1)$  and  $(X, \prec_2)$  be two well-orderings satisfying  $(\forall y \in X)(F(X \setminus \text{pred}(X, \prec_k, y))) = y$  for  $k \in \{1, 2\}$ . Show that  $(\forall a, b \in X)(a \prec_1 b \iff a \prec_2 b)$ .

**Solution.** Observe that both  $(X, \prec_1)$  and  $(X, \prec_2)$  are  $F$ -directed well-orderings. So by Claim (2) in the proof of Zermelo's well-ordering theorem (finish that proof), either  $(X, \prec_1) = (X, \prec_2)$  or one of these is a proper initial segment of the other. Since the latter case is impossible, we must have  $\prec_1 = \prec_2$ .  $\square$

- (8) Let  $f : \mathcal{P}(\omega) \setminus \{\emptyset\} \rightarrow \omega$  be defined by  $f(X) = \min(X)$ . Call a well-orderings  $(A, \prec)$   $f$ -directed iff  $A \subseteq \omega$  and for every  $x \in A$ ,

$$f(\omega \setminus \text{pred}(A, \prec, x)) = x$$

Describe all  $f$ -directed well-orderings.

**Solution:** It is clear that each well-ordering in  $\{(\alpha, <) : \alpha \leq \omega\}$  is  $f$ -directed. Let us show that there is no other  $f$ -directed well-ordering. Suppose  $(A, \prec)$  is an  $f$ -directed well-ordering. First suppose that  $A$  is finite (and nonempty) and let  $x_0 \prec x_1 \prec \dots \prec x_n$  list the members of  $A$  where  $n < \omega$ . Then an easy induction on  $k \leq n$  shows that  $x_k = k$ . Next suppose that  $A$  is infinite. Let  $\text{type}(A, \prec) = \alpha$ . So  $\alpha \geq \omega$ . Let  $\langle x_\beta : \beta < \alpha \rangle$  be an order isomorphism from  $\alpha$  to  $(A, \prec)$ . Once again by induction on  $n < \omega$ , we get  $x_n = n$ . Since  $A \subseteq \omega$ , it follows that  $\alpha = \omega$  and hence  $(A, \prec) = (\omega, <)$ .  $\square$

- (9) Show that if  $\alpha < \beta$  are ordinals, then there is an ordinal  $\gamma$  such that  $\alpha + \gamma = \beta$ . (**Hint:**  $\gamma = \text{type}(\beta \setminus \alpha, \in)$ ).

**Solution:** Following the hint, put  $\gamma = \text{type}(\beta \setminus \alpha, \in)$ . Note that

$$(\beta, \in) \cong (\alpha, \in) \oplus (\beta \setminus \alpha, \in)$$

Hence

$$\alpha + \gamma = \alpha + \text{type}((\beta \setminus \alpha, \in)) = \text{type}((\alpha, \in) \oplus (\beta \setminus \alpha, \in)) = \beta$$

where we used the fact that isomorphic well-orderings have the same type.

- (10) Suppose  $\alpha, \beta, \gamma$  are ordinals and  $\alpha + \beta = \alpha + \gamma$ . Show that  $\beta = \gamma$ .

**Solution** Suppose  $\alpha + \gamma_1 = \alpha + \gamma_2 = \beta$ . We'll show that  $\gamma_1 = \gamma_2$ . Suppose not and without loss of generality assume  $\gamma_1 < \gamma_2$ . Then  $\gamma_1 + 1 \leq \gamma_2$  since  $\gamma_1 + 1$  is the smallest ordinal strictly bigger than  $\gamma_1$ . Now

$$\beta = \alpha + \gamma_2 \geq \alpha + (\gamma_1 + 1) = (\alpha + \gamma_1) + 1 > \alpha + \gamma_1 = \beta$$

So  $\beta > \beta$  which is impossible.  $\square$

- (11) Suppose  $\alpha \cdot \alpha = \beta \cdot \beta$ . Show that  $\alpha = \beta$ .

**Solution:** If  $\alpha$  or  $\beta$  is 0, then this is clear. So assume  $\alpha \geq 1$  and  $\beta \geq 1$ . Towards a contradiction, suppose  $\alpha \neq \beta$  and without loss of generality say  $\alpha < \beta$ . Then  $\alpha + 1 \leq \beta$ . Now

$$\beta \cdot \beta \geq \alpha \cdot (\alpha + 1) = (\alpha \cdot \alpha) + \alpha \geq (\alpha \cdot \alpha) + 1 > \alpha \cdot \alpha$$

which contradicts  $\alpha \cdot \alpha = \beta \cdot \beta$ . □

- (12) Show that there is an uncountable chain in  $(\mathcal{P}(\omega), \subseteq)$ . (**Hint:** Identify  $\omega$  with the set of rationals  $\mathbb{Q}$  and for each real number  $x$ , consider  $\{r \in \mathbb{Q} : r < x\}$ ).

**Solution:** Let  $\mathbb{Q}^+$  be the set of positive rational numbers and  $\mathbb{R}^+$  be the set of positive real numbers. Define  $h : \mathbb{Q}^+ \rightarrow \omega$  by

$$h\left(\frac{m}{n}\right) = 2^m 3^n$$

where  $n, m$  are coprime. Note that  $h$  is injective and hence a bijection from  $\mathbb{Q}^+$  to  $\text{range}(h) \subseteq \omega$ . For each  $x \in \mathbb{R}^+$ , let  $A_x = \{r \in \mathbb{Q}^+ : r < x\}$ . Then  $x < y$  implies  $A_x \subsetneq A_y$ . Hence  $\{A_x : x \in \mathbb{R}^+\}$  is an uncountable chain in  $(\mathcal{P}(\mathbb{Q}^+), \subseteq)$ . It follows that  $\{h[A_x] : x \in \mathbb{R}^+\}$  is an uncountable chain in  $(\mathcal{P}(\omega), \subseteq)$ . □

- (13) Suppose  $f : \mathbb{R} \rightarrow \mathbb{R}$  is additive and  $a = f(1)$ .

(a) Show that  $f(0) = 0$ .

(b) Show that for every  $x \in \mathbb{R}$ ,  $f(-x) = -f(x)$ .

(c) Show that for every  $x \in \mathbb{Q}$ ,  $f(x) = ax$ .

**Solution:** (a) Taking  $x = y = 0$ , we get  $f(0 + 0) = f(0) + f(0)$ . So  $f(0) = 0$ .

(b) Taking  $y = -x$ , we get  $f(x + (-x)) = f(x) + f(-x)$ . So  $f(x) + f(-x) = f(0) = 0$ . Hence  $f(-x) = -f(x)$ .

(c) For each  $m, n \geq 1$ ,  $f(m) = f(n(m/n)) = f(m/n + m/n + \cdots + m/n) = nf(m/n)$ . So  $f(m/n) = f(m)/n$ . Next  $f(m) = f(1 + 1 + \cdots + 1) = mf(1) = ma$ . So  $f(m/n) = a(m/n)$ . Also  $f(-m/n) = -f(m/n) = a(-m/n)$ . It follows that for each nonzero  $x \in \mathbb{Q}$ ,  $f(x) = ax$ . Since  $f(0) = 0$ , part (c) follows. □

- (14) Let  $H \subseteq \mathbb{R}$  be a Hamel basis.

(a) Show that every nonzero  $x \in \mathbb{R}$  can be uniquely written as

$$x = a_1x_1 + a_2x_2 + \cdots + a_nx_n$$

where  $x_1 < x_2 < \cdots < x_n$  are in  $H$  and  $a_1, a_2, \dots, a_n$  are nonzero rational numbers. Uniqueness means the following: Suppose

$$x = a_1x_1 + a_2x_2 + \cdots + a_nx_n = b_1y_1 + b_2y_2 + \cdots + a_my_m$$

where  $x_1 < x_2 < \cdots < x_n$  and  $y_1 < y_2 < \cdots < y_m$  are in  $H$  and  $a_1, \dots, a_n, b_1, \dots, b_m$  are nonzero rationals. Show that  $m = n$  and for every  $1 \leq k \leq n$ ,  $x_k = y_k$  and  $a_k = b_k$ .

(b) Let  $f : H \rightarrow \mathbb{R}$ . Show that there is a unique additive function  $g : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f \subseteq g$ .

**Solution:** (a) First let us check uniqueness: Suppose

$$x = a_1x_1 + a_2x_2 + \cdots + a_nx_n = b_1y_1 + b_2y_2 + \cdots + b_my_m$$

where  $x_1 < x_2 < \cdots < x_n$  and  $y_1 < y_2 < \cdots < y_m$  are in  $H$  and  $a_1, \dots, a_n, b_1, \dots, b_m$  are nonzero rationals. We must show that  $m = n$  and for every  $1 \leq k \leq n$ ,  $x_k = y_k$  and  $a_k = b_k$ . Note that

$$(a_1x_1 + a_2x_2 + \cdots + a_nx_n) - (b_1y_1 + b_2y_2 + \cdots + b_my_m) = 0$$

After collecting like terms this boils down to showing the following. If  $w_1 < w_2 < \cdots < w_p$  are in  $H$ ,  $c_1, c_2, \dots, c_p$  are rationals and

$$c_1w_1 + c_2w_2 + \cdots + c_pw_p = 0$$

then  $c_1 = c_2 = \cdots = c_p = 0$ . But this is true because  $H$  is  $\mathbb{Q}$ -linearly independent.

Next suppose  $x \in \mathbb{R}$  is nonzero. We must show that  $x$  is a finite  $\mathbb{Q}$ -linear combination of members of  $H$ . If  $x \in H$ , then  $x = 1 \cdot x$  hence this is clear. So assume  $x \notin H$ . As  $H$  is a maximal  $\mathbb{Q}$ -linearly independent subset of  $\mathbb{R}$ , it follows that  $H \cup \{x\}$  is not  $\mathbb{Q}$ -linearly independent. As  $H$  is  $\mathbb{Q}$ -linearly independent, this means that there are  $x_1 < x_2 < \cdots < x_n$  in  $H$  and nonzero rationals  $a_1, a_2, \dots, a_n, b$  such that

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n + bx = 0$$

Therefore,

$$x = -\frac{a_1}{b}x_1 - \frac{a_2}{b}x_2 - \cdots - \frac{a_n}{b}x_n$$

(b) Define  $g(x)$  as follows. If  $x = a_1x_1 + \cdots + a_nx_n$  where  $x_1 < \cdots < x_n$  are in  $H$  and  $a_1, \dots, a_n$  are rationals, then

$$g(x) = a_1f(x_1) + \cdots + a_nf(x_n)$$

$g$  is well-defined by part (a). That  $g$  is additive is clear from its definition. To see uniqueness suppose  $g' : \mathbb{R} \rightarrow \mathbb{R}$  is another additive extension of  $f$ . Then for every  $r \in \mathbb{Q}$ ,  $g'(rx) = rg'(x)$ . Hence if  $x = a_1x_1 + \cdots + a_nx_n$  where  $x_1 < \cdots < x_n$  are in  $H$  and  $a_1, \dots, a_n$  are rationals, then

$$g'(x) = a_1g'(x_1) + \cdots + a_ng'(x_n) = a_1f(x_1) + \cdots + a_nf(x_n) = g(x)$$

So  $g' = g$ . □

(15) Suppose  $f : \mathbb{R} \rightarrow \mathbb{R}$  satisfies: For every  $x, y \in \mathbb{R}$ ,  $f(x + y) = f(x)f(y)$ .

(a) Show that either  $f$  is identically zero or  $\text{range}(f) \subseteq \mathbb{R}^+$ .

(b) Suppose  $f$  is continuous and not identically zero. Show that  $f(x) = a^x$  for some  $a > 0$ .

**Solution:** (a) Suppose there is some  $a \in \mathbb{R}$  such that  $f(a) = 0$ . Then for every  $x \in \mathbb{R}$ ,  $f(x + a) = f(x)f(a) = f(x) \cdot 0 = 0$ . Hence  $f$  is identically zero. Next suppose  $f(a) \neq 0$  for every  $a \in \mathbb{R}$ . Then  $f(x) = f(x/2 + x/2) = (f(x/2))^2 > 0$ . So either  $f$  is identically zero or  $\text{range}(f) \subseteq \mathbb{R}^+$ .

(b) By part (a),  $\text{range}(f) \subseteq \mathbb{R}^+$  so we can define  $g(x) = \ln(f(x))$ . Then  $g$  is a continuous additive function and hence  $g(x) = bx$  where  $b = g(1)$ . It follows that  $f(x) = e^{g(x)} = e^{bx} = a^x$  where  $a = e^b > 0$ . □

(16) Show that there is a discontinuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x + y) = f(x)f(y)$  for every  $x, y \in \mathbb{R}$ .

**Solution:** Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be any discontinuous additive function and define  $f(x) = e^{g(x)}$ . □

(17) Prove the following.

(a) For every ordinal  $\alpha$ ,  $|\alpha| \leq \alpha$ .

(b) If  $\kappa$  is a cardinal and  $\alpha < \kappa$ , then  $|\alpha| < \kappa$ .

(c) There is an injection from  $X$  to  $Y$  iff  $|X| \leq |Y|$ .

(d) There is a surjection from  $X$  to  $Y$  iff  $|Y| \leq |X|$ .

(e) There is a bijection from  $X$  to  $Y$  iff  $|X| = |Y|$ .

**Solution:** Let us write  $X \preceq Y$  iff there is an injection from  $X$  to  $Y$  and  $X \sim Y$  iff there is a bijection from  $X$  to  $Y$ .

(a) Let  $|\alpha| = \beta$ . Then for every  $\gamma$ , if  $\gamma \sim \alpha$ , then  $\beta \leq \gamma$ . Since  $\alpha \sim \alpha$ , it follows that  $\alpha \leq \beta = |\alpha|$ .

(b) By part (a),  $|\alpha| \leq \alpha < \kappa$ .

(c) Since  $X \sim |X|$  and  $Y \sim |Y|$ , we get  $X \preceq Y$  iff  $|X| \preceq |Y|$ . So it suffices to show that if  $\kappa, \lambda$  are cardinals, then  $\kappa \preceq \lambda$  iff  $\kappa \leq \lambda$ . It is clear that if  $\kappa \leq \lambda$ , then  $\kappa \preceq \lambda$ . Next suppose  $\lambda < \kappa$ . Since  $|\kappa| = \kappa$  and  $\lambda < \kappa$ ,  $\lambda \not\sim \kappa$ . Since  $\lambda \preceq \kappa$ , by the Schröder-Bernstein theorem, it follows that  $\kappa \not\preceq \lambda$ .

(d) By part (c), it suffices to show that for any  $X$  and  $Y$ , there is a surjection from  $X$  to  $Y$  iff  $Y \preceq X$ . We can assume that  $X, Y$  are nonempty. Suppose  $Y \preceq X$ . Fix an injective function  $f : Y \rightarrow X$ . Then  $f : Y \rightarrow \text{range}(f)$  is a bijection. Fix  $y_0 \in Y$ . Define  $g : X \rightarrow Y$  as follows: If  $x \in \text{range}(f)$ , then  $g(x) = f^{-1}(x)$ , otherwise  $g(x) = y_0$ . Clearly,  $\text{range}(g) = Y$ .

Next suppose  $f : X \rightarrow Y$  and  $\text{range}(f) = Y$ . Let  $\mathcal{F} = \{f^{-1}[\{y\}] : y \in Y\}$ . Then  $\mathcal{F}$  is a partition of  $X$  into nonempty sets. Using the axiom of choice let  $h : \mathcal{F} \rightarrow Y$  be a



choice function. Define  $g : Y \rightarrow X$  by  $g(y) = h(f^{-1}[\{y\}])$ . Then  $g : Y \rightarrow X$  is injective.

(e) Use part (c). □

(18) Prove the following.

(a)  $|\mathbb{R}^\omega| = \mathfrak{c}$ .

(b)  $|C(\mathbb{R})| = \mathfrak{c}$  where  $C(\mathbb{R})$  is the set of all continuous functions from  $\mathbb{R}$  to  $\mathbb{R}$ .

(c) Let  $A$  be the set of all real numbers which are roots of some polynomial equation with rational coefficients. Show that  $|A| = \omega$ .

**Solution:** (a) Let us write  $X \sim Y$  iff there is a bijection from  $X$  to  $Y$ . Then it is easy to check that for any set  $A$ ,

$$(A^\omega)^\omega \sim A^{\omega \times \omega} \sim A^\omega$$

Taking  $A = 2 = \{0, 1\}$  and using the fact that  $|\mathbb{R}| = |2^\omega| = \mathfrak{c}$ , we get  $|\mathbb{R}^\omega| = |(2^\omega)^\omega| = |2^\omega| = \mathfrak{c}$ .

(b) It is clear that  $|C(\mathbb{R})| \geq |\mathbb{R}| = \mathfrak{c}$  since every constant function is continuous. To show that  $|C(\mathbb{R})| \leq \mathfrak{c}$ , we'll construct an injective function from  $C(\mathbb{R})$  to  $\mathbb{R}^\omega$ . This suffices since by part (a),  $|\mathbb{R}^\omega| = \mathfrak{c}$ . Since  $|\mathbb{Q}| = \omega$ , it is enough to construct an injective function  $H : C(\mathbb{R}) \rightarrow \mathbb{R}^\mathbb{Q}$  where  $\mathbb{R}^\mathbb{Q}$  is the set of all functions from  $\mathbb{Q}$  to  $\mathbb{R}$ . Given  $f \in C(\mathbb{R})$ , define  $H(f) = f \upharpoonright \mathbb{Q}$ . We claim that  $H$  is injective. To see this assume that  $H(f) = H(g)$  and we'll show that  $f = g$ . Let  $x \in \mathbb{R}$ . Let  $\langle a_n : n < \omega \rangle$  be a sequence of rationals converging to  $x$ . Since  $f, g$  are continuous,  $f(a_n)$  converges to  $f(x)$  and  $g(a_n)$  converges to  $g(x)$ . As  $f \upharpoonright \mathbb{Q} = g \upharpoonright \mathbb{Q}$ , for every  $n < \omega$  we must have  $f(a_n) = g(a_n)$ . Hence  $f(x) = g(x)$ . So  $f = g$  and  $H$  is injective.

(c) For each  $1 \leq n \leq \omega$ , let  $P_n$  be the set of all polynomials of degree  $n$  with rational coefficients. Any polynomial  $f \in P_n$  is of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

where  $a_n, a_{n-1}, \dots, a_0$  are in  $\mathbb{Q}$  and  $a_n \neq 0$ . It follows that  $|P_n| \leq |\mathbb{Q}^n| = \omega$ . Let  $A_n = \{a \in \mathbb{R} : (\exists p \in P_n)(p(a) = 0)\}$ . Since each polynomial in  $P_n$  has  $\leq n$  real roots, it follows that  $A_n$  is a countable union of finite sets. So each  $A_n$  is countable. Finally,  $A = \bigcup \{A_n : 1 \leq n < \omega\}$  is a countable union of countable sets. Hence  $A$  is also countable. As every rational is in  $A$ ,  $|A| \geq \omega$ . Hence  $|A| = \omega$ . □

(19) Suppose  $(P, \preceq)$  is a partial ordering in which every countable chain has an upper bound. Must  $P$  have a maximal element?

**Solution:** No. Let  $P = \{X \subseteq \mathbb{R} : |X| \leq \omega\}$ . Then every countable chain in  $(P, \subseteq)$  has an upper bound in  $P$  (namely its union, since a countable union of countable sets is countable). Let  $X \in P$  be arbitrary. We will show that  $X$  is not maximal in  $(P, \subseteq)$ .

As  $X$  is countable and  $\mathbb{R}$  is uncountable,  $\mathbb{R} \setminus X \neq \emptyset$ . Let  $Y = X \cup \{a\}$  where  $a \in \mathbb{R} \setminus X$ . Then  $Y$  is countable and  $X$  is a proper subset of  $Y$ . So  $X$  is not a  $\subseteq$ -maximal element of  $P$ .  $\square$

- (20) Use Zorn's lemma to show that  $\mathbb{R}^+ = (0, \infty)$  can be partitioned into two nonempty sets that are both closed under addition.

**Solution:** See Problem 6(a) in Chapter 14 of Komjath-Totik book  
<https://doi.org/10.1007/0-387-36219-3>.

- (21) Is there a subset  $X \subseteq \mathbb{R}^2$  that meets every circle at 2 points?

**Solution:** Suppose  $X$  is such a set. First observe that all the points in  $X$  must be collinear otherwise we can draw a circle  $C$  through 3 non-collinear points in  $X$  so that  $|X \cap C| \geq 3$ . Let  $\ell$  be a line such that  $X \subseteq \ell$ . Choose a circle  $C$  such that  $C \cap \ell = \emptyset$ . Then  $C \cap X = \emptyset$  which is a contradiction.  $\square$

- (22) Show that there is a subset  $X \subseteq \mathbb{R}^2$  that meets every circle at 3 points.

**Solution:** Let  $\mathcal{E}$  be the set of all circles in  $\mathbb{R}^2$ . Since each circle is uniquely determined by its center and radius,  $|\mathcal{E}| \leq |\mathbb{R}^2 \times \mathbb{R}^+| = |\mathfrak{c} \times \mathfrak{c}| = \mathfrak{c}$ . Also there are at least  $\mathfrak{c}$  distinct circles so  $|\mathcal{E}| = \mathfrak{c}$ . Let  $\langle C_\alpha : \alpha < \mathfrak{c} \rangle$  be an injective sequence with range  $\mathcal{E}$ . Using transfinite recursion, we are going to construct a  $\subseteq$ -increasing sequence  $\langle S_\alpha : \alpha < \mathfrak{c} \rangle$  of subsets of  $\mathbb{R}^2$  such that the following hold for every  $\alpha < \mathfrak{c}$ .

1.  $S_0 = \emptyset$ .
2.  $|S_\alpha| \leq \max(|\alpha|, \omega) < \mathfrak{c}$ .
3. No 4 points in  $S_\alpha$  are concyclic.
4.  $|S_{\alpha+1} \cap C_\alpha| = 3$ .
5. If  $\alpha$  is a limit ordinal, then  $S_\alpha = \bigcup_{\beta < \alpha} S_\beta$ .

Limit stage: First observe that at any limit stage  $\alpha$ , defining  $S_\alpha = \bigcup_{\beta < \alpha} S_\beta$  does not violate Clause 3. For suppose  $S_\alpha$  does contain 4 concyclic points. Then all of these 4 points must appear at some stage  $\beta < \alpha$  which is impossible.

Successor stage: Having constructed  $S_\alpha$ ,  $S_{\alpha+1}$  is obtained as follows. By Clause 3,  $|S_\alpha \cap C_\alpha| = n \leq 3$ . We would like to add  $n - 3$  points from  $C_\alpha$  to  $S_\alpha$  to get  $S_{\alpha+1}$  while ensuring that no 4 points in  $S_{\alpha+1}$  are concyclic. We have the following cases.

Case (a):  $n = 3$ . Define  $S_{\alpha+1} = S_\alpha$ .

Case (b):  $n = 2$ . Let  $\mathcal{T}$  be the set of circles that pass through 3 points in  $S_\alpha$ . Then  $|\mathcal{T}| \leq |S_\alpha \times S_\alpha \times S_\alpha| \leq \max(|\alpha|, \omega) < \mathfrak{c}$ . Let  $B$  be the set of points of intersection of  $C_\alpha$  with the circles in  $\mathcal{T}$ . Since each circle in  $\mathcal{T}$  can intersect  $C_\alpha$  in at most 2 points, we get  $|B| \leq \max(|\alpha|, \omega) < \mathfrak{c}$ . Choose any point  $p \in C_\alpha \setminus B$  and define  $S_{\alpha+1} = S_\alpha \cup \{p\}$ .

Case (c):  $n = 1$ . First choose  $p \in C_\alpha \setminus B$  and define  $S' = S_\alpha \cup \{p\}$ . Let  $\mathcal{T}'$  be the set of all circles that pass through 3 points in  $S'$ . As before,  $|\mathcal{T}'| < \mathfrak{c}$ . Let  $B'$  be the set of points of intersection of  $C_\alpha$  with the circles in  $\mathcal{T}'$ . Then  $|B'| < \mathfrak{c}$ . Choose  $q \in C_\alpha \setminus (B' \cup \{p\})$  and define  $S_{\alpha+1} = S_\alpha \cup \{p, q\}$ .

Case (d):  $n = 0$ . Proceed as in Case (c).

In each of the Cases (a)-(d), check that  $S_{\alpha+1}$  does not contain 4 concyclic points and  $|S_{\alpha+1} \cap C_\alpha| = 1$ . This concludes the description of  $\langle S_\alpha : \alpha < \mathfrak{c} \rangle$ .

Define  $S = \bigcup_{\alpha < \mathfrak{c}} S_\alpha$ . It should be clear that  $S$  meets every circle at exactly 3 points. □

- (23) Suppose that the set of propositional variables  $\mathcal{V}ar$  is uncountable. Use Zorn's lemma to show the following: Let  $S$  be a set of propositional formulas such that every finite subset of  $S$  is satisfiable. Then  $S$  is satisfiable. **Hint:** Apply Zorn's lemma to  $(\mathcal{F}, \subseteq)$  where  $\mathcal{F}$  is the set of all functions  $h$  such that  $\text{dom}(h) \subseteq \mathcal{V}ar$ ,  $\text{range}(h) \subseteq \{0, 1\}$  and for every finite  $F \subseteq S$ , there exists a valuation  $val : \mathcal{V}ar \rightarrow \{0, 1\}$  such that  $h \subseteq val$  and every formula in  $F$  is true under  $val$ .

**Solution:** Let  $\mathcal{F}$  be the set of all functions  $h$  such that  $\text{dom}(h) \subseteq \mathcal{V}ar$ ,  $\text{range}(h) \subseteq \{0, 1\}$  and for every finite  $F \subseteq S$ , there exists a valuation  $val : \mathcal{V}ar \rightarrow \{0, 1\}$  such that  $h \subseteq val$  and every formula in  $F$  is true under  $val$ .

We claim that every chain in  $(\mathcal{F}, \subseteq)$  has an upper bound. To see this, fix an arbitrary chain  $\mathcal{C} \subseteq \mathcal{F}$  and define  $g = \bigcup \mathcal{C}$ . Since  $\mathcal{C}$  is a chain, it is easy to see that  $g$  is a function. Clearly,  $\text{dom}(g) \subseteq \mathcal{V}ar$  and  $\text{range}(g) \subseteq \{0, 1\}$ . So it would be sufficient to show that  $g \in \mathcal{F}$  since then  $g$  is an upper bound of  $\mathcal{C}$  in  $(\mathcal{F}, \subseteq)$ . Towards a contradiction, suppose  $g \notin \mathcal{F}$ . Fix a finite  $F \subseteq S$  such that there is no valuation  $val : \mathcal{V}ar \rightarrow \{0, 1\}$  satisfying:  $g \subseteq val$  and every formula in  $F$  is true under  $val$ . Choose a finite  $V \subseteq \mathcal{V}ar$  that contains every propositional variable that occurs in a formula in  $F$ . Put  $W = V \cap \text{dom}(g)$ . Since  $\mathcal{C}$  is a chain, we can find an  $h \in \mathcal{C}$  such that  $W \subseteq \text{dom}(h)$ . Since  $h \in \mathcal{F}$ , there exists a valuation  $val' : \mathcal{V}ar \rightarrow \{0, 1\}$  such that  $h \subseteq val'$  and every formula in  $F$  is true under  $val'$ . Define another valuation  $val : \mathcal{V}ar \rightarrow \{0, 1\}$  as follows:

$$val(p) = \begin{cases} g(p) & \text{if } p \in \text{dom}(g) \\ val'(p) & \text{otherwise} \end{cases}$$

Observe that  $val$  and  $val'$  agree on every propositional variable in  $V$ . Hence every formula in  $F$  is true under  $val$ . But  $g \subseteq val$  so we have a contradiction. So  $g \in \mathcal{F}$  is an upper bound of  $\mathcal{C}$ .

Using Zorn's lemma, fix a  $\subseteq$ -maximal  $f$  in  $\mathcal{F}$ . We claim that  $\text{dom}(f) = \mathcal{V}ar$ . This will complete the proof since it implies that  $f$  is a valuation under which every formula in  $\mathcal{F}$  is true. Towards a contradiction, assume some propositional variable  $p \notin \text{dom}(f)$ . Define  $f_0 = f \cup \{(p, 0)\}$  and  $f_1 = \{(p, 1)\}$ . By the Lemma on Lecture

slide no. 89, it follows that one of  $f_0, f_1$  is in  $\mathcal{F}$ . But this contradicts the maximality of  $g$ . Hence  $\text{dom}(f) = \text{Var}$  and the proof is complete.  $\square$

- (24) Let  $\phi$  be a propositional formula in which  $\neg$  does not occur. Show that  $\phi$  is satisfiable.

**Solution.** Define  $\text{val}(p) = 1$  for every propositional variable  $p$ . By induction on the length of formula  $\phi$ , show that  $\phi$  is true under  $\text{val}$ .  $\square$

- (25) Show that the axiom of extensionality is not a logically valid  $\mathcal{L}_{ZFC}$ -sentence.

**Solution:** Recall that the axiom of extensionality is the following  $\mathcal{L}_{ZFC}$ -sentence

$$\psi \equiv (\forall x)(\forall y)[(\forall v)(v \in x \iff v \in y) \implies (x = y)].$$

Let  $\mathcal{M} = (M, \in^{\mathcal{M}})$  where  $M = \{0, 1, 2\}$  and  $\in^{\mathcal{M}} = \{(0, 1), (0, 2)\}$ . Now  $(\forall v \in M)(v \in^{\mathcal{M}} 1 \iff v \in^{\mathcal{M}} 2) \wedge 1 \neq 2$ . So  $\mathcal{M} \models \neg\psi$ . Since we have shown that there is an  $\mathcal{L}_{ZFC}$ -structure (namely  $\mathcal{M}$ ) in which  $\psi$  is false, it follows that  $\psi$  is not logically valid.  $\square$

- (26) Show that  $(\forall x)(\forall y)(x + y = y + x)$  is not a logically valid  $\mathcal{L}_{PA}$ -sentence.

**Solution:** Consider  $\mathcal{M} = (M, +^{\mathcal{M}})$  where  $M = \mathbb{R}$  and  $x +^{\mathcal{M}} y = x - y$ .  $\square$

- (27) Let  $\mathcal{L}$  be a first order language and  $T$  be an  $\mathcal{L}$ -theory. Assume  $T \vdash \phi$  and  $T \vdash (\phi \implies \psi)$ . Show that  $T \vdash \psi$ .

**Solution:** Let  $\phi_1, \dots, \phi$  be a proof of  $\phi$  in  $T$  and  $\psi_1, \dots, (\phi \implies \psi)$  be a proof of  $(\phi \implies \psi)$  in  $T$ . Check that

$$\phi_1, \dots, \phi, \psi_1, \dots, (\phi \implies \psi), \psi$$

is a proof of  $\psi$  in  $T$ .  $\square$

- (28) Let  $\mathcal{L}$  be a first order language and  $T$  be an  $\mathcal{L}$ -theory. Assume  $T \vdash \phi_1$  and  $T \vdash \phi_2$ . Show that  $T \vdash (\phi_1 \wedge \phi_2)$ .

**Solution:** Use the fact that  $\phi_1 \implies (\phi_2 \implies (\phi_1 \wedge \phi_2))$  is a propositional tautology.  $\square$

- (29) Let  $\mathcal{L}$  be a first order language. Call a set  $S$  of  $\mathcal{L}$ -sentences deductively closed if  $S$  contains every logical axiom of  $\mathcal{L}$  and whenever  $\phi$  and  $(\phi \implies \psi)$  are in  $S$ ,  $\psi$  is also in  $S$ . Show that for any  $\mathcal{L}$ -theory  $T$ , the set of theorems of  $T$  is the smallest set of  $\mathcal{L}$ -sentences that contains  $T$  and is deductively closed.

**Solution:** Let  $S$  be the set of all  $\mathcal{L}$ -sentences  $\psi$  such that  $T \vdash \psi$ . So  $S$  is the set of all theorems of  $T$ .

We first show that  $S$  is deductively closed. If  $\psi$  is a logical axiom of  $\mathcal{L}$ , then  $T \vdash \psi$ , so we get  $\psi \in S$ . Next assume  $\phi$  and  $(\phi \implies \psi)$  are both in  $S$ . We need to show that  $\psi \in S$ . But this follows from problem (27). Thus  $S$  is deductively closed.

Next, to show that  $S$  is the smallest deductively closed set of  $\mathcal{L}$ -sentences containing  $T$ , suppose  $S'$  is another deductively closed set of  $\mathcal{L}$ -sentences such that  $T \subseteq S'$ . We will show that  $S \subseteq S'$ . Let  $\psi \in S$ . Then  $T \vdash \psi$ . Fix a proof  $\psi_1, \dots, \psi_n \equiv \psi$  of  $\psi$  in  $T$ . By induction on  $k \leq n$ , we will show that each  $\psi_k \in S'$ . It will follow that  $\psi \in S'$  and hence  $S \subseteq S'$ .

If  $k = 1$ , it must be the case that either  $\psi_1 \in T$  or  $\psi_1$  is a logical axiom of  $\mathcal{L}$ . If  $\psi_1 \in T$ , then  $\psi_1 \in S'$  since  $T \subseteq S'$ . If  $\psi_1$  is a logical axiom, then as  $S'$  is deductively closed, we must have  $\psi_1 \in S'$ . So in both cases  $\psi_1 \in S'$ .

For the inductive step, assume  $\psi_j \in S$  for each  $j \leq k < n$ . We will show that  $\psi_{k+1} \in S'$ . First suppose that either  $\psi_{k+1}$  is a logical axiom of  $\mathcal{L}$  or  $\psi_{k+1} \in T$ . Then  $\psi_{k+1} \in S'$  as argued above. Finally suppose  $\psi_{k+1}$  was deduced from some  $\psi_i$  and  $\psi_j \equiv (\psi_i \implies \psi_{k+1})$  (where  $i, j \leq k$ ) via modus ponens. Then using the inductive hypothesis we get that  $\psi_i, \psi_j \in S'$  and by using the fact that  $S'$  is deductively closed, we can conclude that  $\psi_{k+1} \in S'$ . This completes the inductive step.  $\square$

- (30) Describe all substructures of  $(\mathbb{Z}, <)$  and  $(\mathbb{Z}, +)$ .

**Solution:** A substructure of  $(\mathbb{Z}, <)$  is of the form  $\mathcal{M} = (M, <)$  where  $M \subseteq \mathbb{Z}$  is any nonempty subset and  $<$  is the usual order on  $M$ . A substructure of  $(\mathbb{Z}, +)$  is of the form  $\mathcal{M} = (M, +)$  where  $M \subseteq \mathbb{Z}$  is nonempty and closed under addition. We leave it as an exercise to describe such subsets of  $\mathbb{Z}$ .  $\square$

- (31) Describe all elementary submodels of  $(\mathbb{Z}, <)$  and  $(\mathbb{Z}, +)$ .

**Solution:** The only elementary submodel of  $(\mathbb{Z}, <)$  is  $(\mathbb{Z}, <)$ . To see this, suppose  $(M, <)$  is an elementary submodel of  $(\mathbb{Z}, <)$  and towards a contradiction, assume  $M \neq \mathbb{Z}$ . Fix some  $n \in \mathbb{Z} \setminus M$ . Since

$$(\mathbb{Z}, <) \models (\forall x)(\exists y)(y < x) \wedge (\forall x)(\exists y)(x < y)$$

we must have

$$(M, <) \models (\forall x)(\exists y)(y < x) \wedge (\forall x)(\exists y)(x < y)$$

So  $(M, <)$  does not have a largest or smallest integer. Using this fact, we can find  $a, b \in M$  such that  $a$  is the largest integer in  $M$  strictly below  $n$  and  $b$  is the smallest integer in  $M$  strictly above  $n$ . Observe that there is no integer in  $M$  that is strictly between  $a$  and  $b$ . So  $(M, <) \models \neg(\exists x)(a < x \wedge x < b)$ . As  $(M, <)$  is an elementary submodel of  $(\mathbb{Z}, <)$ , we get  $(\mathbb{Z}, <) \models \neg(\exists x)(a < x \wedge x < b)$  which is false since  $x = n$  is strictly between  $a$  and  $b$ . This contradiction shows that  $M = \mathbb{Z}$ .

Next, we will show that  $(\mathbb{Z}, +)$  is the only elementary submodel of  $(\mathbb{Z}, +)$ . Suppose  $(M, +)$  is an elementary submodel of  $(\mathbb{Z}, +)$ . Then  $M$  is closed under  $+$ . Moreover, since  $(\mathbb{Z}, +) \models (\forall x)(\forall y)(\exists z)(x = y + z)$ , it follows that  $M$  is also closed under  $-$ . We will show that  $1 \in M$ . Since  $M$  is closed under both  $+$  and  $-$ , it will follow that  $M = \mathbb{Z}$ .

First note that since  $\mathbb{Z}$  is infinite,  $M$  must also be infinite (use  $(\mathbb{Z}, <) \models \exists_{\geq n}$  for every  $n \geq 1$ , Slide 126). So  $M$  must have both positive and negative integers. Let  $p$  be any positive integer in  $M$ . If  $p = 1$ , then we are done. So suppose  $p > 1$ . Observe that  $(\mathbb{Z}, +) \models (\exists x)(px = p)$  where  $px$  is the term  $(x + x + \cdots + x)$  ( $p$  times). Hence  $(M, +) \models (\exists x)(px = p)$ . Therefore  $1 \in M$  and we are done.  $\square$

- (32) For each  $n \geq 1$  and  $\mathcal{L} = \emptyset$ . Give an example of a consistent  $\mathcal{L}$ -theory  $T$  such that for every model  $\mathcal{M} \models T$ ,  $|M| = n$ .

**Solution:** Let  $T = \{\exists_{\geq n}, \neg(\exists_{n+1})\}$  (Slide 126).  $\square$

- (33) Let  $\mathcal{L} = \mathcal{L}_{PA} \cup \{c\}$  where  $c$  is a new constant symbol. Let  $\text{Primes} = \{2, 3, 5, 7, \dots\}$  be the set of all primes numbers. For each  $p \in \text{Primes}$ , let “ $p$  divides  $c$ ” denote the  $\mathcal{L}$ -sentence  $(\exists y)(S^p(0) \cdot y = c)$ . For each  $X \subseteq \text{Primes}$ , let  $T_X$  be the  $\mathcal{L}$ -theory

$$T_X = TA \cup \{(p \text{ divides } c) : p \in X\} \cup \{\neg(p \text{ divides } c) : p \in \text{Primes} \setminus X\}$$

where  $TA = Th(\omega, 0, S, +, \cdot)$  denotes true arithmetic.

(a) Show that  $T_X$  is consistent for every  $X \subseteq \text{Primes}$ .

(b) Show that TA has continuum many pairwise non-isomorphic countable models.

**Solution:** (a) We will show that every finite subset of  $T_X$  has a model. This suffices since then, by compactness theorem, it will follow that  $T_X$  has a model and therefore  $T_X$  is consistent.

Let  $F$  be a finite subset of  $T_X$ . We will construct a model of  $F$ . Let  $W$  be the set of all primes  $p$  such that  $(p \text{ divides } c) \in F$ . Note that  $W$  is a finite subset of  $X$ . Let  $\mathcal{M} = (\omega, 0, S, +, \cdot, c^{\mathcal{M}})$  where  $(\omega, 0, S, +, \cdot)$  is the standard model of arithmetic and  $c^{\mathcal{M}}$  is the product of all the primes in  $W$  (If  $W = \emptyset$ , then define  $c^{\mathcal{M}} = 1$ ). Then a prime  $p$  divides  $c^{\mathcal{M}}$  iff  $p \in W$ . It follows that  $\mathcal{M} \models F$ .

(b) Using part (a), we can fix a family  $\{\mathcal{M}'_X : X \subseteq \text{Primes}\}$  such that for each  $X \subseteq \text{Primes}$ ,  $\mathcal{M}'_X = (M_X, 0^{\mathcal{M}_X}, S^{\mathcal{M}_X}, +^{\mathcal{M}_X}, \cdot^{\mathcal{M}_X}, c^{\mathcal{M}_X})$  is a countable  $\mathcal{L}$ -structure such that  $\mathcal{M}'_X \models T_X$ . Let  $\mathcal{M}_X = (M_X, 0^{\mathcal{M}_X}, S^{\mathcal{M}_X}, +^{\mathcal{M}_X}, \cdot^{\mathcal{M}_X})$ . Then  $\mathcal{M}_X$  is an  $\mathcal{L}_{PA}$ -structure such that  $\mathcal{M}_X \models TA$ .

We claim that for any  $X \subseteq \text{Primes}$ ,  $\{Y \subseteq \text{Primes} : \mathcal{M}_X \cong \mathcal{M}_Y\}$  is countable. Since  $|\{X : X \subseteq \text{Primes}\}| = \mathfrak{c}$ , it will follow that there are continuum many pairwise non-isomorphic models of TA in  $\{\mathcal{M}_X : X \subseteq \text{Primes}\}$ .

Let  $X \subseteq \text{Primes}$ . For each prime  $p$ , let  $\phi_p$  denote the formula “ $p$  divides  $x$ ” where  $x$  is a variable. For each  $a \in M_X$ , let  $T_a = \{p \in \text{Primes} : \mathcal{M}_X \models \phi_p(a/x)\}$ . Then  $T_X = \{T_a : a \in M_X\}$  is a countable family of subsets of  $\text{Primes}$ .

Now observe that if  $Y \subseteq \text{Primes}$  and  $Y \notin T_X$ , then  $\mathcal{M}_X$  cannot be isomorphic to  $\mathcal{M}_Y$ . This is because there exists a member  $a \in M_Y$  (namely,  $a = c^{\mathcal{M}_Y}$ ) such that  $Y = \{p \in \text{Primes} : \mathcal{M}_Y \models \phi_p(a/x)\}$  while there is no such member in  $M_X$ . So  $\{Y \subseteq \text{Primes} : \mathcal{M}_X \cong \mathcal{M}_Y\}$  is countable and we are done.  $\square$

- (34) Show that  $(\mathbb{Q}, <)$  is an elementary submodel of  $(\mathbb{R}, <)$ .

**Solution:** Put  $\mathcal{L} = \{<\}$ . By the Tarski-Vaught criterion, it suffices to show that for every  $\mathcal{L}$ -formula  $\phi(x, y_1, \dots, y_n)$  and  $a_1, \dots, a_n \in \mathbb{Q}$ , if there exists  $b \in \mathbb{R}$  such that  $(\mathbb{R}, <) \models \phi(b, a_1, \dots, a_n)$ , then there exists  $c \in \mathbb{Q}$  such that  $(\mathbb{R}, <) \models \phi(c, a_1, \dots, a_n)$ . Fix some  $b \in \mathbb{R}$  such that  $(\mathbb{R}, <) \models \phi(b, a_1, \dots, a_n)$ . If  $b \in \mathbb{Q}$ , then we are done. So we can assume that  $b \in \mathbb{R} \setminus \mathbb{Q}$ . In particular,  $b \notin \{a_1, \dots, a_n\}$ . Using the fact that  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , choose  $c \in \mathbb{Q} \setminus \{a_1, \dots, a_n\}$  such that for every  $1 \leq k \leq n$ ,  $(a_n < c) \iff (a_n < b)$ . Now check that there is a (piecewise linear) order-preserving bijection  $h : \mathbb{R} \rightarrow \mathbb{R}$  such that  $h(a_k) = a_k$  for every  $1 \leq k \leq n$  and  $h(b) = c$  (Draw a picture on the number line and use the fact that for any two bounded closed intervals in  $\mathbb{R}$ , there is an order-preserving linear function between them). Since  $h$  is an isomorphism from  $(\mathbb{R}, <) \rightarrow (\mathbb{R}, <)$ , by the Lemma on Slide 125, it follows that  $(\mathbb{R}, <) \models \phi(h(b), h(a_1), \dots, h(a_n))$ . As  $h(b) = c$  and  $h(a_k) = a_k$  for every  $1 \leq k \leq n$ , we get  $(\mathbb{R}, <) \models \phi(c, a_1, \dots, a_n)$ .  $\square$

- (35) Show that every countable linear ordering  $(L, <_L)$  is isomorphic to  $(A, <)$  for some  $A \subseteq \mathbb{Q}$ .

**Solution:** If  $L$  is finite, this is clear – Say  $L = \{x_0 <_L x_1 <_L x_2 <_L \dots <_L x_k\}$ . Take  $A = \{0, 1, 2, \dots, k\}$ . Then  $f(x_i) = i$  is an order isomorphism from  $(L, <_L)$  to  $(A, <)$ .

So assume  $|L| = \omega$  and fix an injective enumeration  $\langle x_k : k < \omega \rangle$  of  $L$ . By induction on  $n < \omega$ , define  $f_n : \{x_k : k \leq n\} \rightarrow \mathbb{Q}$  as follows. Start by defining  $f_0(x_0) = 0$ . Suppose  $f_n : \{x_k : k \leq n\} \rightarrow \mathbb{Q}$  has been defined such that for every  $i < j \leq n$ ,  $x_i <_L x_j \iff f(x_i) < f(x_j)$ .

Define  $f_{n+1} : \{x_k : k \leq n+1\} \rightarrow \mathbb{Q}$  as follows. Define  $f_{n+1}(x_k) = f_n(x_k)$  for every  $k \leq n$ . Next, define  $f_{n+1}(x_{n+1}) = r \in \mathbb{Q}$  that satisfies: For every  $i \leq n$ ,  $x_i <_L x_{n+1} \iff f_n(x_i) < r$ . This is always possible because  $(\mathbb{Q}, <) \models DLO$ .

Having defined  $f_n$  for every  $n < \omega$ , put  $f = \bigcup_{n < \omega} f_n$  and note that  $f : L \rightarrow \mathbb{Q}$  is an order preserving function. Put  $A = \text{range}(f)$ . Then  $(L, <_L) \cong (A, <)$ .  $\square$

- (36) Let  $\mathcal{L} = \{<\}$  where  $<$  is a binary relation symbol. Let  $T$  be the  $\mathcal{L}$ -theory obtained by replacing Axiom (6) of DLO (Slide 142) with the following

$$(\exists y)(\forall x)(y < x \vee y = x)$$

Show that  $T$  is  $\omega$ -categorical and therefore complete. Use this to also show that for every  $\mathcal{L}$ -sentence  $\phi$

$$T \vdash \phi \iff (\mathbb{Q}_{\geq 0}, <) \models \phi$$

where  $\mathbb{Q}_{\geq 0}$  is the set of all non-negative rationals.

**Solution:** Let  $(L_1, <_1)$  and  $(L_2, <_2)$  be two models of  $T$  such that  $|L_1| = |L_2| = \omega$ . Then both  $L_1, L_2$  have a least member since the sentence  $(\exists y)(\forall x)(y < x \vee y = x)$  is true in both of them. Let  $a_1$  be the  $<_1$ -least member of  $L_1$  and  $a_2$  be the  $<_2$ -least member of  $L_2$ . Put  $L'_1 = L_1 \setminus \{a_1\}$  and  $L'_2 = L_2 \setminus \{a_2\}$ . Then  $(L'_1, <_1) \models DLO$  and

$(L'_2, <_2) \models DLO$ . As DLO is  $\omega$ -categorical, it follows that  $(L'_1, <_1) \cong (L'_2, <_2)$ . Let  $f : L'_1 \rightarrow L'_2$  be an isomorphism. Extend  $f$  to  $L_1$  by defining  $f(a_1) = a_2$ . It is easy to see that  $f$  is an isomorphism from  $L_1$  to  $L_2$ .

Note that  $T$  does not have any finite model since any linear ordering that models  $T$  does not have a largest member. So by the theorem on slide 141, as  $T$  is  $\omega$ -categorical, it must be complete. Since  $(\mathbb{Q}_{\geq 0}, <) \models T$ , by soundness theorem, it follows that if  $T \vdash \phi$ , then  $(\mathbb{Q}_{\geq 0}, <) \models \phi$ . Finally, suppose  $(\mathbb{Q}_{\geq 0}, <) \models \phi$  and towards a contradiction, assume  $T \not\vdash \phi$ . Then since  $T$  is complete, we must have  $T \vdash \neg\phi$  and so  $(\mathbb{Q}_{\geq 0}, <) \models \neg\phi$  which is impossible.  $\square$

- (37) Let  $W \subseteq \omega$  be nonempty. Show that  $W$  is c.e. iff there exists a computable function  $f : \omega \rightarrow \omega$  such that  $\text{range}(f) = W$ .

**Solution:** First assume that  $W$  is c.e. Fix a program  $P$  such that for each  $n < \omega$ ,  $P$  halts on input  $n$  iff  $n \in W$ .

Define a program  $Q$  as follows. On input  $n$ ,  $Q$  runs  $P$  on each one of the inputs  $0, 1, \dots, n$  for  $n$  steps. Let  $S_n$  be the set of those  $k \leq n$  such that  $P$  halts on input  $k$  in at most  $n$  steps. Let  $W_n$  be the set of outputs of  $Q$  on inputs  $0, 1, \dots, n-1$ . If  $S_n \setminus W_n \neq \emptyset$ , then  $Q$  outputs  $\min(S_n \setminus W_n)$ . Otherwise,  $Q$  outputs  $\min(W)$ .

It is clear that  $Q$  halts on every input. Let  $f : \omega \rightarrow \omega$  be the function computed by  $Q$ . We claim that  $\text{range}(f) = W$ . That  $\text{range}(f) \subseteq W$  is obvious. For the other inclusion, towards a contradiction, suppose  $W \setminus \text{range}(f) \neq \emptyset$  and let  $n_* = \min(W \setminus \text{range}(f))$ . Choose  $m > n_*$  large enough such that  $W \cap n_* \subseteq \text{range}(f \upharpoonright m)$  and for every  $n \leq n_*$ , if  $n \in W$ , then  $P$  halts on input  $n$  in less than  $m$  steps. Now observe that on input  $m$ ,  $Q$  must output  $n_*$ : A contradiction. So we must have  $W \subseteq \text{range}(f)$ . It follows that  $W = \text{range}(f)$ .

Next assume that  $f : \omega \rightarrow \omega$  is computable. Put  $W = \text{range}(f)$ . Let  $P$  be a program that on input  $n$  starts computing  $f(0), f(1), f(2), \dots$  and halts iff  $n$  appears in this list. Then  $P$  witnesses that  $W$  is c.e.  $\square$

- (38) Let  $W \subseteq \omega$  be infinite. Show that  $W$  is c.e. iff there exists an injective computable function  $f : \omega \rightarrow \omega$  such that  $\text{range}(f) = W$ .

**Solution:** Assume  $W \subseteq \omega$  is c.e. and infinite. By the previous problem, we can fix a computable  $g : \omega \rightarrow \omega$  such that  $\text{range}(g) = W$ . Define  $f : \omega \rightarrow \omega$  as follows.  $f(0) = g(0)$  and  $f(n+1)$  is the first number that appears in the list  $g(0), g(1), g(2), \dots$ , that is not in  $\{f(0), f(1), \dots, f(n)\}$  (such a number exists since  $\text{range}(g)$  is infinite). It is clear that  $f$  is computable, injective and  $\text{range}(f) = \text{range}(g) = W$ . The other implication follows from the previous problem.  $\square$

- (39) Let  $W \subseteq \omega$  be an infinite c.e. set. Show that  $W$  has an infinite computable subset.

**Solution:** By the previous problem, we can fix an injective computable  $f : \omega \rightarrow \omega$  such that  $\text{range}(f) = W$ . Define  $g : \omega \rightarrow \omega$  as follows.  $g(0) = f(0)$  and  $g(n+1)$  is the



first entry in the list  $f(0), f(1), f(2), \dots$ , that is strictly bigger than  $g(n)$ . Then  $g : \omega \rightarrow \omega$  is computable and strictly increasing. Put  $\text{range}(g) = A$ . Then  $A \subseteq W$  is infinite. To see that  $A$  is computable, note that to check whether  $n \in A = \text{range}(g)$ , we only need to check if  $n \in \{g(0), g(1), \dots, g(n)\}$  (because  $g$  is strictly increasing).  $\square$

- (40) Let  $W \subseteq \omega$  be nonempty. Show that  $W$  is c.e. iff there exists a computable  $A \subseteq \omega^2$  such that  $W = \{n \in \omega : (\exists m)((n, m) \in A)\}$ .

**Solution:** First assume  $W$  is c.e. and fix a program  $P$  that halts on input  $n$  iff  $n \in W$ . Define  $A$  to be the set of  $(n, m) \in \omega^2$  such that on input  $n$ ,  $P$  halts in  $\leq m$  steps. Then  $A$  is computable and  $W = \{n \in \omega : (\exists m)((n, m) \in A)\}$ .

Next suppose  $A \subseteq \omega^2$  is computable and  $W = \{n \in \omega : (\exists m)((n, m) \in A)\}$ . Consider a program  $Q$  that on input  $n$ , starts checking if  $(n, m) \in A$  for  $m = 0, 1, 2, 3, \dots$  and halts as soon as it finds some  $m$  for which  $(n, m) \in A$ . Then  $Q$  halts on input  $n$  iff  $(\exists m)((n, m) \in A)$  iff  $n \in W$ . So  $W$  is c.e.  $\square$

- (41) Suppose  $X \subseteq \omega$  is numeralwise representable in PA. Show that  $X$  is computable.

**Solution:** Fix an  $\mathcal{L}_{PA}$  formula  $\phi(x)$  such that for every  $n < \omega$ , if  $n \in A$ , then  $PA \vdash \phi(\bar{n})$  and if  $n \notin A$ , then  $PA \vdash \neg\phi(\bar{n})$ . Since the set of theorems in PA is c.e. (see Slides 159-160), we can fix a program  $P$  such that for any  $\mathcal{L}_{PA}$ -sentence  $\psi$ ,  $P$  halts on input  $\psi$  iff  $PA \vdash \psi$ . Consider the program  $Q$  which on input  $n$ , runs  $P$  with inputs  $\phi(\bar{n})$  and  $\neg\phi(\bar{n})$ . If  $P$  halts on input  $\phi(\bar{n})$ , then  $Q$  returns 1. If  $P$  halts on input  $\neg\phi(\bar{n})$ , then  $Q$  returns 0. It is easy to see that  $Q$  computes  $X$ .  $\square$

- (42) Let  $E \subseteq \omega$  be a non-computable c.e. set. Show that  $E$  is definable in  $\mathcal{N} = (\omega, 0, S, +, \cdot)$  but not numeralwise representable in PA.

**Solution:** By problem (40), we can fix a computable  $A \subseteq \omega^2$  such that  $E = \{n : (\exists m)((n, m) \in A)\}$ . Since  $A$  is computable, it is definable in  $\mathcal{N}$ . So there is an  $\mathcal{L}_{PA}$ -formula  $\phi(y, x)$  such that for every  $(n, m) \in \omega^2$ ,  $(n, m) \in A$  iff  $\mathcal{N} \models \phi(n, m)$ . Let  $\psi(y)$  be the formula  $(\exists x)(\phi(y, x))$ . Then for every  $n < \omega$ ,  $n \in E$  iff  $(\exists m)((n, m) \in A)$  iff  $\mathcal{N} \models \psi(n)$ . Hence  $E$  is definable in  $\mathcal{N}$  via  $\psi(y)$ . That  $E$  is not numeralwise representable in PA follows from problem (41) and the fact that  $E$  is non-computable.  $\square$

- (43) Do the Exercise on the last slide (180) of notes.

**Solution:** Let  $m < \omega$ . We must show that if  $m \in H$ , then  $Q$  returns 1 on input  $m$  and if  $m \notin H$ , then  $Q$  returns 0 on input  $m$ .

First suppose  $m \in H$ . Then for some  $n < \omega$ ,  $f(n) = m$ . By Clause 1,  $PA \vdash \psi(\bar{m}, \bar{n})$ . Note that  $\psi(\bar{m}, \bar{n}) \implies (\exists x)(\psi(\bar{m}, x))$  is a logical axiom of type 5. So by Modus Ponens,  $PA \vdash (\exists x)(\psi(\bar{m}, x))$ . Hence  $Q$  returns 1 on input  $m$ .

Next suppose  $m \notin H$ . We must show that  $PA \not\vdash (\exists x)(\psi(\bar{m}, x))$ . Towards a contradiction, suppose  $PA \vdash (\exists x)(\psi(\bar{m}, x))$ . Since  $\mathcal{N}$  is a model of PA, it follows that

$\mathcal{N} \models (\exists x)(\psi(m, x))$ . Fix  $n < \omega$  such that  $\mathcal{N} \models \psi(m, n)$ . Since  $m \notin H = \text{range}(f)$ , we must have  $f(n) \neq m$ . By Clause 2, this implies that  $PA \vdash \neg\psi(\overline{m}, \overline{n})$ . As  $\mathcal{N}$  models PA, we get  $\mathcal{N} \models \neg\psi(m, n)$ . So  $\mathcal{N} \models \psi(m, n)$  and  $\mathcal{N} \models \neg\psi(m, n)$ : A contradiction.

It follows that  $Q$  computes  $H$ . □