Representation Theory — Examples Sheet 2

On this sheet all representations are complex representations

- 1. Let (ρ, V) be a representation of a finite group G with character χ . Show that $\ker \rho = \{g \in G \mid \chi(g) = \chi(1)\}$. Show further that $|\chi(g)| \leq \chi(1)$ for all $g \in G$, with equality precisely if $\rho(g) = \lambda \operatorname{id}_V$ for some $\lambda \in \mathbb{C}^{\times}$. Explain how the set of normal subgroups of G may be calculated directly from the character table.
- 2. Let χ be the character of a representation of a group G and let $g \in G$. If g has order 2 show that $\chi(g) \in \mathbb{Z}$ and that $\chi(g) \equiv \chi(1) \mod 2$. Show that if in addition G is a non-cyclic simple group then $\chi(g) \equiv \chi(1) \mod 4$. If instead g has order 3 and is conjugate to g^2 show that $\chi(g) \equiv \chi(1) \mod 3$.
- 3. Construct the character tables of the dihedral group D_8 and the quaternion group Q_8 . What do you notice? Compare the determinants of their respective two dimensional representations.
- 4. Construct the character tables of the dihedral groups D_{10} and D_{12} . How do the irreducible representations decompose when restricted to the subgroups of rotations?
- 5. Construct the character tables of A_4 , S_4 , A_5 and S_5 . The action of S_n on A_n by conjugation induces an action on the character table of A_n by permuting the conjugacy classes. Describe what this does to the rows of the character table for n = 4, 5.
- 6. Show that there is only one non-abelian group of order 21 up to isomorphism. Construct its character table.
- 7. A group of order 720 has 11 conjugacy classes. Two representations of the group are known and have corresponding characters α and β . The table below summarises the sizes of the conjugacy classes and the values of α and β on them. Prove that the group has an irreducible representation of degree 16 and calculate its character.

8. A group of order 168 has 6 conjugacy classes. Three representations of this group are known and have characters α, β and γ summarised in the table below. Construct the character table of the group. You may assume if required that $\sqrt{7}$ is not in the field generated by \mathbb{Q} and a primitive 7^{th} root of unity.

- 9. Consider the action of a finite group G by conjugation. What is the character of the corresponding permutation representation $\mathbb{C}G$? Prove that the sum of elements in any row of the character table of G is a non-negative integer.
- 10. Show that the character table of a finite group G is invertible when viewed as a matrix. By considering the actions induced on the rows and on the columns of the character table by complex conjugation, show that the number of irreducible characters of G that only take real values is the number of self-inverse conjugacy classes.
- 11. Let G be a finite group and χ be an irreducible character of G. By beginning with the irreducible representations, show that if (ρ, V) is any representation of G then $\frac{\chi(1)}{|G|} \sum_{g \in G} \overline{\chi(g)} \rho(g)$ is a G-linear projection onto a subspace of V. Deduce that every representation can be decomposed uniquely into isotypical components.

Comments and Corrections to S.J.Wadsley@dpmms.cam.ac.uk.