

ElegantNote: An Elegant L^AT_EX Template for Notes

FirstName LASTNAME

Example University

Version: 2.60

Update: June 5, 2026

1 ElegantNote Instructions

The ElegantNote template is based on the L^AT_EX `article` class. It supports both pdfL^AT_EX (recommended for English) and X_YL^AT_EX.

Key features:

- Two modes: geye (green background) and hazy (light blue)
- Five color themes: blue, green, cyan, sakura, black
- Customisable font sizes: 8pt – 20pt
- Theorem environments: theorem, lemma, proposition, corollary, definition, conjecture, example, remark, note

Note Select `device=normal` for a standard A4 PDF output.

2 Writing Sample

We define the integral of a measurable function in three steps.

Definition 2.1 (Left Coset) Let H be a subgroup of a group G . A left coset of H in G is a subset of G of the form $xH = \{xh : h \in H\}$.

Lemma 2.1 (Size of Left Coset) Let H be a finite subgroup of a group G . Then each left coset of H in G has the same number of elements as H .

Theorem 2.2 (Lagrange's Theorem) Let G be a finite group, and let H be a subgroup of G . Then the order of H divides the order of G .

Proof. Let $z \in xH \cap yH$. Then $z = xa$ for some $a \in H$ and $z = yb$ for some $b \in H$. For any $h \in H$, $zh = x(ah)$ and $xh = z(a^{-1}h)$, so $zH = xH$. Similarly $yH = zH$, hence $xH = yH$.




Figure 1: Matplotlib: Scatter Plot Example

Regression analysis examines the relationship between variables of interest.

Table 1: Auto MPG and Price

	(1)	(2)
mpg	−238.90***	−49.51
weight		1.75***
constant	11,253***	1,946
obs	74	74
R^2	0.220	0.293

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$