

# Homework 4

Jaden Wang

**Problem (7.1).** Recall that 1 is a generator of  $\mathbb{Z}_{12}$ . If 1 can be represented by a word of  $\{2, 3\}$ , then we can just let the  $\{2, 3\}$  representation of 1 generate  $\mathbb{Z}_{12}$ . This is indeed possible here:

$$1 = 3 +_{12} 3 +_{12} 3 +_{12} 2 +_{12} 2.$$

Therefore,  $\{2, 3\}$  should generate the whole  $\mathbb{Z}_{12}$ .

**Problem (7.2).** Since even numbers under  $+_{12}$  can only output even numbers, we know that the odd numbers  $\{1, 3, 5, 7, 9, 11\}$  cannot be in the subgroup. Let's check the rest:

$$0 = 6 +_{12} 6$$

$$2 = 6 +_{12} 4 +_{12} 4$$

$$4 = 4$$

$$6 = 6$$

$$8 = 4 +_{12} 4$$

$$10 = 4 +_{12} 6$$

Hence, the elements of the subgroup are  $\{0, 2, 4, 6, 8, 10\}$ .

**Problem (7.7).**

$$\text{a) } (a^2b)a^3 = eaabaaa = a^3b$$

$$\text{b) } (ab)(a^3b) = eabaaab = a^2$$

$$\text{c) } b(a^2b) = ebaab = a^2$$

**Problem (7.8).** To fill out the table, we simply need to represent  $c$  in terms of  $a, b$ . The graph tells us  $c = ab = ba$ . Hence we have the following:  $aa = e, ab = c, ac = aab = eb = b, ba = c, bb = e, bc = bba = ea = a, ca = b, cb = a, cc = c(ab) = ce = c$ . That is,

*	e	a	b	c
e	e	a	b	c
a	a	e	c	b
b	b	c	e	a
c	c	b	a	e

**Problem (7.10).** Let's represent other elements using  $a, b$ :  $c = aad = ba, f = ab$ . Thus we have the following:  $aa = c, ab = f, ac = aaa = e, ad = aba = b, af = aab = d, ba = d, bb = e, bc = baa = f, bd = bba = a, bf = bab = c, ca = e, cb = d, cc = caa = a, cd = cba = f, cf = cab = b, da = f, db = c, dc = daa = b, dd = dba = e, df = dab = a, fa = b, fb = a, fc = faa = d, fd = fba = c, ff = fab = e$ .

*	e	a	b	c	d	f
e	e	a	b	c	d	f
a	a	c	f	e	b	d
b	b	d	e	f	a	c
c	c	e	d	a	f	b
d	d	f	c	b	e	a
f	f	b	a	d	c	e

**Problem (8.1).**

$$\begin{aligned}\tau\sigma &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 4 & 1 & 3 & 6 & 5 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 4 & 1 & 3 & 6 & 5 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 6 & 5 & 4 \end{pmatrix}\end{aligned}$$

**Problem (8.2).**

$$\begin{aligned}\tau^2\sigma &= \tau(\tau\sigma) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 4 & 1 & 3 & 6 & 5 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 6 & 5 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 4 & 1 & 5 & 6 & 3 \end{pmatrix}\end{aligned}$$

**Problem (8.6).**

$$\begin{aligned}
\sigma^2 &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 5 & 6 & 2 & 1 \end{pmatrix} \\
\sigma^3 &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 5 & 6 & 2 & 1 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 4 & 6 & 2 & 1 & 3 \end{pmatrix} \\
\sigma^4 &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 4 & 6 & 2 & 1 & 3 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 5 & 2 & 1 & 3 & 4 \end{pmatrix} \\
\sigma^5 &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 5 & 2 & 1 & 3 & 4 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 6 & 1 & 3 & 4 & 5 \end{pmatrix} \\
\sigma^6 &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 4 & 5 & 6 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 6 & 1 & 3 & 4 & 5 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{pmatrix}
\end{aligned}$$

Since  $\sigma^6$  is clearly the identity, we know that  $\sigma^6 = \sigma^0$ . Therefore,  $|\langle \sigma \rangle| = 6$ .

**Problem (8.9).** Notice

$$\begin{aligned}
\mu^2 &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 2 & 4 & 3 & 1 & 6 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 2 & 4 & 3 & 1 & 6 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{pmatrix}
\end{aligned}$$

Hence  $\mu^2 = \mu^0$ . It is easy to see that  $\mu^{100} = \mu^0$  which is the identity above.

**Problem (8.20).**

	$\rho^0$	$\rho$	$\rho^2$	$\rho^3$	$\rho^4$	$\rho^5$
$\rho^0$	$\rho^0$	$\rho$	$\rho^2$	$\rho^3$	$\rho^4$	$\rho^5$
$\rho$	$\rho$	$\rho^2$	$\rho^3$	$\rho^4$	$\rho^5$	$\rho^0$
$\rho^2$	$\rho^2$	$\rho^3$	$\rho^4$	$\rho^5$	$\rho^0$	$\rho$
$\rho^3$	$\rho^3$	$\rho^4$	$\rho^5$	$\rho^0$	$\rho$	$\rho^2$
$\rho^4$	$\rho^4$	$\rho^5$	$\rho^0$	$\rho$	$\rho^2$	$\rho^3$
$\rho^5$	$\rho^5$	$\rho^0$	$\rho$	$\rho^2$	$\rho^3$	$\rho^4$

This cannot be isomorphic to  $S_3$ , since it is symmetric about the diagonal and thus abelian yet  $S_3$  is nonabelian.

**Problem (8.32).** It suffices to show that  $f_3$  has a two-sided inverse to determine it is a permutation. Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  where  $g(x) = -x^{\frac{1}{3}}$ . It is clear that given  $x \in \mathbb{R}$ ,  $f_3 \circ g(x) = -\left(-x^{\frac{1}{3}}\right)^3 = x = -(-x^3)^{\frac{1}{3}} = g \circ f_3(x)$ . Therefore,  $g(x) = f^{-1}(x)$  thus  $f_3$  is bijective and hence is a permutation.

**Problem (8.33).**  $f_4$  is not a permutation because it is not bijective on  $\mathbb{R} \rightarrow \mathbb{R}$ . Choose  $y = -1 \in \mathbb{R}$ , notice that since  $e^x > 0$  for all  $x \in \mathbb{R}$ , there is no  $x \in \mathbb{R}$  such that  $f_4(x) = y < 0$ . This violates the surjectivity requirement.

**Problem (8.34).**  $f_5$  is not a permutation.  $f_5(x) = x^3 - x^2 - 2x = x(x - 2)(x + 1)$ . We can immediately see that it is not bijective because although  $f_4(0) = f_4(-1) = f_4(2) = 0$ , we have  $-1 \neq 0 \neq 2$ , violating the injective requirement.