

5

VITALS & NORMAL RANGES

⚠ Don't let vitals collection delay treatment or evacuation. Write on skin/tape with Sharpie for 911.

VITAL	ABBREV.	NORMAL RANGE
Heart Rate	HR	60–100 beats/min
Respiration	RR	12–20 breaths/min
Blood Pressure	BP	Strong radial pulse
Eyes (Pupils)	PEARL	Equal, Round, Reactive to Light
Skin	SCTM	Warm and Dry
Hands & Feet	CSM	Circulation, Sensation, Movement
Responsiveness	LoR	Alert and oriented



SUPERHERO FIRST AID SYSTEMS

FIELD GUIDE



⚠ STOP — BEFORE YOU HELP

- 1 Is the scene safe?
- 2 How many people need help?
- 3 Call 911
- 4 Gloves + eye protection

Take a breath — you've got this.

⚡ CRITICAL INTERVENTIONS

1

AIRWAY
BREATHING

2

BLEEDING

3

DISABILITY
ENVIRONMENT
GLUCOSE

4

EVACUATION

🔪 ONGOING CARE

5

VITALS

6

MEDICATIONS

7

BLISTERS

⚠ PROTOTYPE · BETA V1.0.0 · FOR TESTING ONLY · NOT FOR CLINICAL USE

6

MEDICATIONS

MEDICATION	USE / NOTES
💊 Acetaminophen (Tylenol)	Fever, pain. Can combine with Ibuprofen. ⚠ Avoid if liver issues
😓 Ibuprofen (Advil)	Pain, inflammation. Can combine with Acetaminophen. ⚠ Avoid if kidney issues
❤ Aspirin	Suspected heart attack (angina or MI) ⚠ Avoid if major bleeding
💊 Loperamide HCl	Diarrhea
😓 Diphenhydramine	Allergic reactions, nausea
🔥 Calcium Carbonate	Acid reflux, heartburn
🌿 Triple Antibiotic (Neosporin)	Infection prevention, keep wounds moist
🔥 Lidocaine Gel	Thermal burns
🦷 Benzocaine	Toothaches, oral pain
🌿 Anti-Itch Gel	Itching from poisonous plants
🐝 Anti-Sting Wipes	Bites and stings irritation
💧 Hydrocortisone 1%	Eczema, bites, jewelry irritation, general rash

Disclaimer: This guide is for informational purposes only and is not a substitute for formal first aid training. Professional CPR and Wilderness First Aid (WFA) training are strongly recommended. Use is at your own risk. I CPR protocols are updated regularly — always reference the American Heart Association (heart.org) as the official authority.

7

BLISTERS

- 1 Pop with a **sterile** safety pin
- 2 Leave intact skin — remove **only** loose/dead skin
- 3 Clean with warm soapy water (if available)
- 4 Clean again with an **alcohol wipe**
- 5 Apply triple antibiotic **over** the blister
- 6 Cover with a band-aid
- 7 Cut 2+ moleskin "**donuts**" — place **AROUND** blister, **NOT ON IT**. Stack to create space between blister and force causing blister.
- 8 Apply Leukotape or duct tape **over** moleskin to secure
- 9 Repeat steps 3–8 every 12–24 hrs

💡 **Bonus — Blister Prevention:** At first sign of a hot spot, alcohol wipe the area and apply Leukotape directly over it.

1 A–B: AIRWAY & BREATHING

⚡ **Major bleeding?** Skip to Section 2, return here after.

A — AIRWAY

- 1 **Check & clear** the airway — remove any visible obstruction.
- 2 **Choking?** Abdominal thrusts from behind.
- 3 **Unresponsive?** Open airway:
→ Jaw-Thrust or Head-Tilt, Chin-Lift

B — BREATHING

- 1 **No breath + no pulse?**
→ **CPR:** 30 compressions, 2 breaths — repeat
→ Ask 911 or bystanders for an AED
→ Maintain CPR → go to Section 2
- 2 **No breath, has pulse?**
→ 1 rescue breath every 6 seconds
→ Maintain rescue breaths → go to Section 2
- 3 **Breathing + pulse present?** → Go to Section 2

2 C — BLEEDING / CIRCULATION

- 1 **Head wound?**
→ ABD pads on wound · gentle pressure
- 2 **Neck, core, back, pelvis?**
→ **Pack** hemostatic gauze deep · firm pressure
- 3 **Arm or leg?**
→ **A.** ABD pads · firm pressure
→ **B.** No stop → **TOURNIQUET** as high as possible on limb
→ **C.** Not on a knee or elbow joint
→ **D.** Tighten until bleeding **stops**
→ **E.** Write the time on their skin
- 4 **Bleeding stopped?** No → repeat 1–3 · Yes → Section 3

↩ Came here for urgent bleeding? Return to Section 1 — Airway.

3 D–E–G: DISABILITY, ENVIRONMENT & GLUCOSE

D — DISABILITY (NEURO CHECK)

- 1 **Ask:** name · time · location · what happened
→ Wrong/no answer → prepare for evacuation
- 2 **Check hands & feet** — movement, sensation, warmth?
→ Any absent → minimize movement (spine/brain injury risk)

E — ENVIRONMENT

- 1 **Cold, wet, or major bleeding?**
→ Remove wet clothing
→ Insulate from the ground
→ Wrap in space blanket

G — GLUCOSE

- 1 **Anxious, confused, or sweating?**
→ Administer glucose between teeth and cheeks

⚠ **Untreated hypoglycemia can cause cardiac arrest.**

4 EVACUATION

EVACUATE IMMEDIATELY IF ANY APPLY

• Major bleeding or burns

• Tourniquet or hemostatics used

• Difficulty breathing

• Confusion, agitation, anxiety

• Hands/feet numb or immobile

• Vitals outside normal ranges

Never be too proud to call for a helicopter and Search & Rescue.