

5 VITALS & NORMAL RANGES

⚠ Don't let vitals collection delay treatment or evacuation. Write on skin/tape with Sharpie for 911.

VITAL	ABBREV.	NORMAL RANGE
Heart Rate	HR	60–100 beats/min
Respiration	RR	12–20 breaths/min
Blood Pressure	BP	Strong radial pulse
Eyes (Pupils)	PEARL	Equal, Round, Reactive to Light
Skin	SCTM	Warm and Dry
Hands & Feet	CSM	Circulation, Sensation, Movement
Responsiveness	LoR	Alert and oriented

6 MEDICATIONS

MEDICATION	USE / NOTES
 Acetaminophen (Tylenol)	Fever, pain. Can combine with Ibuprofen. ⚠ Avoid if liver issues
 Ibuprofen (Advil)	Pain, inflammation. Can combine with Acetaminophen. ⚠ Avoid if kidney issues
 Aspirin	Suspected heart attack (angina or MI) ⚠ Avoid if major bleeding
 Loperamide HCl	Diarrhea
 Diphenhydramine	Allergic reactions, nausea
 Calcium Carbonate	Acid reflux, heartburn
 Triple Antibiotic (Neosporin)	Infection prevention, keep wounds moist
 Lidocaine Gel	Thermal burns
 Benzocaine	Toothaches, oral pain
 Anti-Itch Gel	Itching from poisonous plants
 Anti-Sting Wipes	Bites and stings irritation
 Hydrocortisone 1%	Eczema, bites, jewelry irritation, general rash

Disclaimer: This guide is for informational purposes only and is not a substitute for formal first aid training. Professional CPR and Wilderness First Aid (WFA) training are strongly recommended. Use is at your own risk. I CPR protocols are updated regularly — always reference the American Heart Association (heart.org) as the official authority.



SUPERHERO FIRST AID SYSTEMS

FIELD GUIDE



⚠ STOP — BEFORE YOU HELP

- 1 Is the scene safe?
- 2 How many people need help?
- 3 Call 911
- 4 Gloves + eye protection

Take a breath — you've got this.

⚡ CRITICAL INTERVENTIONS

1

AIRWAY
BREATHING

2

BLEEDING

3

DISABILITY
ENVIRONMENT
GLUCOSE

4

EVACUATION

🔪 ONGOING CARE

5

VITALS

6

MEDICATIONS

7

BLISTERS

⚠ **PROTOTYPE · BETA V1.0.1 · FOR TESTING ONLY · NOT FOR CLINICAL USE**

7 BLISTERS

- 1 Pop with a **sterile** safety pin
- 2 Leave intact skin — remove **only** loose/dead skin
- 3 Clean with warm soapy water (if available)
- 4 Clean again with an **alcohol wipe**
- 5 Apply triple antibiotic **over** the blister
- 6 Cover with a band-aid
- 7 Cut 2+ moleskin "**donuts**" — place **AROUND** blister, **NOT ON IT**. Stack to create space between blister and force causing blister.
- 8 Apply Leukotape or duct tape **over** moleskin to secure
- 9 Repeat steps **3–8** every **12–24 hrs**

💡 **Bonus — Blister Prevention:** At first sign of a hot spot, alcohol wipe the area and apply Leukotape directly over it.

1 A-B: AIRWAY & BREATHING

⚡ **Major bleeding?** Skip to Section 2, return here after.

A — AIRWAY

- 1 **Check & clear** the airway — remove any visible obstruction.
- 2 **Choking?** Abdominal thrusts from behind.
- 3 **Unresponsive?** Open airway:
→ Jaw-Thrust or Head-Tilt, Chin-Lift

B — BREATHING

- 1 **No breath + no pulse?**
→ **CPR:** 30 compressions, 2 breaths — repeat
→ Ask 911 or bystanders for an AED
→ Maintain CPR → go to Section 2
- 2 **No breath, has pulse?**
→ 1 rescue breath every 6 seconds
→ Maintain rescue breaths → go to Section 2
- 3 **Breathing + pulse present?** → Go to Section 2

2 C — BLEEDING / CIRCULATION

- 1 **Head wound?**
→ ABD pads on wound · gentle pressure
→ **Do NOT pack — brain tissue is fragile**
- 2 **Neck, core, back, pelvis?**
→ **Pack** z-folded gauze deep · ABD pads on top · firm pressure
- 3 **Arm or leg?**
→ **A.** ABD pads · firm pressure
→ **B.** If direct pressure isn't stopping the bleeding → **TOURNIQUET** as high as possible on limb
→ **C.** Not on a knee or elbow joint
→ **D.** Tighten until bleeding **stops**
→ **E.** **Write the time** on their skin
- 4 **Bleeding stopped?** No → repeat 1–3 · Yes → Section 3

↩ Came here for urgent bleeding? Return to Section 1 — Airway.

3 D-E-G: DISABILITY, ENVIRONMENT & GLUCOSE

D — DISABILITY (NEURO CHECK)

- 1 **Ask:** name · time · location · what happened
→ Wrong/no answer → prepare for evacuation
- 2 **Check hands & feet** — movement, sensation, warmth?
→ Any absent → minimize movement (spine/brain injury risk)

E — ENVIRONMENT

- 1 **Cold, wet, or major bleeding?**
→ Remove wet clothing
→ Insulate from the ground
→ Wrap in space blanket

G — GLUCOSE

- 1 **Anxious, confused, or sweating?**
→ Administer glucose between teeth and cheeks

⚠ **Untreated hypoglycemia can cause cardiac arrest.**

4 EVACUATION

EVACUATE IMMEDIATELY IF ANY APPLY

- | | |
|-------------------------------|----------------------------------|
| • Major bleeding or burns | • Tourniquet or hemostatics used |
| • Difficulty breathing | • Confusion, agitation, anxiety |
| • Hands/feet numb or immobile | • Vitals outside normal ranges |

Never be too proud to call for a helicopter and Search & Rescue.