

## 5 VITALS & NORMAL RANGES

⚠ Don't let vitals collection delay treatment or evacuation. Write on skin/tape with Sharpie for 911.

| VITAL          | ABBREV. | NORMAL RANGE                     |
|----------------|---------|----------------------------------|
| Heart Rate     | HR      | 60–100 beats/min                 |
| Respiration    | RR      | 12–20 breaths/min                |
| Blood Pressure | BP      | <b>Strong</b> radial pulse       |
| Eyes (Pupils)  | PEARL   | Equal, Round, Reactive to Light  |
| Skin           | SCTM    | Warm and Dry                     |
| Hands & Feet   | CSM     | Circulation, Sensation, Movement |
| Responsiveness | LoR     | Alert and oriented               |

## 6 MEDICATIONS

| MEDICATION                                                                                                             | USE / NOTES                                                                            |
|------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------|
|  <b>Acetaminophen (Tylenol)</b>        | Fever, pain. Can combine with Ibuprofen.<br>⚠ <b>Avoid if liver issues</b>             |
|  <b>Ibuprofen (Advil)</b>             | Pain, inflammation. Can combine with Acetaminophen.<br>⚠ <b>Avoid if kidney issues</b> |
|  <b>Aspirin</b>                       | Suspected heart attack (angina or MI)<br>⚠ <b>Avoid if major bleeding</b>              |
|  <b>Loperamide HCl</b>                | Diarrhea                                                                               |
|  <b>Diphenhydramine</b>               | Allergic reactions, nausea                                                             |
|  <b>Calcium Carbonate</b>             | Acid reflux, heartburn                                                                 |
|  <b>Triple Antibiotic (Neosporin)</b> | Infection prevention, keep wounds moist                                                |
|  <b>Lidocaine Gel</b>                 | Thermal burns                                                                          |
|  <b>Benzocaine</b>                    | Toothaches, oral pain                                                                  |
|  <b>Anti-Itch Gel</b>                 | Itching from poisonous plants                                                          |
|  <b>Anti-Sting Wipes</b>              | Bites and stings irritation                                                            |
|  <b>Hydrocortisone 1%</b>             | Eczema, bites, jewelry irritation, general rash                                        |

Disclaimer: This guide is for informational purposes only and is not a substitute for formal first aid training. Professional CPR and Wilderness First Aid (WFA) training are strongly recommended. Use is at your own risk. I CPR protocols are updated regularly — always reference the American Heart Association (heart.org) as the official authority.



SUPERHERO FIRST AID SYSTEMS

# FIELD GUIDE



## ⚠ STOP — BEFORE YOU HELP

- 1 Is the scene safe?
- 2 How many people need help?
- 3 Call 911
- 4 Gloves + eye protection

Take a breath — you've got this.

## ⚡ CRITICAL INTERVENTIONS

1

AIRWAY  
BREATHING

2

BLEEDING

3

DISABILITY  
ENVIRONMENT  
GLUCOSE

4

EVACUATION

## 🔪 ONGOING CARE

5

VITALS

6

MEDICATIONS

7

BLISTERS

⚠ **PROTOTYPE · BETA V1.0.1 · FOR TESTING ONLY · NOT FOR CLINICAL USE**

## 7 BLISTERS

- 1 Pop with a **sterile** safety pin
- 2 Leave intact skin — remove **only** loose/dead skin
- 3 Clean with warm soapy water (if available)
- 4 Clean again with an **alcohol wipe**
- 5 Apply triple antibiotic **over** the blister
- 6 Cover with a band-aid
- 7 Cut 2+ moleskin "**donuts**" — place **AROUND** blister, **NOT ON IT**. Stack to create space between blister and force causing blister.
- 8 Apply Leukotape or duct tape **over** moleskin to secure
- 9 Repeat steps **3–8** every **12–24 hrs**

💡 **Bonus — Blister Prevention:** At first sign of a hot spot, alcohol wipe the area and apply Leukotape directly over it.

## 1 A-B: AIRWAY & BREATHING

⚡ **Major bleeding?** Skip to Section 2, return here after.

### A — AIRWAY

- 1 **Check & clear** the airway — remove any visible obstruction.
- 2 **Choking?** Abdominal thrusts from behind.
- 3 **Unresponsive?** Open airway:  
→ Jaw-Thrust or Head-Tilt, Chin-Lift

### B — BREATHING

- 1 **No breath + no pulse?**  
→ **CPR:** 30 compressions, 2 breaths — repeat  
→ Ask 911 or bystanders for an AED  
→ Maintain CPR → go to Section 2
- 2 **No breath, has pulse?**  
→ 1 rescue breath every 6 seconds  
→ Maintain rescue breaths → go to Section 2
- 3 **Breathing + pulse present?** → Go to Section 2

## 2 C — BLEEDING / CIRCULATION

- 1 **Head wound?**  
→ ABD pads on wound · gentle pressure  
→ **Do NOT pack — brain tissue is fragile**
- 2 **Neck, core, back, pelvis?**  
→ **Pack** hemostatic gauze deep · firm pressure
- 3 **Arm or leg?**  
→ **A.** ABD pads · firm pressure  
→ **B.** If direct pressure isn't stopping the bleeding → **TOURNIQUET** as high as possible on limb  
→ **C.** Not on a knee or elbow joint  
→ **D.** Tighten until bleeding **stops**  
→ **E.** **Write the time** on their skin
- 4 **Bleeding stopped?** No → repeat 1–3 · Yes → Section 3

↩ Came here for urgent bleeding? Return to Section 1 — Airway.

## 3 D-E-G: DISABILITY, ENVIRONMENT & GLUCOSE

### D — DISABILITY (NEURO CHECK)

- 1 **Ask:** name · time · location · what happened  
→ Wrong/no answer → prepare for evacuation
- 2 **Check hands & feet** — movement, sensation, warmth?  
→ Any absent → minimize movement (spine/brain injury risk)

### E — ENVIRONMENT

- 1 **Cold, wet, or major bleeding?**  
→ Remove wet clothing  
→ Insulate from the ground  
→ Wrap in space blanket

### G — GLUCOSE

- 1 **Anxious, confused, or sweating?**  
→ Administer glucose between teeth and cheeks

⚠ **Untreated hypoglycemia can cause cardiac arrest.**

## 4 EVACUATION

### EVACUATE IMMEDIATELY IF ANY APPLY

- |                               |                                  |
|-------------------------------|----------------------------------|
| • Major bleeding or burns     | • Tourniquet or hemostatics used |
| • Difficulty breathing        | • Confusion, agitation, anxiety  |
| • Hands/feet numb or immobile | • Vitals outside normal ranges   |

Never be too proud to call for a helicopter and Search & Rescue.