

5

VITALS & NORMAL RANGES

⚠ Don't let vitals collection delay treatment or evacuation. Write on skin/tape with Sharpie for 911.

VITAL	ABBREV.	NORMAL RANGE
Heart Rate	HR	60–100 beats/min
Respiration	RR	12–20 breaths/min
Blood Pressure	BP	Strong radial pulse
Eyes (Pupils)	PEARL	Equal, Round, Reactive to Light
Skin	SCTM	Warm and Dry
Hands & Feet	CSM	Circulation, Sensation, Movement
Responsiveness	LoR	Alert and oriented

6

MEDICATIONS

MEDICATION	USE / NOTES
💊 Acetaminophen (Tylenol)	Fever, pain. Can combine with Ibuprofen. ⚠ Avoid if liver issues
💊 Ibuprofen (Advil)	Pain, inflammation. Can combine with Acetaminophen. ⚠ Avoid if kidney issues
❤ Aspirin	Suspected heart attack (angina or MI) ⚠ Avoid if major bleeding
💊 Loperamide HCl	Diarrhea
💊 Diphenhydramine	Allergic reactions, nausea
🔥 Calcium Carbonate	Acid reflux, heartburn
🌿 Triple Antibiotic (Neosporin)	Infection prevention, keep wounds moist
🔥 Lidocaine Gel	Thermal burns
🦷 Benzocaine	Toothaches, oral pain
🌿 Anti-Itch Gel	Itching from poisonous plants
🦋 Anti-Sting Wipes	Bites and stings irritation
👤 Hydrocortisone 1%	Eczema, bites, jewelry irritation, general rash

Disclaimer: This guide is for informational purposes only and is not a substitute for formal first aid training. Professional CPR and Wilderness First Aid (WFA) training are strongly recommended. Use is at your own risk. | CPR protocols are updated regularly — always reference the American Heart Association (heart.org) as the official authority.

SUPERHERO FIRST AID SYSTEMS

FIELD GUIDE

⚠ STOP — BEFORE YOU HELP

- 1

Is the scene safe?
- 2

How many people need help?

3

Call 911

4

Gloves + eye protection

Take a breath — you've got this.

⚡ CRITICAL INTERVENTIONS

1

AIRWAY
BREATHING

2

BLEEDING

3

DISABILITY
ENVIRONMENT
GLUCOSE

4

EVACUATION

🩹 ONGOING CARE

5

VITALS

6

MEDICATIONS

7

BLISTERS

⚠ PROTOTYPE · BETA V1.0.0 · FOR TESTING ONLY · NOT FOR CLINICAL USE

7

BLISTERS

- 1

Pop with a **sterile** safety pin
- 2

Leave intact skin — remove **only** loose/dead skin
- 3

Clean with warm soapy water (if available)
- 4

Clean again with an **alcohol wipe**
- 5

Apply triple antibiotic **over** the blister
- 6

Cover with a band-aid
- 7

Cut 2+ moleskin "**donuts**" — place **around** blister, **not on it**. Stack to create space between blister and force causing blister.
- 8

Apply Leukotape or duct tape **over** moleskin to secure
- 9

Repeat steps 3–8 every **12–24 hrs**

💡 **Bonus — Blister Prevention:** At first sign of a hot spot, alcohol wipe the area and apply Leukotape directly over it.

⚡ Major bleeding? Skip to Section 2, return here after.

A — AIRWAY

1 Check & clear the airway — remove any visible obstruction.

2 Choking? Abdominal thrusts from behind.

3 Unresponsive? Open airway:

Jaw-Thrust or Head-Tilt, Chin-Lift

B — BREATHING

1 No breath + no pulse?

→ CPR: 30 compressions, 2 breaths — repeat

→ Ask 911 or bystanders for an AED

→ Maintain CPR → go to Section 2

2 No breath, has pulse?

→ 1 rescue breath every 6 seconds

→ Maintain rescue breaths → go to Section 2

3 Breathing + pulse present? → Go to Section 2

2 C — BLEEDING / CIRCULATION

1 Head wound?

ABD pads on wound · gentle pressure

2 Neck, core, back, pelvis?

Pack hemostatic gauze deep · firm pressure

3 Arm or leg?

A ABD pads · firm pressure

B No stop → **TOURNIQUET** as high as possible on limb

C Not on a knee or elbow joint

D Tighten until bleeding stops

E Write the time on their skin

4 Bleeding stopped? No → repeat 1-3 · Yes → Section 3

↔ Came here for urgent bleeding? Return to Section 1 — Airway.

D — DISABILITY (NEURO CHECK)

1 Ask: name · time · location · what happened

Wrong/no answer → prepare for evacuation

2 Check hands & feet — movement, sensation, warmth?

Any absent → minimize movement (spine/brain injury risk)

E — ENVIRONMENT

1 Cold, wet, or major bleeding?

→ Remove wet clothing

→ Insulate from the ground

→ Wrap in space blanket

G — GLUCOSE

1 Anxious, confused, or sweating?

→ Administer glucose between teeth and cheeks

⚠ Untreated hypoglycemia can cause cardiac arrest

4 EVACUATION

EVACUATE IMMEDIATELY IF ANY APPLY

• Major bleeding or burns

• Difficulty breathing

• Hands/feet numb or immobile

• Tourniquet or hemostatics used

• Confusion, agitation, anxiety

• Vitals outside normal ranges

Never be too proud to call for a helicopter and Search & Rescue.