



Primary pulmonary meningioma and minute pulmonary meningotheial-like nodules: Rare pulmonary nodular lesions requiring more awareness in clinical practice

Li-Dan Liu, Ke-Xin Zhang, Hai-Ning Zhang, Yi-Wen Zheng, Hong-Tao Xu

Specialty type: Pathology

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Handra-Luca A, France

Received: November 28, 2023

Peer-review started: November 28, 2023

First decision: January 17, 2024

Revised: January 25, 2024

Accepted: March 21, 2024

Article in press: March 21, 2024

Published online: April 16, 2024



Li-Dan Liu, Ke-Xin Zhang, Hai-Ning Zhang, Yi-Wen Zheng, Hong-Tao Xu, Department of Pathology, The First Hospital and College of Basic Medical Sciences of China Medical University, Shenyang 110001, Liaoning Province, China

Corresponding author: Hong-Tao Xu, MD, PhD, Professor, Department of Pathology, The First Hospital and College of Basic Medical Sciences of China Medical University, No. 155 Nanjingbei Street, Heping District, Shenyang 110001, Liaoning Province, China.

xuht@cmu.edu.cn

Abstract

In this editorial, we comment on an article by Ruan *et al* published in a recent issue of the *World Journal of Clinical Case*. Pulmonary meningotheial proliferative lesions, including primary pulmonary meningiomas, minute pulmonary meningotheial-like nodules, and metastatic pulmonary meningiomas are rare pulmonary lesions. These lesions are difficult to differentiate from lung cancers based on clinical and imaging manifestations. Herein, we briefly introduce the clinical, imaging, and pathological characteristics of these lesions and discuss their pathogenesis to strengthen the current understanding of pulmonary meningotheial proliferative lesions in clinical diagnosis and therapy.

Key Words: Pulmonary meningotheial proliferation; Primary pulmonary meningioma; Minute pulmonary meningotheial-like nodule; Lung neoplasm; Rare pulmonary nodular lesion

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Primary pulmonary meningiomas (PPM) and minute pulmonary meningothelial-like nodules (MPMN) are rare pulmonary lesions, which are difficult to differentiate from lung cancers due to similarities in clinical and imaging manifestations. To avoid misdiagnosis and overtreatment, PPM and MPMN should be considered in the differential diagnoses of pulmonary nodules of difficult clinical and/or imaging procedure diagnosis, particularly of asymptomatic patients. Differentiating PPM and MPMN from other pulmonary nodule diseases and assessing the malignancy of PPMs are difficult using only imaging-procedures, and histopathological examination thus remains the gold standard diagnostic procedure. Enhancing our understanding of MPMPNs and PPMs and elucidating their pathogenesis will aid in the accurate differentiation of these lesions, to improve the clinical management of patients.

Citation: Liu LD, Zhang KX, Zhang HN, Zheng YW, Xu HT. Primary pulmonary meningioma and minute pulmonary meningothelial-like nodules: Rare pulmonary nodular lesions requiring more awareness in clinical practice. *World J Clin Cases* 2024; 12(11): 1857-1862

URL: <https://www.wjgnet.com/2307-8960/full/v12/i11/1857.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i11.1857>

INTRODUCTION

Pulmonary meningothelial proliferative lesions, including primary pulmonary meningiomas (PPMs), minute pulmonary meningothelial-like nodules (MPMPNs) and metastatic pulmonary meningiomas (MPMs), are rare pulmonary lesions[1]. PPMs and MPMPNs are usually discovered incidentally during physical examination or lung surgeries performed for other indications. In recent years, pulmonary nodules have become more commonly, due to the popularization of regular physical examinations and the development of medical imaging technology, thus raising the attention of clinical physicians and pathologists. Consequently, the incidence of MPMPNs and PPMs have also significantly increased. Additionally, MPM is very rare, and diagnosis depends on whether the patient has a medical history of intracranial or spinal meningioma. These lesions are difficult to differentiate from lung cancers based solely on clinical and imaging manifestations. Furthermore, due to differences in biological behavior and treatment methods, it is also necessary to differentiate between PPM, MPMPN, and MPM. In a recent issue of the *World Journal of Clinical Cases*, Ruan *et al*[2] reported a rare case of MPMPN and explained its histomorphology and immunohistochemical features. In this editorial, we review the recent developments in the clinical, imaging, and pathological characteristics of PPM and MPMPN and discuss their pathogenesis, in order to strengthen the current clinical understanding of pulmonary meningothelial proliferative lesion diagnosis and treatment.

CLINICAL FEATURES OF PPM AND MPMPN

Meningiomas are common tumors of the nervous system originating from meningothelial cells of the arachnoid layer, often occurring in the intracranial or spinal canal. Meningiomas occurring in tissues and organs not covered by the meninges are termed ectopic meningiomas. PPM is defined as a primary meningioma in the lung tissue[3] and is classified as a rare condition under the category of lung tumors of ectopic tissues according to the World Health Organization (WHO) Classification of Thoracic Tumors[4].

Patients with PPMs and MPMPNs usually present with no obvious symptoms, other than some atypical respiratory symptoms, although this can depend on tumor size, location, and accompanying diseases[5-7]. PPMs are usually found through imaging examinations[5,8]. Patients with PPMs are mainly middle-aged or older women[1,7,9]. The vast majority of PPMs are benign, well-circumscribed and solitary, with a favorable prognosis[5,10-12]. Therefore, the major therapeutic method for PPM is surgery, such as wedge resection or lobectomy[8,10,13,14]. It has also been suggested that a conservative approach, using close medical monitoring, is the best option for patients when the location, size, and number of smaller nodules challenge the accessibility for biopsy or resection[13,15]. Malignant PPMs are prone to recurrence or metastasis, leading to a poor prognosis[3,16,17]. Conversely, some reports have indicated that the prognosis of patients with malignant PPM is actually favorable owing to a typically slow progression[6].

IMAGING FEATURES OF PPM AND MPMPN

Radiologically, PPMs most commonly appear in the form of pulmonary nodules[5]. However, the visual features of PPM nodules are not universal, for example, they can appear either as a solitary nodule or in clusters[7,15]. The distribution of nodules is frequently subpleural (less than 1 cm away from the visceral pleura), whereas others are intrapulmonary (greater than 1 cm away from the visceral pleura), with these two presentations accounting for 89.1% and 10.9% of cases, respectively[18]. The edges of PPM nodules can also either be close to pulmonary blood vessels or visually away from the bronchi, blood vessels, or pleura[7,18]. There is usually no preferred location for PPM in the lung[19].

PPM nodules usually manifest as a round homogeneous mass with well-demarcated borders on imaging examinations [5,7,13,18]. PPMs can appear with ground-glass opacity and cyst-like structures or calcifications, depending on the changes in the lesion [7,18]. Small, early-stage PPMs can manifest as ground-glass like lesions, which are easily confused with early-stage lung cancers. As the lesions grow, it can become difficult to differentiate them from other benign lung tumors, such as pulmonary hamartoma, sclerosing pneumocytoma, lipoma, and certain infections (such as cryptococcal infection or virus infection). It is also difficult to differentiate PPMs from aggressive growths or multiple advanced lung cancer nodules [10].

Approximately 74.0% of PPMs are less than 3 cm in diameter [7]. The size of benign PPM tumors ranges from 0.4 to 6 cm in diameter (median: 2 cm), whereas malignant PPM tumors range in size from 1.5 to 15 cm in diameter (median: 6.4 cm) [7]. Although the average diameter of malignant PPMs is larger than that of benign PPMs, it is impossible to diagnose them based solely on tumor size. In one PPM case, the tumor measured 9.5 cm × 8.4 cm × 5.3 cm in size, but pathological examination confirmed it was benign [3].

MPMNs are usually smaller than PPMs, ranging from 2.5 to 5.0 mm in diameter on CT scans, with an average of 3.04 ± 1.12 mm [18]. Therefore, MPMNs are difficult to discover during imaging examinations. In addition, some larger nodules may be misrepresented as PPM due to their increasing prevalence [13].

PET/CT has been proven to be more accurate than traditional CT in evaluating solitary pulmonary nodules. Low uptake of 18F-fluorodeoxyglucose (18F-FDG), defined as a maximum standardized uptake value (SUVmax) less than 1.5–2.0, is generally associated with a benign diagnosis [20], however, PPMs can present with various CT manifestations in enhanced imaging [6,7,21]. Therefore, the pattern of enhancement may not help determine whether the PPM is benign or malignant [7,22,23]. PPMs exhibit anywhere from mild to high metabolic activity, and the average value of SUVmax is 4.36, ranging from 0.6 to 12.9, although this factor has no relevance for malignant evaluation [3]. A recent study reported increased 18F-FDG uptake in synchronous benign and malignant PPMs [23]. In patients with a history of cancer, PPMs or MPMNs can be easily mistaken as lung metastases due to deceptive increases in 18F-FDG uptake [24,25]. As such, a single 18F-FDG PET or contrast-enhanced CT examination may not be sufficient to evaluate patients with PPM.

Overall, the imaging manifestations of PPMs are often heterogeneous between different patients. Although accurate diagnoses based on imaging results is of great significance for developing appropriate treatment plans, it is currently impossible to differentiate PPM from other pulmonary nodule diseases or to differentiate between benign and malignant PPM through radiographic observation alone [6,10]. Thus, pathological examination is the most accurate method of diagnosis of PPM [7,8,10].

PATHOLOGICAL FEATURES OF PPM AND MPMN HISTOLOGICAL FEATURES

PPM, MPMN, and MPM have similar morphological and immunohistochemical features [1]. PPMs share the same histomorphology and WHO grading with that of central nerve system (CNS) meningiomas [8,19]. The premise underlying the diagnosis of PPM is that no CNS lesions are found in imaging modalities and that MPM can be ruled out [26]. In histopathological analyses, PPMs are typically categorized as epithelial, fibrous or transitional [13]. Further, they often exhibit spindle cells in whirlpool arrangements, while psammoma bodies can also be observed [5,6,8,13,15,19]. Pseudonuclear inclusions have also been reported in PPM histopathology [5,8,13]. No mitoses or necrosis has been identified in benign PPMs [5,8]. Finally, high proliferative activity with several mitotic figures and a high Ki-67 index may be indicative of malignant PPM [23].

MPMNs are interstitial cellular proliferations first identified in 1960 [27]. MPMNs can often appear with no visible abnormal lesions, or sometimes as pale, reddish gray or brown, and regular or irregular solid nodules, with moderate or soft texture, and unclear boundaries [28]. Like PPMs, MPMN can also present as a solitary or multiple nodules. In cases of MPMN which present in a diffuse, miliary like distribution in both lungs (up to tens, hundreds, or uncountable), it is defined as “diffuse pulmonary meningeal epithelial neoplasia”. Under the microscope, MPMNs exhibit elliptical nodules ranging in size from 1.0 to 5.0 mm. Most lesions tend to be subpleural, or located in the pulmonary interstitium, and distributed along the alveolar septa. Tumor cells are spindle shaped or epithelioid and arranged in a nest or vortex shape around small veins. The nuclei are ovoid with delicate chromatin and indistinct nucleoli with a rich and eosinophilic cytoplasm and an ambiguous cell boundary. Pseudoinclusion bodies can be seen in some nuclei, and mitotic figures are rare [1]. It is difficult to differentiate MPMN from PPM because of their morphologic similarity, with an exception to their size differences [29].

Even though histological diagnosis of PPM may not appear challenging according to morphological and immunostaining characteristics, misdiagnosis remains an issue. When PPM contains psammoma bodies, it can be misdiagnosed as metastatic papillary thyroid carcinoma [30]. In addition, approximately 90% of patients with MPMNs have synchronous lung cancer-related lesions, including atypical adenomatous hyperplasia, adenocarcinoma *in situ*, minimally invasive adenocarcinoma, and invasive adenocarcinoma [1].

IMMUNOHISTOCHEMICAL FEATURES

PPM and MPMN often overlap in terms of histological and immunohistochemical phenotypes, which are consistent with the manifestations of CNS meningioma [31]. Specific immunohistochemical staining is useful for diagnosing PPM and MPMN and excluding other types of lung tumors.

Positive immunohistochemical staining for vimentin, epithelial cell membrane antigen, and progesterone receptor (PR) is common in most cases of PPM and MPMN[5,6,8,13,18], whereas markers such as S100, cytokeratin (CK), and CD34 may exhibit variable or inconsistent staining results[1,8,15,19,26]. The expression of thyroid transcription factor-1 and neuroendocrine markers, such as synaptophysin and chromogranin, are negative[6,31]. Tao *et al*[32] previously reported that MPMNs showed strong and diffuse cytoplasmic expression of somatostatin receptor 2A (SSTR2A) and that the expression rate of SSTR2A was higher than that of conventional markers of meningioma.

PATHOGENESIS OF PPM AND MPMN

The incidence of ectopic meningioma is very rare (2%) and can be found in various anatomical sites such as the head and neck region, and specifically in the sinonasal tract, ear and temporal bone, and scalp[33]. However, meningiomas originating from the lungs are even more rare. Although the pathogenesis of PPM and MPMN remains unclear, it is generally believed that ectopic meningioma is derived from pluripotential subpleural mesenchymal cells or heterotopic embryonic rests of arachnoid cells[3]. Some scholars believe that PPMs originate from MPMNs based on their similar morphological, immunohistochemical and ultrastructural features[8,34]. Kraushaar *et al*[35] suggested that PPM can appear as a giant form of MPMN. Presentation of MPMNs as a single lesion lacking genetic alterations may indicate its origin of reactivity. MPMN-omatosis syndrome, defined as the diffuse distribution of MPMN, may represent the transition between reactive and neoplastic proliferations[31].

PPM and MPMN are more common in females. Further, as both PPM and MPMN cells express PR, the occurrence of these tumors may be related to PR activity. Moreover, PR develop from normal meningeal epithelial cells and the expression in meningiomas has been well confirmed, with results indicating the potential presence of lung meningeal epithelioid cells in normal lung tissues[2].

Patients with malignant lung tumors tend to have a higher incidence of MPMN[18,36]. Further, the microenvironment theory suggests that the external environment is the main contributing factor to the occurrence of MPMNs. Therefore, lung diseases may provide a localized microenvironment that promotes the occurrence and development of MPMN[36]. One case report of a patient with a *BRCA2* germline mutation exhibited quadruple neoplasms, including PPM. *BRCA2* mutations can increase tumor sensitivity in patients, but further research is needed to determine whether the *BRCA2* germline mutation could be related to PPMs[37].

CONCLUSION

The accurate diagnosis of PPM and MPMN is challenging due to their rarity. To avoid misdiagnosis and overtreatment in clinical practice, PPM and MPMN should be considered as differential diagnoses of pulmonary nodules of difficult clinical and/or imaging procedure diagnosis, in particular asymptomatic patients. Differentiating PPM and MPMN from other pulmonary nodule diseases and assessing the malignancy of PPMs are difficult using only imaging procedures. As such, histopathological examination remains the gold standard for diagnosing PPM and MPMN. Moreover, as most PPM and MPMN have synchronous lung cancer-related lesions, ground-glass-like changes which PPM and MPMN often exhibit in imaging examinations may cause the primary malignant lung tumor to exhibit metastatic malignancy, leading to delayed or canceled treatment for the primary malignant lung tumor[29]. Wedge resection is so far the best option for the diagnosis and treatment of these lesions, as no recurrence has been reported in benign cases after complete resection. As PPM is usually benign, conservative monitoring by biopsy is also a good treatment strategy, particularly if the locations of the lesions are inconvenient for surgical operation. In these cases, asymptomatic patients can undergo long-term follow-up observation instead. Enhancing our understanding of MPMN and PPM features and elucidating the pathogenesis underlying the development of these lesions could help to increase the clinical accuracy of PPM and MPMN differentiation, thus improving the clinical management of patients.

FOOTNOTES

Author contributions: Liu LD was responsible for investigation and writing the original draft; Zhang KX, Zhang HN, and Zheng YW were responsible for investigation; Xu HT was responsible for conceptualization, writing, reviewing and editing; all authors have read and approve the final manuscript.

Conflict-of-interest statement: Dr. Xu has nothing to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Ke-Xin Zhang 0000-0002-4275-6418; Hai-Ning Zhang 0000-0002-7608-7080; Hong-Tao Xu 0000-0001-5062-3108.

S-Editor: Lin C

L-Editor: A

P-Editor: Yu HG

REFERENCES

- 1 Lin D, Yu Y, Wang H, Fang Y, Yin J, Shen Y, Tan L. Radiological manifestations, histological features and surgical outcomes of pulmonary meningotheial proliferation: a case series and rethinking. *Transl Lung Cancer Res* 2020; **9**: 1159-1168 [PMID: 32953494 DOI: 10.21037/tlcr-19-699]
- 2 Ruan X, Wu LS, Fan ZY, Liu Q, Yan J, Li XQ. Pathological diagnosis and immunohistochemical analysis of minute pulmonary meningotheial-like nodules: A case report. *World J Clin Cases* 2023; **11**: 8022-8029 [PMID: 38075568 DOI: 10.12998/wjcc.v11.i33.8022]
- 3 Feng Y, Wang P, Liu Y, Dai W. PET/CT imaging of giant primary pulmonary meningioma: a case report and literature review. *J Cardiothorac Surg* 2023; **18**: 171 [PMID: 37138316 DOI: 10.1186/s13019-023-02276-4]
- 4 Radiopaedia. WHO classification of tumours. [cited 25 January 2024]. Available from: <https://radiopaedia.org/articles/who-classification-of-skin-tumours?lang=us>
- 5 Baş A, Valiyev E, Özkan ND, Tombul İ, Yonat S, Sayan M, Kurul İC. A Rare Entity: Primary Pulmonary Meningioma. *Türk Patoloji Derg* 2023; **39**: 98-99 [PMID: 34514565 DOI: 10.5146/tjpath.2021.01535]
- 6 Jiang M, Chen P, Huang R, Zhang J, Zheng J. A case report of primary pulmonary meningioma masquerading as lung metastasis in a patient with rectal carcinoma: role of (18)F-FDG PET/CT. *J Cardiothorac Surg* 2021; **16**: 153 [PMID: 34051819 DOI: 10.1186/s13019-021-01546-3]
- 7 Zhang DB, Chen T. Primary pulmonary meningioma: A case report and review of the literature. *World J Clin Cases* 2022; **10**: 4196-4206 [PMID: 35665099 DOI: 10.12998/wjcc.v10.i13.4196]
- 8 Yang B, Qiu J. Primary pulmonary meningioma with associated multiple micronodules: a case report and literature review. *J Surg Case Rep* 2023; **2023**: rjad034 [PMID: 36755931 DOI: 10.1093/jscr/rjad034]
- 9 Gou XN, Liu QY. [Minute pulmonary meningotheial-like nodules: a clinicopathological study of 97 cases]. *Zhonghua Bing Li Xue Za Zhi* 2022; **51**: 1260-1262 [PMID: 36480838 DOI: 10.3760/cma.j.cn112151-20220413-00275]
- 10 Huang X, Mou YF, Ren FQ, Wang Y, Yang Y. Multiple primary pulmonary meningioma: A case report and literature review. *Thorac Cancer* 2022; **13**: 2257-2259 [PMID: 35734847 DOI: 10.1111/1759-7714.14542]
- 11 Tang H, He Y, Wang L, Wu G, Xu Y, Guo D. Coexistence of primary pulmonary meningioma and metastatic papillary renal cell carcinoma of the lung: A rare case report with review of the literature. *Medicine (Baltimore)* 2022; **101**: e32157 [PMID: 36550861 DOI: 10.1097/MD.00000000000032157]
- 12 Satoh Y, Ishikawa Y. Multiple primary pulmonary meningiomas: 20-year follow-up findings for a first reported case confirming a benign biological nature. *Int J Surg Case Rep* 2017; **31**: 58-60 [PMID: 28107759 DOI: 10.1016/j.ijscr.2017.01.002]
- 13 Chakrabarti R, Ghuman D. Diagnostic and Management Considerations in a Patient With Primary Pulmonary Meningioma With Associated Micro-Solid Nodules. *Cureus* 2021; **13**: e15700 [PMID: 34277287 DOI: 10.7759/cureus.15700]
- 14 Gürcay N, Öztürk A, Demirağ F, İncekara F. Primary pulmonary meningioma mimicking pulmonary metastasis: A rare case report. *Türk Gogus Kalp Damar Cerrahisi Derg* 2020; **28**: 699-701 [PMID: 33403148 DOI: 10.5606/tgkdc.dergisi.2020.19370]
- 15 Oh JH, Cho HS, Hwang HS, Ji W. Primary pulmonary meningioma presenting as multiple lung nodules: A case report. *Thorac Cancer* 2022; **13**: 141-143 [PMID: 34878222 DOI: 10.1111/1759-7714.14270]
- 16 Minami Y, Sato S, Koyanagi H, Kinowaki Y. Malignant primary pulmonary meningioma with bone metastasis. *Oxf Med Case Reports* 2020; **2020**: omaa005 [PMID: 32128218 DOI: 10.1093/omcr/omaa005]
- 17 Xu KK, Tian F, Cui Y. Primary pulmonary meningioma presenting as a micro solid nodule: A rare case report. *Thorac Cancer* 2018; **9**: 874-876 [PMID: 29718593 DOI: 10.1111/1759-7714.12639]
- 18 Zhang Y, Wu J, Zhao Y, Zhang T, Xu H, Chen YC. Minute pulmonary meningotheial-like nodules: associations between computed tomography and pathology features. *Quant Imaging Med Surg* 2023; **13**: 462-470 [PMID: 36620132 DOI: 10.21037/qims-21-1250]
- 19 Moran CA, Hochholzer L, Rush W, Koss MN. Primary intrapulmonary meningiomas. A clinicopathologic and immunohistochemical study of ten cases. *Cancer* 1996; **78**: 2328-2333 [PMID: 8941002]
- 20 Fletcher JW, Kymes SM, Gould M, Alazraki N, Coleman RE, Lowe VJ, Marn C, Segall G, Thet LA, Lee K; VA SNAP Cooperative Studies Group. A comparison of the diagnostic accuracy of 18F-FDG PET and CT in the characterization of solitary pulmonary nodules. *J Nucl Med* 2008; **49**: 179-185 [PMID: 18199626 DOI: 10.2967/jnumed.107.044990]
- 21 Hong S, Jiang J, Zhou F, Liu J. Computed tomography findings of primary pulmonary meningioma: A case report. *Medicine (Baltimore)* 2018; **97**: e9651 [PMID: 29480880 DOI: 10.1097/MD.0000000000009651]
- 22 Hsu CC, Tsai YM, Yang SF, Hsu JS. Primary pulmonary meningioma. *Kaohsiung J Med Sci* 2023; **39**: 1155-1156 [PMID: 37698283 DOI: 10.1002/kjm2.12754]
- 23 Cimini A, Ricci F, Pugliese L, Chiaravallotti A, Schillaci O, Floris R. A Patient with a Benign and a Malignant Primary Pulmonary Meningioma: An Evaluation with 18F Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography and Computed Tomography with Iodinated Contrast. *Indian J Nucl Med* 2019; **34**: 45-47 [PMID: 30713380 DOI: 10.4103/ijnm.IJNM_101_18]
- 24 Fujikawa R, Arai Y, Otsuki Y, Nakamura T. A case of a primary pulmonary meningioma mimicking a metastasis from a papillary thyroid carcinoma due to a size reduction after radioactive iodine therapy. *Surg Case Rep* 2020; **6**: 57 [PMID: 32221747 DOI: 10.1186/s40792-020-00823-y]
- 25 Juan CM, Chen ML, Ho SY, Huang YC. Primary Pulmonary Meningioma Simulating a Pulmonary Metastasis. *Case Rep Pulmonol* 2016; **2016**: 8248749 [PMID: 27974986 DOI: 10.1155/2016/8248749]
- 26 Liu D, Lu X, Shen J, Huang S, Wang H. Primary pulmonary meningioma: A case report. *Türk Gogus Kalp Damar Cerrahisi Derg* 2023; **31**: 286-288 [PMID: 37484635 DOI: 10.5606/tgkdc.dergisi.2023.21196]
- 27 Korn D, Bensch K, Liebow AA, Castleman B. Multiple minute pulmonary tumors resembling chemodectomas. *Am J Pathol* 1960; **37**: 641-672 [PMID: 13753180]

- 28 **Li H**, Xue J, Li P, Xu Y, Zheng Z, Li S, Liang N. [Recent Advances and Controversies in Minute Pulmonary Meningothelial-like Nodules]. *Zhongguo Fei Ai Za Zhi* 2023; **26**: 621-629 [PMID: [37752542](#) DOI: [10.3779/j.issn.1009-3419.2023.102.30](#)]
- 29 **Masago K**, Hosada W, Sasaki E, Murakami Y, Sugano M, Nagasaka T, Yamada M, Yatabe Y. Is primary pulmonary meningioma a giant form of a meningothelial-like nodule? A case report and review of the literature. *Case Rep Oncol* 2012; **5**: 471-478 [PMID: [23109924](#) DOI: [10.1159/000342391](#)]
- 30 **Ohashi-Nakatani K**, Shibuki Y, Fujima M, Watanabe R, Yoshida A, Yoshida H, Matsumoto Y, Tsuchida T, Watanabe SI, Motoi N. Primary pulmonary meningioma: A rare case report of aspiration cytological features and immunohistochemical assessment. *Diagn Cytopathol* 2019; **47**: 330-333 [PMID: [30548187](#) DOI: [10.1002/dc.24126](#)]
- 31 **Ionescu DN**, Sasatomi E, Aldeeb D, Omalu BI, Finkelstein SD, Swalsky PA, Yousem SA. Pulmonary meningothelial-like nodules: a genotypic comparison with meningiomas. *Am J Surg Pathol* 2004; **28**: 207-214 [PMID: [15043310](#) DOI: [10.1097/00000478-200402000-00008](#)]
- 32 **Tao L**, Chen Y, Huang Q, Yong J, Yan S, Huang Y. Constant expression of somatostatin receptor 2a in minute pulmonary meningothelial-like nodules. *J Clin Pathol* 2019; **72**: 525-528 [PMID: [31208995](#) DOI: [10.1136/jclinpath-2019-205913](#)]
- 33 **Rushing EJ**, Bouffard JP, McCall S, Olsen C, Mena H, Sandberg GD, Thompson LD. Primary extracranial meningiomas: an analysis of 146 cases. *Head Neck Pathol* 2009; **3**: 116-130 [PMID: [19644540](#) DOI: [10.1007/s12105-009-0118-1](#)]
- 34 **Gaffey MJ**, Mills SE, Askin FB. Minute pulmonary meningothelial-like nodules. A clinicopathologic study of so-called minute pulmonary chemodectoma. *Am J Surg Pathol* 1988; **12**: 167-175 [PMID: [2830799](#) DOI: [10.1097/00000478-198803000-00001](#)]
- 35 **Kraushaar G**, Ajlan AM, English JC, Müller NL. Minute pulmonary meningothelial-like nodules: a case of incidentally detected diffuse cystic micronodules on thin-section computed tomography. *J Comput Assist Tomogr* 2010; **34**: 780-782 [PMID: [20861786](#) DOI: [10.1097/RCT.0b013e3181e47c28](#)]
- 36 **Mizutani E**, Tsuta K, Maeshima AM, Asamura H, Matsuno Y. Minute pulmonary meningothelial-like nodules: clinicopathologic analysis of 121 patients. *Hum Pathol* 2009; **40**: 678-682 [PMID: [19144385](#) DOI: [10.1016/j.humpath.2008.08.018](#)]
- 37 **Zhao Z**, Sun K, Yan T, Wei R, Guo W. Multiple primary tumors: a case report and review of the literature. *BMC Musculoskelet Disord* 2020; **21**: 394 [PMID: [32571290](#) DOI: [10.1186/s12891-020-03426-8](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

