

Case 323: Minute Pulmonary Meningothelial-like Nodules

Furkan Ufuk, MD¹ • Emel Kilicarslan, MD • Ferda Bir, MD • Goksel Altinisik, MD

From the Departments of Radiology (F.U.), Pathology (E.K., F.B.), and Pulmonology (G.A.), University of Pamukkale, Denizli, Turkey. Received October 3, 2022; revision requested November 29; revision received February 6, 2023; accepted February 10. Address correspondence to F.U. (email: furkan.ufuk@hotmail.com).

¹ Current address: Department of Radiology, University of Chicago, 5841 S Maryland Ave, Chicago, IL 60637

Conflicts of interest are listed at the end of this article.

Radiology 2024; 310(3):e222512 • <https://doi.org/10.1148/radiol.222512> • Content code: CH • © RSNA, 2024

History A 70-year-old woman with a 6-year history of asthma, a 12-year history of diabetes mellitus, and who did not smoke presented to the pulmonology clinic with dyspnea and cough. Chest CT performed 5 years earlier for similar symptoms revealed multiple pulmonary nodules. However, she was lost to follow-up before the work-up was concluded. Otherwise, her medical history was unremarkable. Family history included maternal endometrial cancer. Physical examination revealed partial oxygen saturation of 98%, respiratory rate of 18 breaths per minute, and heart rate of 77 beats per minute. Her breath sounds and other systemic findings were normal. Pulmonary function test results were as follows: forced expiratory volume in 1 second, 108% predicted (normal range, 80%–120%); total lung capacity, 72% predicted (normal range, 80%–120%); forced vital capacity, 101% predicted (normal range, 80%–120%); diffusing capacity for carbon monoxide, 69% predicted (normal range, 60%–120%); and forced midexpiratory flow, 85% predicted (normal range, 40%–160%). Complete blood count, erythrocyte sedimentation rate, C-reactive protein level, rheumatoid factor, and antinuclear antibody levels were within normal limits. The patient underwent volumetric thin-section CT of the chest using a multidetector CT scanner (Ingenuity Core 128; Philips Medical Systems) to evaluate lung nodules and pulmonary interstitium.

Part one of this case appeared 4 months previously and may contain larger images.

Imaging Findings

The initial unenhanced chest CT images obtained 5 years earlier showed randomly distributed small (≤ 6 mm) ground-glass nodules (Fig 1). High-resolution CT of the chest revealed numerous small ground-glass nodules with a random distribution in both lungs. Some of the nodules showed central lucencies with a ringlike appearance (Cheerio sign) (Fig 2) (1). The pulmonary nodules were essentially unchanged when compared with the chest CT images 5 years earlier. There was no mediastinal or hilar lymphadenopathy. The patient underwent a surgical lung biopsy. Histopathology revealed numerous bland spindle cell proliferations expanding the alveolar septa and resembling meningothelial cells with a whorl formation. Immunohistochemical staining was positive for progesterone receptors and epithelial membrane antigen but negative for synaptophysin (Fig 3). On the basis of the radiologic and histopathologic findings, the patient was diagnosed with minute pulmonary meningothelial-like nodules (MPMNs) and diffuse pulmonary meningotheliomatosis. The patient was observed clinically, without any additional treatment recommendations.

Discussion

MPMNs, also known as minute meningothelioid nodules, are a rare pathologic condition identified with increasing frequency and characterized by meningothelial-like cellular proliferation within the lung interstitium (2,3). The characteristic CT findings of MPMNs include multiple, small, and randomly distributed ground-glass nodules with central lucencies (ringlike appearance) (3–6).

The etiopathogenesis of MPMNs is unknown. It usually affects nonsmoking women in their sixth or seventh decade of life, as in the present case (4,5). MPMNs are usually incidentally discovered in surgical or autopsy lung specimens. Patients with MPMNs may be asymptomatic or may present with dyspnea or cough (4,5). MPMNs should be considered in the differential diagnosis of multiple pulmonary nodules (6), which includes pulmonary Langerhans cell histiocytosis (PLCH), multifocal lung adenocarcinoma, metastases, rheumatoid nodule, granulomatous polyangiitis, multifocal micronodular pneumocyte hyperplasia (MMPH), diffuse idiopathic pulmonary neuroendocrine cell hyperplasia (DIPNECH), nonfibrotic hypersensitivity pneumonitis, and cavitary sarcoidosis (6,7).

On the basis of clinical findings alone, it can be challenging to distinguish these causes of multiple pulmonary nodules. Thin-section CT of the chest can help in diagnosing MPMN. Histopathologic examination and immunohistochemical staining are required to establish the definitive diagnosis.

PLCH can also cause multiple cavitary pulmonary nodules. However, it usually affects young male smokers and shows a predilection for the upper and mid lung, while the lung bases are often spared (8). Moreover, PLCH is usually progressive. Therefore, the long-term stability seen in this case is inconsistent with PLCH.

Persistent multiple ground-glass nodules on follow-up CT scans may be associated with early-stage adenocarcinoma of the lung (6). While it can grow slowly, 5-year stability is inconsistent with malignancy. Similarly, pulmonary nodules associated with metastasis, rheumatoid




Figure 1: Axial unenhanced chest CT images obtained 5 years before current presentation. Selected images from (A) apices, (B) arcus aorta, (C) lingular bronchus, and (D) basal lung levels using lung window settings (16×1.5 -mm collimation, 120-kV tube voltage, 130-mA tube current, 512×512 matrix, 1.5-mm section thickness). Images were also reconstructed with 1.0-mm section thickness. There are multiple, randomly distributed small (≤ 6 mm) ground-glass nodules. Note that some nodules include central lucencies (ringlike appearance) (arrows).




Figure 2: Axial unenhanced thin-section chest CT images obtained at current presentation. Selected images from (A) apices, (B) arcus aorta, (C) lingular bronchus, and (D) lower lung using lung window settings (64×0.625 -mm collimation, 100-kV tube voltage, 350-mA tube current, 512×512 matrix, and 1.0-mm section thickness). Images were also reconstructed with 0.5-mm section thickness using a high-spatial-frequency algorithm. There are multiple, randomly distributed small (≤ 6 mm) ground-glass nodules. Again, some of the nodules include central lucencies (ringlike appearance) (arrows).

nodule, and granulomatous polyangiitis are not expected to remain stable for 5 years (6). Moreover, the patient had normal laboratory findings inconsistent with rheumatoid arthritis, granulomatous polyangiitis, or infection.

MMPH is a rare cause of multiple pulmonary nodules and is seen frequently in patients with the tuberous sclerosis complex. Histopathologically, MMPH is characterized by benign hamartomatous proliferation of type II pneumocytes, often occurring concurrently with pulmonary lymphangioleiomyomatosis (8). Chest CT shows randomly distributed, well-defined small ground-glass nodules and thin-walled, well-defined, scattered round lung cysts. Although pulmonary nodules in MMPH can be stable for many years, as in the present case, central lucency is not an expected CT finding (8). Moreover, the present case did not have lung cysts consistent with lymphangioleiomyomatosis at CT and did not have a clinical history of tuberous sclerosis complex.

DIPNECH is a disease of unknown origin characterized by abnormal proliferation of pulmonary neuroendocrine cells secreting hormones, causing airway inflammation and fibrosis, which may result in constructive bronchiolitis (9). In patients with DIPNECH, uniform nodules are seen throughout both lungs and have well-defined margins on chest images. These nodules do not grow or change over time. Moreover, CT usually shows mosaic attenuation due to air trapping with thickened bronchial walls, which is absent in the present case. While DIPNECH is commonly seen in nonsmoking middle-aged women, central lucency is not an expected CT finding in DIPNECH nodules (9).

Nonfibrotic hypersensitivity pneumonitis (HP), previously known as subacute HP or extrinsic allergic alveolitis, is characterized by an immune-mediated inflammation of the bronchi and peribronchiolar lung tissue caused by inhalation of organic or nonorganic allergens (10). Histopathologically, HP is characterized by poorly defined noncaseating granulomas, cellular bronchiolitis, and giant cells in the pulmonary interstitium or alveoli. Radiologically, HP is characterized by small (<5 mm) centrilobular ground-glass nodules with indistinct borders, mosaic attenuation with air trapping, and patchy areas of ground-glass opacity. Ground-glass nodules in the nonfibrotic HP may contain a small central lucency caused by air in a patent bronchiole and can mimic MPMN radiologically (10). However, unlike MPMN, in nonfibrotic HP, the nodules are smaller and more numerous, and the subpleural lung is spared. While patients with MPMN are generally asymptomatic, nonfibrotic patients with HP usually have signs of small airway disease, and CT usually shows evidence of air trapping.

Pulmonary cavitory sarcoidosis (PCS) is a rare manifestation of sarcoidosis. The prevalence of PCS has been reported as 2.2%, and cavitory lesions are usually seen in patients with severe and active sarcoidosis (7). Radiologically, PCS is generally characterized by thin-walled, multiple, and multiloculated cavitory lesions with the upper lobe predominance, and rarely can septae inside the cavity be found. In PCS cases, the pericavitory space is often affected by nodules, micronodules, ground-glass opacity, consolidation, or fibrosis (7). However, the lung parenchyma surrounding the nodules was normal in the present case. Hours et al (7) reported the median diameter of cavitory lesions in patients with PCS was 20 mm (range, 11–100 mm). However, unlike PCS, the nodules in the present case were much smaller and did not contain apparent cavitation.

In conclusion, this patient had nonspecific respiratory symptoms and randomly distributed ground-glass nodules with central lucencies that were stable over 5 years, suggesting a benign process. The CT appearance and long-term stability favored a diagnosis of MPMNs, confirmed at histopathology. Thin-section CT of the chest helps in diagnosing MPMN and its differentials.




Figure 3: Right lower lobe lung biopsy specimen. (A) Photomicrograph shows numerous foci of cellular proliferation expanding the alveolar septa. (Hematoxylin-eosin stain; original magnification, $\times 20$.) (B) Photomicrograph shows numerous cellular proliferations expanding the alveolar septa and resembling meningothelial cells with a streaming arrangement in syncytial sheets and whorl formation (arrows). (Hematoxylin-eosin stain; original magnification, $\times 100$.) (C) Immunohistochemical staining for epithelial membrane antigen shows positive staining (arrowheads). (Original magnification, $\times 20$.) (D) Immunohistochemical staining for progesterone receptors shows positive staining. (Original magnification, $\times 100$.)

Disclosures of conflicts of interest: E.U. No relevant relationships. E.K. No relevant relationships. F.B. No relevant relationships. G.A. No relevant relationships.

References

1. Chou SH, Kicska G, Kanne JP, Pipavath S. Cheerio sign. *J Thorac Imaging* 2013;28(1):W4.
2. Mizutani E, Tsuta K, Maeshima AM, Asamura H, Matsuno Y. Minute pulmonary meningothelial-like nodules: clinicopathologic analysis of 121 patients. *Hum Pathol* 2009;40(5):678–682.
3. Huang EC, Zhang Y, Bishop JW, Gandour-Edwards RF, Afify AM. Diffuse pulmonary meningotheliomatosis: A diagnostically challenging entity on fine-needle aspiration cytology. *Diagn Cytopathol* 2015;43(9):727–730.
4. Yun G, Huang T, O'Dwyer D, Chugtai A, Agarwal P. Diffuse pulmonary meningotheliomatosis. *Clin Imaging* 2021;70:111–113.
5. Fadl SA, Pillappa R, Parker MS. Minute Pulmonary Meningothelial-like Nodules. *Radiol Cardiothorac Imaging* 2021;3(5):e210219.
6. Ghosh S, Mehta AC, Abuquyyas S, Raju S, Farver C. Primary lung neoplasms presenting as multiple synchronous lung nodules. *Eur Respir Rev* 2020;29(157):190142.
7. Hours S, Nunes H, Kambouchner M, et al. Pulmonary cavitory sarcoidosis: clinico-radiologic characteristics and natural history of a rare form of sarcoidosis. *Medicine (Baltimore)* 2008;87(3):142–151.
8. Wang MX, Segaran N, Bhalla S, et al. Tuberous Sclerosis: Current Update. *RadioGraphics* 2021;41(7):1992–2010.
9. Little BP, Junn JC, Zheng KS, et al. Diffuse Idiopathic Pulmonary Neuroendocrine Cell Hyperplasia: Imaging and Clinical Features of a Frequently Delayed Diagnosis. *AJR Am J Roentgenol* 2020;215(6):1312–1320.
10. Fernández Pérez ER, Travis WD, Lynch DA, et al. Diagnosis and Evaluation of Hypersensitivity Pneumonitis: CHEST Guideline and Expert Panel Report. *Chest* 2021;160(2):e97–e156.

Congratulations to the 59 individuals and nine resident groups that submitted the most likely diagnosis (minute pulmonary meningothelial-like nodules) for Diagnosis Please, Case 323. The names and locations of the individuals and resident groups, as submitted, are as follows:

Individual responses

Gholamali Afshang, MD, *Tinley Park, IL*
 Hiroaki Arakawa, MD, *Shimotsuga-Gun, Japan*
 Firat Atak, MD, *Ankara, Turkey*
 Shin-Ichi Cho, *Tokyo, Japan*
 Yves-Sebastien Cordoliani, MD, *Rutali, France*
 Mustafa Kemal Demir, MD, *Istanbul, Turkey*
 Seyed A. Emamian, MD, PhD, *Bethesda, MD*
 Francis T. Flaherty, MD, *Danbury, CT*
 Toshihiro Furuta, MD, PhD, *Minato-Ku, Japan*
 Taku Gomi, *Tokyo, Japan*
 Vasco Goncalves-Matoso, MD, *Morges, Switzerland*
 Harish Gudi, MD, MBBS, *Tucson, AZ*
 Heidi S. Hagen-Kersten, MD, *Amotfors, Sweden*
 Yuki Hayashi, *Nagoya, Japan*
 Christoph Hefel, *Feldkirch, Austria*
 Sodai Hoshiai, MD, PhD, *Tsukuba, Japan*
 Kengo Ikejima, MD, *Tokyo, Japan*
 Akitoshi Inoue, MD, PhD, *Otsu, Japan*
 Koki Kato, MD, *Utsunomiya, Japan*
 Takao Kiguchi, MD, PhD, *Ichinomiya, Japan*
 Takuji Kiryu, MD, PhD, *Gifu, Japan*
 Jakob Kist, MD, PhD, *Naarden, Netherlands*
 Osamu Kizu, MD, *Otsu, Japan*
 Jeffrey S. Klein, MD, *Williston, VT*
 Ryo Kurokawa, MD, PhD, *Tokyo, Japan*
 Naganathan B. Mani, MD, *Chesterfield, MO*
 Maria D. Martin, MD, *Madison, WI*
 Satoshi Matsushima, MD, *Tokyo, Japan*
 Kyoko Nagai, MD, *Yokohama, Japan*
 Moto Nakaya, MD, *Tokyo, Japan*
 Ryan Narbutas, MD, *Norwalk, CT*
 Tammam N. Nehme, MD, *Hinsdale, IL*
 Michael D. Orsi, MD, *San Antonio, TX*
 Tatsushi Oura, *Osaka City, Japan*

Hande Ozen Atalay, MD, *Istanbul, Turkey*
 Mayur Pankhania, *Rajkot, India*
 Diogo L. Pinheiro, MD, *Curitiba, Brazil*
 Jip F. Prince, MD, PhD, *Utrecht, Netherlands*
 Kazuhiro Saito, MD, *Tokyo, Japan*
 Akihiko Sakata, MD, PhD, *Akita, Japan*
 Yusuke Sakurai, *Nagoya, Japan*
 Yoshinao Sato, MD, PhD, *Tokyo, Japan*
 Taro Shimono, MD, PhD, *Osaka, Japan*
 Yuki Shinohara, MD, PhD, *Akita, Japan*
 Ichiro Shirouzu, MD, *Tokyo, Japan*
 Evan S. Siegelman, MD, *Media, PA*
 Liz J. Silverstone, MBChB, *Sydney, Australia*
 Ken Simmons, MD, *Sydney, Australia*
 Taku Tajima, MD, PhD, *Minato-Ku, Japan*
 Taro Takeda, MD, *Hashima-Gun, Japan*
 Takashi Tanaka, MD, PhD, *Okayama, Japan*
 Yuki Tashiro, MD, *Yokohama, Japan*
 Suzune Tsukamoto, MD, *Amagasaki-Shi, Japan*
 Meric Tuzun, MD, *Ankara, Turkey*
 Yusuke Utsunomiya, *Osaka, Japan*
 Christopher P. Vittore, MD, *Belvidere, IL*
 Yukari Wakabayashi, MD, *Tokyo, Japan*
 Fumiko Yagi, MD, *Shinjuku-Ku, Japan*
 Takayuki Yamamoto, MD, PhD, *Kyoto, Japan*

Resident group responses

Amsterdam UMC Radiology Residents, *Amsterdam, Netherlands*
 Grupo de Residentes CT Scanner-INCICH, *Mexico City, Mexico*
 Hospital De Santa Maria Radiology Residents, *Lisbon, Portugal*
 Hospital of the University of Pennsylvania Radiology Residents, *Philadelphia, PA*
 Mater Dei Hospital Radiology Residents, *Msida, Malta*
 Mie University Hospital Radiology Residents, *Tsu, Japan*
 Norwalk Hospital Radiology Residents, *Norwalk, CT*
 Virginia Commonwealth University Health System Radiology Residents, *Richmond, VA*
 Western University Radiology Residents, *London, Canada*