

**WEIGHING MATRICES: GENERALIZATIONS, RELATED  
CONFIGURATIONS**

**THOMAS PENDER  
BACHELOR OF SCIENCE, UNIVERSITY OF LETHBRIDGE, 2020**

A thesis submitted  
in partial fulfillment of the requirements for the degree of

**MASTER OF SCIENCE**

in

**MATHEMATICS**

Department of Mathematics and Computer Science  
University of Lethbridge  
LETHBRIDGE, ALBERTA, CANADA

©Thomas Pender, 2022



WEIGHING MATRICES: GENERALIZATIONS, RELATED  
CONFIGURATIONS

THOMAS PENDER

Date of Defence: June 22, 2022

Dr. H. Kharaghani Supervisor	Professor	Ph.D.
---------------------------------	-----------	-------

Dr. W. Holzmann Examination Committee Member	Professor	Ph.D.
---	-----------	-------

Dr. M. Walton Examination Committee Member	Professor	Ph.D.
---	-----------	-------

Dr. J. Sheriff Chair, Examination Committee	Professor	Ph.D.
--	-----------	-------



## **DEDICATION**

To my wife and son who changed everything.



## **ABSTRACT**

It is the purpose of this thesis to explore the relationships that exist between weighing matrices, including their generalizations, and various other combinatorial configurations.

Principally, it will be shown that any balanced generalized weighing matrix with entries from a finite abelian group is equivalent to the existence of families of commutative association schemes.

Additionally, novel constructions of related combinatorial configurations are presented such as balancedly splittable orthogonal designs and new families of balanced weighing matrices.





## **ACKNOWLEDGEMENTS**

I would like to thank Dr. H. Kharaghani for his remarkable teaching prowess and his singular patience. There can be no doubt that if I have found any success during my studies, it is precisely because of my teacher, Hadi.

Furthermore, thanks are owed to Dr. S. Suda who, with an abundance of charity, agreed to include me in his collaboration with Dr. H. Kharaghani.

The author's correspondence with the aforementioned researchers has proven to be most valuable. Indeed, their patience in allowing me to "catch-up" in order to provide my own insights and contributions in equal measure has been indispensable.

Finally, I would like to recognize the Department of Mathematics and Computer Science of the University of Lethbridge. Truly, here is a group of talented individuals with a passion for teaching and sharing their expertise.



# Contents

<b>Dedication</b>	<b>i</b>
<b>Abstract</b>	<b>iii</b>
<b>Acknowledgements</b>	<b>v</b>
<b>List of Symbols</b>	<b>xi</b>
<b>Chapter 1. Introduction</b>	
§1. Balanced Incomplete Block Designs . . . . .	1
1.1. Definition and Necessary Conditions . . . . .	1
1.2. Related Configurations . . . . .	3
1.3. Isomorphisms of Designs . . . . .	5
§2. Error-Correcting Codes . . . . .	6
2.1. Definitions . . . . .	6
2.2. The Hamming and Simplex Codes . . . . .	8
§3. Weighing Matrices and Their Generalizations . . . . .	9
3.1. Weighing Matrices . . . . .	9
3.2. Generalized Bhaskar Rao Designs . . . . .	10
3.3. Isomorphisms of BGW matrices . . . . .	11
3.4. Orthogonal Designs . . . . .	12
3.5. Sequences and Circulants . . . . .	13

§4. Association Schemes . . . . .	16
4.1. Defintion . . . . .	16
4.2. Adjacency Algebras . . . . .	17
4.3. Intersection Matrices . . . . .	19
Notes . . . . .	20

## **Chapter 2. Balancedly Splittable Orthogonal Designs**

§5. Balancedly Splittable Hadamard Matrices . . . . .	21
5.1. Definition . . . . .	21
5.2. Equiangular Lines . . . . .	23
5.3. Constructions . . . . .	24
§6. Balancedly Splittable Orthogonal Designs . . . . .	27
6.1. Definition . . . . .	28
6.2. Unfaithful Constructions . . . . .	29
6.3. Faithful Construction . . . . .	34
§7. Related Configurations . . . . .	41
7.1. Quasi-Symmetric BIBDs . . . . .	41
7.2. Equiangular Tight Frames . . . . .	43
7.3. Unbiased Orthogonal Designs . . . . .	47
Notes . . . . .	49

## **Chapter 3. A New Family of Balanced Weighing Matrices and Association Schemes**

§8. Generalized Kronecker Product . . . . .	51
8.1. Definitions . . . . .	52
8.2. A First Application: Block designs . . . . .	54
8.3. A Second Application: Bkaskar Rao designs . . . . .	56
§9. A New Family of Balanced Weighing Matrices . . . . .	57

9.1.	Lemmata . . . . .	57
9.2.	Construction . . . . .	61
§10.	Weighing Matrices and Association Schemes . . . . .	63
10.1.	Adjacency Matrices . . . . .	64
10.2.	Character Tables . . . . .	66
	Notes . . . . .	69
 <b>Chapter 4. A Unified Construction of Weighing Matrices</b>		
§11.	A General Method . . . . .	71
11.1.	A First Application: Weighing matrices . . . . .	72
11.2.	A Second Application: Block designs . . . . .	75
§12.	BGW Matrices and Association Schemes . . . . .	78
12.1.	Adjacency Matrices . . . . .	78
12.2.	Character Tables . . . . .	81
	Notes . . . . .	86
 <b>Appendix</b>		<b>87</b>
 <b>Bibliography</b>		<b>97</b>
 <b>Index</b>		<b>101</b>



## LIST OF SYMBOLS

$\mathbf{N}$	The additive semigroup of natural numbers $\{0, 1, 2, \dots\}$ .
$\mathbf{Z}$	The ring of integers $\{0, \pm 1, \pm 2, \dots\}$ .
$\mathbf{Q}$	The rational number field $\{\frac{n}{m} : n, m \in \mathbf{Z}\}$ .
$\mathbf{R}$	The real number field.
$\mathbf{C}$	The complex number field $\{x + y\sqrt{-1} : x, y \in \mathbf{R}\}$ .
$\mathbf{Z}/n\mathbf{Z}$	The integers modulo $n$ $\{\bar{0}, \dots, \overline{n-1}\}$ .
$\text{GF}(q)$	The Galois field of $q$ elements.
$\mathbf{D} = (X, \mathcal{B})$	A balanced incomplete block design with point set $X$ and block set $\mathcal{B}$ .
$\text{BIBD}(v, k, \lambda)$	A balanced incomplete block design with associated parameters.
$\mathfrak{C}(\mathbf{D})$	Complementary design of $\mathbf{D}$ .
$\mathfrak{R}(\mathbf{D})$	Residual design of $\mathbf{D}$ .
$\mathfrak{D}(\mathbf{D})$	Derived design of $\mathbf{D}$ .
$A(\mathbf{D})$	Incidence matrix of the design $\mathbf{D}$ .
$[n, k, d, w]_q$ -code	A linear code in $\text{GF}(q^n)$ of dimension $k$ , minimum distance $d$ , and minimum weight $w$ .
$A_q(n, d, w)$	The maximum size of any $(n, M, d, w)_q$ -code, i.e. $M \leq A_q(n, d, w)$ .
$\mathbf{W}(v, k)$	A weighing matrix of order $v$ and weight $k$ .
$\mathbf{BW}(v, k; n)$	A Butson weighing matrix of order $v$ and weight $k$ over the $n$ -th roots of unity.
$\text{GBRD}(v, k, \lambda; G)$	A generalized Bhaskar Rao design over the finite group $G$ with associated parameters.
$\text{BGW}(v, k, \lambda; G)$	A balanced generalized weighing matrix of order $v$ , weight $k$ , and index $\lambda/ G $ , over the finite group $G$ .
$\mathfrak{X} = (X, \mathcal{R})$	An association scheme with point set $X$ and relations $\mathcal{R}$ .
$\mathcal{S}_{q,n}$	A generalized simplex code.
$\mathcal{H}_{q,n}$	A generalized Hamming code.
$A \otimes B$	The Kronecker product of $A$ by $B$ .

# 1

## Introduction

This preliminary chapter introduces the objects with which we will be working. We intend to move briskly, so only references to proofs and more in-depth discussions will be given, and moreover, examples will be reserved for later chapters. The following sections will define in turn: balanced incomplete block designs, error-correcting codes, weighing matrices and their generalizations, and finally association schemes.

### §1. Balanced Incomplete Block Designs

This section presents some basic definitions and results about block designs that will be used throughout this work. Particular emphasis will be placed on matrix representations of such objects.

\* \* \*

*1.1. Definition and Necessary Conditions.* A block design is simply a collection of subsets of a finite point set such that the points are regularly distributed



## 1. Introduction

amongst the subsets in question. There are numerous directions to take with this vague understanding, but we only require the following.

**1.1. Definition.** Let  $X$  be a set of order  $v$ , called the set of varieties; and let  $\mathcal{B} \subset \binom{X}{k}$ , called the set of blocks, have order  $b$ . The ordered pair  $\mathbf{D} = (X, \mathcal{B})$  is a *balanced incomplete block design* (henceforth BIBD) if there is a positive integer  $\lambda$  such that each 2-subset of varieties appears in  $\lambda$  blocks of  $\mathcal{B}$ .

The conditions placed on a finite set and a collection of its subsets in order to form a BIBD are quite strong, and we have at once the following result which can be shown using the elementary techniques of double counting (see Cameron, 1994, Chapter 2 for an abstract discussion).

**1.2. Proposition.** Let  $\mathbf{D} = (X, \mathcal{B})$  be a BIBD with  $|X| = v$ , and  $\mathcal{B} \subset \binom{X}{k}$  for which  $|\mathcal{B}| = b$ . Then:

(1.2.a) Every point of  $X$  occurs in  $r = \frac{\lambda(v-1)}{k-1}$  blocks, and

(1.2.b) there are  $b = \frac{vr}{k} = \frac{\lambda v(v-1)}{k(k-1)}$  blocks in  $\mathcal{B}$ .

**1.3. Corollary.** For the parameters  $v, k, \lambda$  of a BIBD, it must hold that

(1.3.a)  $\lambda(v-1) \equiv 0 \pmod{k-1}$ , and

(1.3.b)  $\lambda v(v-1) \equiv 0 \pmod{k(k-1)}$ .

If  $\mathbf{D} = (X, \mathcal{B})$  is a BIBD with the parameters shown above, then we denote this property as  $\text{BIBD}(v, b, r, k, \lambda)$ . As we have seen, however, the parameters  $b$  and  $r$  are expressible in terms of  $v, k$ , and  $\lambda$ ; hence, we will usually shorten the denotation to  $\text{BIBD}(v, k, \lambda)$  whenever no confusion will arise.

Corollary 1.3 imposes some necessary conditions on the parameters of a BIBD. Our next result, due to Fisher (1940), is a strong necessary condition relating the

## 1. Introduction

number of points to the number of blocks of a BIBD, and it has far reaching consequences in the applications of designs to fields like statistics.

This most important result admits several interesting derivations employing techniques ranging from determinants to variance counting <sup>1</sup>).

**1.4. Fisher's Inequality.** Let  $\mathbf{D} = (X, \mathcal{B})$  be a  $\text{BIBD}(v, b, r, k, \lambda)$ . It follows that

$$(1.4.a) \quad b \geq v.$$

The extremal case of Fisher's inequality is naturally very interesting and important. We single this case out thus.

**1.5. Definition.** Let  $\mathbf{D} = (X, \mathcal{B})$  be some  $\text{BIBD}(v, b, r, k, \lambda)$ . If  $v = b$  (equiv.  $k = r$ ), then we say that  $\mathbf{D}$  is a *symmetric* balanced incomplete block design, or simply symmetric.

\* \* \*

**1.2. Related Configurations.** Thus far, we have been thinking of designs strictly as subsets of some finite set. We can, however, broaden our view to include the following tool, and in so doing the theory of linear algebra can be brought to bear on the subject.

**1.6. Definition.** Let  $\mathbf{D} = (\{x_0, \dots, x_{v-1}\}, \{B_0, \dots, B_{b-1}\})$  be a  $\text{BIBD}(v, b, r, k, \lambda)$ , and let  $A$  be a  $v \times b$   $(0, 1)$ -matrix defined by

$$(1.6.a) \quad A_{ij} = \begin{cases} 1 & \text{if } x_i \in B_j, \text{ and} \\ 0 & \text{if } x_i \notin B_j. \end{cases}$$

We call  $A$  the *incidence matrix* of the design.

## 1. Introduction

The next result is immediate. Note that we use  $I_n$  and  $J_n$  to denote the identity matrix and the all ones matrix, respectively, of  $n$  rows and  $n$  columns. Similarly,  $\mathbf{1}_n$  and  $\mathbf{0}_n$  will denote the column with  $n$  ones and the column with  $n$  zeros. For simplicity, the indices will at times be omitted.

**1.7. Proposition.** Let  $\mathbf{D} = (X, \mathcal{B})$  be a BIBD( $v, b, r, k, \lambda$ ), and let  $A$  be a  $v \times b$   $(0, 1)$ -matrix. Then  $A$  is the incidence matrix of the design if and only if the following hold.

$$(1.7.a) \quad AA^t = rI_v + \lambda(J_v - I_v), \text{ and}$$

$$(1.7.b) \quad \mathbf{1}_v^t A = k\mathbf{1}_b^t.$$

As balanced incomplete block designs can more generally be regarded as finite incidence structures <sup>2)</sup>, there can be related a number of further such objects. For our purposes, we will be interested in the following.

**1.8. Definition.** Let  $\mathbf{D} = (X, \mathcal{B})$  be a BIBD, and let  $B \in \mathcal{B}$ . Then the *complement design*  $\mathfrak{C}(\mathbf{D})$  is the pair  $(X, \binom{X}{k} \setminus \mathcal{B})$ . If  $\mathbf{D}$  is symmetric, we have that the *derived design*  $\mathfrak{D}(\mathbf{D})$  is the pair  $(B_0, \{B \cap B_0 : B \in \mathcal{B} \text{ and } B \neq B_0\})$ , and the *residual design*  $\mathfrak{R}(\mathbf{D})$  is the pair  $(X \setminus B_0, \{B - B_0 : B \in \mathcal{B} \text{ and } B \neq B_0\})$ . When convenient, we will simply denote the complement, derived, and residual designs as  $\mathfrak{C}$ ,  $\mathfrak{D}$ , and  $\mathfrak{R}$ , respectively.

The following result shows why, in part, these substructures are interesting.

**1.9. Proposition.** Let  $\mathbf{D} = (X, \mathcal{B})$  be a BIBD( $v, b, r, k, \lambda$ ). Then

$$(1.9.a) \quad \mathfrak{C} \text{ is a BIBD}(v, b, b - r, v - k, b - 2r + \lambda).$$

If  $\mathbf{D}$  is symmetric, then we further have that

$$(1.9.b) \quad \mathfrak{D} \text{ is a BIBD}(k, b - 1, k - 1, \lambda, \lambda - 1), \text{ and}$$

## 1. Introduction

(1.9.c)  $\mathfrak{R}$  is a BIBD( $v - k, b - 1, k, k - \lambda, \lambda$ ).

\* \* \*

**1.3. Isomorphisms of Designs.** We conclude this section by briefly discussing isomorphisms of designs.

**1.10. Definition.** Let  $\mathbf{D}_1 = (X_1, \mathcal{B}_1)$  and  $\mathbf{D}_2 = (X_2, \mathcal{B}_2)$  be two BIBDs with the same parameters, and let  $f : X_1 \rightarrow X_2$  be some bijection. If  $f(\mathcal{B}_1) = \mathcal{B}_2$ , then we say that  $f$  is an *isomorphism* and that the two designs are *isomorphic*. For the case in which  $\mathbf{D}_1 = \mathbf{D}_2$ , we say that  $f$  is an *automorphism*. The collection of all automorphisms of a design  $\mathbf{D}$  forms a group under composition called the *automorphism group* of the design.

In practice, one is usually concerned with the actions of isomorphisms on the incidence matrices of designs. In particular, two BIBD( $v, b, r, k, \lambda$ )s with incidence matrices  $A_1$  and  $A_2$  are isomorphic if and only if there is a permutation matrix  $P$  of order  $v$  and a permutation matrix  $Q$  of order  $b$  such that

$$(1.10.a) \quad PA_1Q = A_2^3).$$

**1.11. Definition.** As nothing essential is changed under the action of an isomorphism, one can then assume that the incidence matrix of a symmetric design has the following form

$$(1.11.a) \quad \begin{pmatrix} \mathbf{0}_{v-k} & A_1 \\ \mathbf{1}_k & A_2 \end{pmatrix}.$$

We will say that such an incidence matrix is in *normal form*.

\* \* \*

## 1. Introduction

### §2. Error-Correcting Codes

In this section, the definitions of linear error-correcting codes will be given. We then move on to consider the famous generalized Hamming and simplex codes. As these are the only family of codes that we require, this section will be brief. The interested reader is referred to the standard references of Huffman and Pless (2003) and MacWilliams and Sloane (1977) for a greater exposition of the subject.

\* \* \*

**2.1. Definitions.** In essence a code is simply a finite collection of words over a given finite alphabet. There are many ways to formalize such a concept; for instance, we may consider a code to be a subset of functions from one finite set into another.

For the purposes at hand, however, we are interested in the case the alphabet is endowed with arithmetic sufficient to form a field in which case we may consider the code to be a linear subspace of an extension of the alphabet.

**2.1. Definition.** Let  $H = (A \mid I_{n-k})$  be an  $(n - k) \times n$  matrix over  $\text{GF}(q)$ . The linear code  $\mathcal{C}$  with parity check matrix  $H$  is given by  $\mathcal{C} = \text{Null}(H) = \{x \in \text{GF}(q^n) : Hx^t = 0\}$ , where the extension  $\text{GF}(q^n)$  is interpreted here as a linear space over  $\text{GF}(q)$ . We say that  $\mathcal{C}$  is a linear  $[n, k]_q$ -code, where clearly  $\dim(\mathcal{C}) = k$ .

We have defined a linear code by its parity check matrix. An equivalent definition is to use the so-called *generator matrix*  $G$ . If  $H = (A \mid I_{n-k})$  is the parity check matrix of the code, then  $G = (I_k \mid -A^t)$ . The code is then given by the linear span of the rows of  $G$ . Note that by construction, it follows that  $GH^t = HG^t = O$ .

## 1. Introduction

The dual of a linear code  $\mathcal{C}$  is given in the usual way by  $\mathcal{C}^\perp = \{x \in \text{GF}(q^n) : xy^t = 0, \text{ for every } y \in \mathcal{C}\}$ . If  $H$  and  $G$  are the parity check and generator matrices of  $\mathcal{C}$ , then  $G$  and  $H$  are the parity check and generator matrices of  $\mathcal{C}^\perp$ , respectively.

In Definition 2.1, there are two parameters explicitly given of a code, namely, the length  $n$  of the codewords and the dimension  $k$  of the linear space consisting of the codewords. Two more fundamental parameters for us are the minimum distance and the minimum weight defined thus.

**2.2. Definition.** Let  $\mathcal{C}$  be a linear  $[n, k]_q$ -code, and let  $x = x_0 \cdots x_{n-1}$  and  $y = y_0 \cdots y_{n-1}$  be any two codewords of  $\mathcal{C}$ . Then:

(2.2.a) The *Hamming weight* of  $x$  is  $\text{wt}(x) = \#\{i : x_i \neq 0\}$ ;

(2.2.b) the *Hamming distance* between  $x$  and  $y$  is  $\text{dist}(x, y) = \#\{i : x_i \neq y_i\}$ ;

(2.2.c) the *minimum weight* of the code is  $\text{wt}(\mathcal{C}) = \min_{x \in \mathcal{C} \setminus \{0\}} \text{wt}(x)$ ; and

(2.2.d) the *minimum distance* of the code is  $\text{dist}(\mathcal{C}) = \min_{\substack{x, y \in \mathcal{C} \\ x \neq y}} \text{dist}(x, y)$ .

If we wish to emphasize the distance  $d = \text{dist}(\mathcal{C})$  of a code, we write  $[n, k, d]_q$ -code.

The Hamming distance can easily be seen to form a metric <sup>4)</sup> on  $\text{GF}(q^n)$ . The next result is then clear (see Hill, 1986, Theorem 1.9).

**2.3. Proposition.** If  $\mathcal{C}$  is any  $[n, k, d]_q$ -code, then

(2.3.a) for every  $x, y \in \mathcal{C}$ , it holds that  $\text{dist}(x, y) = \text{wt}(x - y)$ , hence  $\text{dist}(\mathcal{C}) = \text{wt}(\mathcal{C})$ .

\* \* \*

## 1. Introduction

**2.2. The Hamming and Simplex Codes.** The theory of error-correcting codes began with the seminal paper of Richard W. Hamming (1950) who introduced redundancy for the purpose of error correction. In this paper, he also constructed a most useful family of codes aptly called the *Hamming codes*. The codes constructed by Hamming were binary; however, it is a simple matter to extend the construction to an arbitrary finite field.

**2.4. Definition.** Let  $H$  be the  $n \times (q^n - 1)/(q - 1)$  matrix over  $\text{GF}(q)$  whose columns are representatives of the nonzero 1-dimensional subspaces of the extension field  $\text{GF}(q^n)$  over  $\text{GF}(q)$ . The linear code  $\mathcal{H}_{q,n}$  whose parity check matrix is  $H$  is called a *Hamming code*. The linear code  $\mathcal{S}_{q,n}$  whose generator matrix is  $H$  is called a *simplex code*.

The following result is immediate.

**2.5. Proposition.** Let  $q$  be a prime power,  $n > 1$ , and let  $v = (q^n - 1)/(q - 1)$ . Then:

(2.5.a)  $\mathcal{S}_{q,n}$  is a linear  $[v, n]_q$ -code, and

(2.5.b)  $\mathcal{H}_{q,n}$  is a linear  $[v, v - n]_q$ -code.

It can be shown that  $\mathcal{H}_{q,n}$  is a linear single-error correcting code. In what follows, however, the simplex code will play a central role. The reader may consult Theorem 3.9.27 of Ionin and Shrikhande (2006) for a derivation.

**2.6. Theorem.** Let  $q$  be a prime power,  $n > 1$ , and let  $v = (q^n - 1)/(q - 1)$ . Then a linear  $[v, n]_q$ -code  $\mathcal{C}$  is a  $\mathcal{S}_{q,n}$  code if and only if  $\text{wt}(x) = q^{n-1}$ , for every  $x \in \mathcal{C}$ .

### §3. Weighing Matrices and Their Generalizations

This section focuses on a third combinatorial configuration, namely, weighing matrices together with their generalizations. We begin with a brief look at weighing matrices themselves before moving onto to consider two generalizations. The first generalization is allowing the entries of a weighing matrix to be chosen from a finite group. This idea is then synthesized with the ideas of §1. Finally, we allow the entries of a weighing matrix to be taken from sets of indeterminants and consider the utility of such an approach.

\* \* \*

**3.1. Weighing Matrices.** We begin the following definition.

**3.1. Definition.** Let  $W$  be a  $(-1, 0, 1)$ -matrix of order  $n$ .  $W$  is a *weighing matrix* of order  $n$  and *weight*  $k$  if

$$(3.1.a) \quad WW^t = kI_n.$$

If  $n = k$ , then we say that  $W$  is a *Hadamard matrix*. If  $n - 1 = k$ , then we say that  $W$  is a *conference matrix*. In any event, we write  $W(n, k)$  to denote this property.

There are many useful generalizations of weighing matrices. We will take this up generally in the following sections, but for now we note the following special case. Note that we use  $A^*$  to denote the conjugate (Hermitian) transpose of a complex matrix  $A$ .

**3.2. Definition.** Let  $G = \{\exp(\frac{2\pi im}{p}) : 0 \leq m < p\}$ , and let  $W$  be a  $(0, G)$ -matrix of order  $v$ . We say that  $W$  is a *Butson weighing matrix* of order  $v$  and *weight*  $k$  if

$$(3.2.a) \quad WW^* = kI_n,$$



## 1. Introduction

and we write  $\text{BW}(v, k; p)$  to denote this property.

A result that is useful in studying properties of complex weighing matrices is the following that can be found in Lam and Leung (2000).

**3.3. Lemma.** Let  $p$  be a prime, and let  $\xi$  be a primitive complex  $p$ -th root of unity. Then  $\sum_{i=0}^n a_i \xi^i = 0$  for some  $n < p$  and  $a_i \in \mathbb{N}$  if and only if  $n = p - 1$  and  $a_0 = \dots = a_n$ .

We now move on to consider some generalizations of a weighing matrix.

\* \* \*

**3.2. Generalized Bhaskar Rao Designs.** In the previous section, we defined a weighing matrix as a square matrix over  $\{-1, 0, 1\}$ . We then extended this definition to include those matrices over  $\{0\}$  together with the complex  $p$ -th roots of unity. More generally, we can have weighing matrices over any finite group.

Before we can do this, however, we need to extend the conjugate transpose to group matrices. To accomplish this, let  $A$  be some matrix over a finite group  $G$ , and define  $\bar{A}$  by  $\bar{A}_{ij} = A_{ij}^{-1}$ , that is, the matrix obtained by taking the group inverse of the nonzero entries of  $A$ . Finally, define  $A^* = \bar{A}^t$ . We then have the following.

**3.4. Definition.** Let  $G$  be some finite group, and let  $A$  be a  $v \times b$   $(0, G)$ -matrix such that

$$(3.4.a) \quad AA^* = rI_v + \frac{\lambda}{|G|} \left( \sum_{g \in G} g \right) (J_v - I_v),$$

for some positive integers  $r$  and  $\lambda$ , and such that there are  $k$  non-zero entries in every column. We then say that  $A$  is a *generalized Bhaskar Rao design* (henceforth GBRD), and we write  $\text{GBRD}(v, k, \lambda; G)$  to denote this property. If we need to stress the remaining parameters, then we write  $\text{GBRD}(v, b, r, k, \lambda; G)$ .

## 1. Introduction

It is clear that replacing the nonzero entries of a GBRD with unity furnishes a BIBD. Hence, we see that Fisher's Inequality applies. Again we single out the extremal case of the inequality.

**3.5. Definition.** A *balanced generalized weighing matrix* is a  $\text{GBRD}(v, b, r, k, \lambda; G)$  in which  $v = b$  (equiv.  $k = r$ ). We use the denotation  $\text{BGW}(v, k, \lambda; G)$ . A  $\text{BGW}(v, k, \lambda; G)$  in which  $v = k$  is called a *generalized Hadamard matrix*, and we denote this as  $\text{GH}(G, \lambda)$  where  $\lambda = v/|G|$ . If  $G = \text{EA}(q)$ , the elementary abelian group<sup>5</sup> of order  $q$ , then we write  $\text{GH}(q, \lambda)$  instead.

Using Lemma 3.3, we see that a Butson weighing matrix over prime complex roots of unity forms a BGW precisely in the case that the nonzero entries yield the incidence matrix of a symmetric design.

BGW matrices are quite useful in the construction of other combinatorial designs (see, for example, Ionin and Kharaghani 2003a and 2003b), and satisfy a number of properties. For a detailed discussion of BGW matrices, the interested reader may consult Ionin and Shrikhande (2006).

\* \* \*

**3.3. Isomorphisms of BGW matrices.** As before, we can impose an equivalence on the set of all  $v \times b$   $(0, G)$ -matrices, which will play an important part in what is to come.

**3.6. Definition.** Two  $v \times b$   $(0, G)$ -matrices  $A_1$  and  $A_2$  are said to be *monomially equivalent* if there are monomial  $(0, G)$ -matrices  $P$  and  $Q$  of orders  $v$  and  $b$ , respectively, such that

$$(3.6.a) \quad PA_1Q = A_2 \text{ } ^6).$$

In order to extend normality to BGW matrices, we begin by altering somewhat Definition 3.4 as in Part V of Colbourn and Dinitz (2007).

## 1. Introduction

**3.7. Definition.** let  $G$  be some finite group, and let  $A$  be a  $v \times b$   $(0, G)$ -matrix.

If  $A$  has  $k$  non-zero entries in every column, and if there is an element  $c \in (\frac{\lambda}{|G|}G) \setminus \{1\} \subset \mathbf{Z}[G]$  such that

$$(3.7.a) \quad AA^* = rI_v + c(J_v - I_v),$$

then we say that  $A$  is a  $c$ -GBRD( $v, k, \lambda - 1; G$ ), or a  $c$ -GBRD( $v, b, r, k, \lambda - 1; G$ ) if more precision required.

We can now properly extend the idea of normality to BGW matrices.

**3.8. Definition.** A BGW( $v, k, \lambda; G$ ) is said to be in *normal form* if it has the form

$$(3.8.a) \quad \begin{pmatrix} \mathbf{0}_{v-k} & A_1 \\ \mathbf{1}_k & A_2 \end{pmatrix}.$$

Note that of necessity,  $A_1$  is a GBRD( $v - k, v - 1, k, k - \lambda, \lambda; G$ ) and  $A_2$  is a  $c$ -GBRD( $k, v - 1, k - 1, \lambda, \lambda - 1; G$ ). These have the parameters of the residual and derived designs of a square BIBD( $v, k, \lambda$ ), hence we call them, respectively, a residual GBRD and a derived  $c$ -GBRD.

\* \* \*

**3.4. Orthogonal Designs.** In the previous subsection, we covered one generalization of weighing matrices, namely, we allowed the nonzero enetries to come from a finite group. In this subsection, we pursue another generalization in a different direction. In particular, instead of allowing the nonzero entries to come from some group, we will take the nonzero entries to be indeterminants—real, complex, or quaternary <sup>7</sup>).

We begin with the case of real indeterminants.

## 1. Introduction

**3.9. Definition.** Let  $x_1, \dots, x_u$  be real, commuting indeterminants, and let  $X$  be an  $n \times n$  matrix with entries from  $\{0, \pm x_1, \dots, \pm x_u\}$ . We say that  $X$  is an *orthogonal design* if

$$(3.9.a) \quad XX^t = \left( \sum_i s_i x_i^2 \right) I_n.$$

We say that the orthogonal design is of order  $n$  and type  $(s_1, \dots, s_u)$ , and we write  $X$  is an  $OD(n; s_1, \dots, s_u)$ . If  $\sum_i s_i = n$ , then we say that the OD is *full*.

We now extend Definition 3.9 to the case of complex and quaternary numbers.

**3.10. Definition.** Let  $z_1, \dots, z_u$  be complex (resp. quaternary), commuting indeterminants, and let  $X$  be a matrix of order  $n$  with entries from  $\{0, \varepsilon_1 z_1, \dots, \varepsilon_u z_u\}$  and  $\{\varepsilon_1 z_1^*, \dots, \varepsilon_u z_u^*\}$ , where each  $\varepsilon_t \in \{\pm 1, \pm i\}$  (resp.  $\varepsilon_t \in \{\pm 1, \pm i, \pm j, \pm k\}$ ).

In the event that

$$(3.10.a) \quad XX^* = \left( \sum_i s_i |z_i|^2 \right) I_n,$$

then we say that  $X$  is a *complex* (resp. *quaternary*) orthogonal design of type  $(s_1, \dots, s_u)$ . We write  $X$  is a  $COD(n; s_1, \dots, s_u)$  (resp.  $QOD(n; s_1, \dots, s_u)$ ).

Equivalence of orthogonal designs is the same as for BGW matrices where the monimal matrices have nonzero entries in set  $\{1, i, j, k\}$ , and where we also allow a permutation of symbols.

\* \* \*

**3.5. Sequences and Circulants.** Let  $A$  be an  $n \times n$  matrix with first row  $(a_0, \dots, a_{n-1})$ . Recall that  $A$  is *circulant* if  $A_{ij} = a_{j-i}$ , where the indices are calculated modulo  $n$ . In this way, the entire matrix is determined by its first row; moreover, if  $A$  and  $B$  are two circulants of the same dimension, then  $A^t$ ,  $A+B$ , and  $AB$  are also circulant matrices. Therefore, to effect a study of circulant matrices, we can profitably study sequences.

What we are particularly interested with here is the following.

## 1. Introduction

**3.11. Definition.** Let  $\mathcal{A} = \{A_i\}$  be a finite collection of circulant matrices of the same dimension over a commutative ring  $R$  endowed with an involution  $\cdot^*$ . The collection  $\mathcal{A}$  is said to be *complementary* if

$$(3.11.a) \quad \sum_i A_i A_i^* = aI, \text{ for some } a \in R,$$

where, as usual,  $(m_{ij})^* = (m_{ji}^*)$ .

Note, however, that we can state this in terms of sequences. First, a definition.

**3.12. Definition.** Let  $a_0 = (a_{0,0}, \dots, a_{0,n-1})$  be a sequence in a commutative ring  $R$  with the involution  $\cdot^*$ . The *j-th aperiodic* and *j-th periodic autocorrelations* of the sequence  $a$  are given respectively by

$$(3.12.a) \quad N_j(a) = \sum_{i=0}^{n-j-1} a_{0,i} a_{0,i+j}^*, \text{ and}$$

$$(3.12.b) \quad P_j(a) = \sum_{i=0}^{n-1} a_{0,i} a_{0,i+j}^*, \text{ indices calculated modulo } n.$$

If  $a_1 = (a_{1,0}, \dots, a_{1,n-1}), \dots, a_m = (a_{m,0}, \dots, a_{m,n-1})$  are any other sequence in  $R$ , then  $a_0, \dots, a_m$  are *complementary* if

$$(3.12.c) \quad \sum_i P_j(a_i) = 0, \text{ for every } j \in \{1, \dots, n-1\}.$$

For the case in which  $m = 1$ , we say that we have a *Golay pair*.

We see immediately that  $P_j(a) = N_j(a) + N_{n-j}(a)^*$ ; hence, if  $N_j(a) + N_j(b) = 0$ , for every  $j \in \{1, \dots, n-1\}$ , then  $a$  and  $b$  are complementary. However, vanishing periodic autocorrelations does not in general imply vanishing aperiodic autocorrelations.

The importance of the periodic correlation is given by the fact that if the first row of the circulant  $A$  continues to be  $a = (a_0, \dots, a_{n-1})$ , then the first row of  $AA^*$  is given by  $(\sum_i |a_i|^2, P_{n-1}(a), \dots, P_1(a))$ . So we see that complementary circulants and complementary sequences are one and the same.

## 1. Introduction

Complementary sequences and orthogonal designs are connected in an intimate way; in fact, complementary sequences offer many elegant constructions of ODs. To make the connection precise, we need to allow sequence elements to be indeterminants whether real, complex, or quaternary, and where the involution is taken to be conjugation.

**3.13. Proposition.** Let  $\{z_1, \dots, z_u\}$  be commuting quaternary indeterminants, and let  $a = (a_0, \dots, a_{n-1})$  and  $b = (b_0, \dots, b_{n-1})$  be complementary sequences with entries from  $\{0, \varepsilon_0 z_0, \dots, \varepsilon_u z_u\}$ , where  $\varepsilon_t \in \{\pm 1, \pm i, \pm j, \pm k\}$ , such that  $\sum_i (|a_i|^2 + |b_i|^2) = \sum_i s_i x_i^2$ . Then

$$(3.13.a) \quad \begin{pmatrix} A & B \\ -B^* & A^* \end{pmatrix}$$

is a QOD( $2n; s_1, \dots, s_u$ ), where  $A$  and  $B$  are the circulants with first rows  $a$  and  $b$ , respectively.

The matrix (3.13.a) will feature as a submatrix in our later work where the sequences will be composed of matrices. We will need one further idea before we proceed.

**3.14. Definition.** Let  $a = (a_0, \dots, a_{n-1})$  and  $b = (b_0, \dots, b_{n-1})$  be two sequences over a ring with involution  $\cdot^*$ . The  $j$ -th cross-correlation of  $a$  by  $b$  is given by

$$(3.14.a) \quad C_j(a, b) = \sum_{i=0}^{n-1} a_i b_{i+j}^*, \text{ for each } j \in \{1, \dots, n-1\}.$$

If  $A$  has first row  $a = (a_0, \dots, a_{n-1})$ , and if  $B$  has first row  $b = (b_0, \dots, b_{n-1})$ , then the first row of the circulant  $AB^*$  is  $(\sum_i a_i b_i^*, C_{n-1}(a, b), \dots, C_1(a, b))$ . Note, however, that  $C_j(a, b)$  is not in general equal to  $C_j(b, a)$ . So, care must be taken in extending Proposition 3.13 since amicability is required in maintaining orthogonality when substituting into an OD.

## 1. Introduction

Much more can be said about this most useful topic. The interested reader is may consult Seberry (2017) and Seberry and Yamada (1992) for a wealth of material.

## §4. Association Schemes

This fourth and final preliminary section briefly touches on association schemes, a fundamental abstract object used as a unifying tool across the otherwise disparate fields of combinatorics.

\* \* \*

**4.1. Definition.** We begin with the following definition.

**4.1. Definition.** Let  $X$  be a finite set of  $v$  elements, and let  $\mathcal{R} = \{R_0, \dots, R_d\}$  be a collection of relations on  $X$ . We say that the ordered pair  $\mathfrak{X} = (X, \mathcal{R})$  is an *association scheme* with  $d$  classes whenever the following are satisfied.

$$(4.1.a) \ R_0 = \{(x, x) : x \in X\};$$

$$(4.1.b) \ R_i \cap R_j = \emptyset \text{ and } X \times X \text{ is the disjoint union of } R_0, \dots, R_d;$$

$$(4.1.c) \ R_i^t = R_{i'}, \text{ for some } i' \in \{0, \dots, d\}, \text{ where } R_i^t = \{(x, y) : (y, x) \in R_i\};$$

and

$$(4.1.d) \ \text{Given } (x, y) \in R_k, \text{ the number of } z \in X \text{ such that } (x, z) \in R_i \text{ and } (z, y) \in R_j \text{ is a constant } p_{ij}^k. \text{ We call the } p_{ij}^k, \text{ the } \textit{intersection numbers} \text{ of the scheme.}$$

The scheme  $\mathfrak{X}$  is *commutative* if

$$(4.1.e) \ p_{ij}^k = p_{ji}^k, \text{ for all } i, j, k \in \{0, \dots, d\}.$$

The scheme is *symmetric* if

## 1. Introduction

$$(4.1.f) \ i = i', \text{ for every } i \in \{0, \dots, d\}.$$

We denote  $p_{ii}^0$  as  $k_i$ , the *valency* of the relation  $R_i$ .

Unless otherwise stated, we will assume that the association schemes we are working with are commutative.

\* \* \*

**4.2. Adjacency Algebras.** The importance of association schemes resides in the following equivalent definition.

**4.2. Definition.** Let  $\mathfrak{X} = (X, \mathcal{R})$  be a  $d$ -class association scheme. For  $i \in \{0, \dots, d\}$ , define the  $v \times v$   $(0, 1)$ -matrix  $A_i$  with rows and columns indexed by elements of  $X$  by  $(A_i)_{xy} = 1$  if and only if  $(x, y) \in R_i$ . We call  $A_i$  the *adjacency matrix* of the relation  $R_i$ . Then Definition 4.1 is equivalent to the following.

$$(4.2.a) \ A_0 = I;$$

$$(4.2.b) \ A_0 + \dots + A_d = J;$$

$$(4.2.c) \ A_i^t = A_{i'} \text{ for some } i' \in \{0, \dots, d\}; \text{ and}$$

$$(4.2.d) \ A_i A_j = \sum_k p_{ij}^k A_k, \text{ for every } i, j \in \{0, \dots, d\}.$$

If  $\mathfrak{X}$  is commutative, then

$$(4.2.e) \ A_i A_j = A_j A_i, \text{ for each } i, j \in \{0, \dots, d\}.$$

If  $\mathfrak{X}$  is symmetric, then

$$(4.2.f) \ A_i^t = A_i, \text{ for every } i \in \{0, \dots, d\}.$$

By (4.2.b), the adjacency matrices of the scheme are  $\mathbf{C}$ -linearly independent and generate a subspace  $\mathfrak{U}$  of  $\text{Mat}_v(\mathbf{C})$  of dimension  $d+1$ . By (4.2.d),  $\mathfrak{U}$  is closed under



## 1. Introduction

standard matrix multiplication. We call  $\mathfrak{U}$  the *adjacency algebra* of the association scheme  $\mathfrak{X}$ .

We further have that  $A_i \circ A_j = \delta_{ij} A_i$ . Therefore,  $A_0, \dots, A_d$  also generate a commutative algebra  $\hat{\mathfrak{U}}$  with Schur multiplication for which they are primitive idempotents<sup>8</sup>). We therefore also call the adjacency matrices the *Schur idempotents* of the scheme.

Assume an ordering of  $X = \{x_0, \dots, x_{v-1}\}$ , and take  $e_{x_i}$  to be the standard vector with  $i$ -th position equal to 1 and 0 elsewhere. Take  $V$  to be the Hermitian space with the standard orthonormal basis  $\{e_x : x \in X\}$ .

Since we are assuming commutativity, the adjacency matrices  $A_0, \dots, A_d$  are pairwise commuting, normal matrices. So, they share an eigenbasis, and by the Spectral Theorem for normal matrices,  $V = \bigoplus_{i=1}^r V_i$ , where the  $V_i$  are maximal common eigenspaces.

Since  $J = \sum_i A_i$ , we find the eigenspace corresponding to the eigenvalue  $v$  is spanned by  $\mathbf{1}_v$ , i.e. it is 1-dimensional and hence maximal. It follows that this space is equal to  $V_i$  for some  $i$ . We can assume, then, that  $i = 0$ .

If we take  $E_i$  to be the orthogonal projection  $V \rightarrow V_i$  with respect to the basis  $\{e_x : x \in X\}$ , then we can assume that  $E_0 = |X|^{-1} J$ , and we have  $E_0 + \dots + E_d = I$ . Moreover, there is a unitary matrix  $\Lambda$  such that  $\Lambda^* E_i \Lambda = \text{diag}(0, \dots, 0, \underbrace{1, \dots, 1}_{m_i}, 0, \dots, 0)$ , where we have used  $m_i$  to denote  $\dim(V_i)$ . The numbers  $m_i$ , for  $i \in \{0, \dots, d\}$ , are called the *multiplicities* of the scheme.

It can be shown (see Bannai and Ito, 1984, Theorem 3.1) that the projection matrices  $E_0, \dots, E_d$  are primitive idempotents of  $\mathfrak{U}$  and form a dual basis of  $\mathfrak{U}$ . It follows that there are constants  $P_{ij}$  and  $Q_{ij}$ , for  $i, j \in \{0, \dots, d\}$ , called the *eigenvalues* and *dual-eigenvalues* of the scheme, such that  $A_j = \sum_i P_{ij} E_i$  and  $E_j = |X|^{-1} \sum_i Q_{ij} A_i$ . Using these constants, we form the matrices  $P$  and  $Q$  with  $(i, j)$ -th entry given by  $P_{ij}$  and  $Q_{ij}$ , respectively, and we call these matrices the

## 1. Introduction

*first and second character tables* of the scheme. By what has been said, we have at once that  $PQ = QP = vI$ .

\* \* \*

**4.3. Intersection Matrices.** If we regard left multiplication of  $\mathfrak{U}$  as linear transformations of  $\mathfrak{U}$  and express them in terms of the basis  $\{A_0, \dots, A_d\}$ , then we have an algebra homomorphism  $\mathfrak{U} \rightarrow \text{Mat}_v(\mathbb{C})$  called the *left regular representation* of  $\mathfrak{U}$  with respect to  $\{A_0, \dots, A_d\}$ .

For  $i \in \{0, \dots, d\}$ , define the matrix  $B_i$  by  $(B_i)_{jk} = p_{ij}^k$ , called the *i-th intersection matrix*. It then follows by (4.2.d) that the image of  $A_i$  under the above homomorphism is  $B_i^t$ , whence  $A_i \leftrightarrow B_i^t$  is an isomorphism. Since, however, we are assuming that our schemes are commutative, transposition is an isomorphism; thus,  $A_i \leftrightarrow B_i$  is an isomorphism. If we denote  $\mathfrak{B} = \langle B_0, \dots, B_d \rangle$ , then we have shown  $\mathfrak{U} \simeq \mathfrak{B}$ .

That the intersection matrices and the isomorphism given above are important for us is shown in the next result. First, however, we say that a vector is in *standard form* if the first entry is 1.

**4.3. Theorem.** Let  $\mathfrak{X} = (X, \mathcal{R})$  be a commutative association scheme with  $d$  classes. Let  $v_i = (P_{i0}, \dots, P_{id})$  and  $u_i = (\overline{P_{i0}}/k_0, \dots, \overline{P_{id}}/k_d)$  be the row vectors obtained by standardizing the first row and column of  $P$  and  $Q$ , respectively. Then  $u_i$  is the unique standardized common left eigenvector  $u$  of the matrices  $B_j$  such that  $uB_j = P_{ij}u$ . Similarly,  $v_i^t$  is the common right eigenvector  $v^t$  of the matrices  $B_j$  such that  $B_jv^t = P_{ij}v^t$ .

As stated in Bannai and Ito (1984), by virtue of this result, one can conceivably derive the character tables  $P$  and  $Q$  from the intersection matrices. In particular, if the algebra  $\mathfrak{B}$  is generated by a single  $B_j$ , we can use the single matrix  $B_j$  to calculate the character tables. Following Lemma 2.2.1 of Brouwer et al. (1989),

## 1. Introduction

the authors go on to point out that if some  $B_j$  has  $d + 1$  distinct eigenvalues, then we can similarly use this intersection matrix. Of course, if  $\mathfrak{B}$  is cyclic, then this property is satisfied by the generator matrix.

## Notes

1. This technique is famously used in the justification of Hoffman's co-clique bound (see Brouwer et al., 1989, Proposition 1.3.2). In Cameron (1994) and Cameron and van Lint (1991), it is used to great effect in the context of designs.
2. By an incidence structure is meant a triple  $S = (X, \mathcal{B}, I)$ , where  $I \subseteq X \times \mathcal{B}$ . We say that  $X$  is the point set,  $\mathcal{B}$  the block set, and  $I$  the set of flags. If  $(x, B) \in I$ , then the point  $x$  and block  $B$  are said to be incident. See Batten (1997) and Dembowski (1997) for a standard treatment of incidence. See Beth et al. (1999) for study of incidence in the context of design theory.
3. This is more generally given as the orbits of the action of  $S_v \times S_b$  on the set of binary  $v \times b$  matrices defined by  $A \cdot (P, Q) = P^t A Q$ , where the transposition is necessary in order to properly define a right action. Note that we are using the notation given in the seminal work Cameron (1999).
4. Recall that a metric space is a pair  $(X, \varrho)$ , where  $X$  is a nonempty set, and where  $\varrho : X \rightarrow \mathbf{R}_{\geq 0}$  is a map satisfying, for all  $x, y, z \in X$ , (a)  $\varrho(x, y) \geq 0$  with equality iff  $x = y$ ; (b)  $\varrho(x, y) = \varrho(y, x)$ ; and (c)  $\varrho(x, y) \leq \varrho(x, z) + \varrho(z, y)$ . We then say that  $\varrho$  is a metric.
5. If  $q = p^n$ , for some prime  $p$ , then  $\text{EA}(q) \simeq \underbrace{C_p \times \cdots \times C_p}_n$ .
6. Let  $H$  and  $K \leq S_v$  be finite groups. Recall that the wreath product  $H \wr K$  is the semi direct product  $H^v \rtimes K$  where  $k(h_0, \dots, h_{v-1})k^{-1} = (h_{k^{-1}0}, \dots, h_{k^{-1}(v-1)})$ . Let  $H_1 = G \wr S_v$ , let  $H_2 = G \wr S_b$ , and let  $H = H_1 \times H_2$ . We define an action of  $H$  on the set of all  $v \times b$   $(0, G)$ -matrices by  $A \cdot (P, Q) = P^* A Q$ . The monomially equivalent matrices are composed precisely by the orbits of this action.
7. Recall that the quaternion group  $\{\pm 1, \pm i, \pm j, \pm k\}$  is defined with anticommutative multiplication given by  $i^2 = j^2 = k^2 = -1$ ,  $ij = k, jk = i, ki = j$ . We then take the quaternion algebra to be the  $\mathbf{R}$ -algebra  $\{a_1 + a_i i + a_j j + a_k k : a_1, a_i, a_j, a_k \in \mathbf{R}\}$ . Conjugation is extended to the quaternion algebra by  $(a_1 + a_i i + a_j j + a_k k)^* = a_1 - a_i i - a_j j - a_k k$ .
8. The algebra  $\hat{\mathfrak{U}}$  is in many senses the dual of the algebra  $\mathfrak{U}$ . This is explored in Bannai and Ito (1984), Bannai et al. (2021), and Delsarte (1973).

# 2

## Balancedly Splittable Orthogonal Designs

This is the first chapter constituting the novel results of this work. Here balancedly splittable orthogonal designs are defined and various constructions are presented. These most interesting objects are connected to several objects such as frames and pairs of unbiased orthogonal designs.

### §5. Balancedly Splittable Hadamard Matrices

The results of this section are due to Kharaghani and Suda (2019). We include this material here to evince the fact that the concept of a balancedly splittable orthogonal design is a generalization of previous work and to provide a more intuitive introduction to the concept.

\* \* \*

**5.1. Definition.** Recall that a Hadamard matrix is a weighing matrix  $W(n, n)$  (see §3). These elusive objects have vexed combinatorialists for over a century

## 2. Balancedly Splittable Orthogonal Designs

now. Ever more clever techniques from ever more branches of mathematics are needed in order to construct these objects.

We will consider one such construction in preparation for our study of orthogonal designs. The matrices we will study are the so-called balancedly splittable Hadamard matrices. First, an example.

**5.1. Example.** Consider a Hadamard matrix of order 4 shown below

$$(5.1.a) \quad \begin{pmatrix} + & + & + & + \\ + & + & - & - \\ + & - & + & - \\ + & - & - & + \end{pmatrix}.$$

Label the rows  $h_0, h_1, h_2$ , and  $h_3$ . We then form the block matrix with  $(i, j)$ -th entry given by  $h_j^t h_i$ .

$$(5.1.b) \quad \begin{pmatrix} + & + & + & + & + & + & + & + & + & + & + & + & + & + & + & + \\ + & + & + & + & + & + & + & + & - & - & - & - & - & - & - & - \\ + & + & + & + & - & - & - & - & + & + & + & + & - & - & - & - \\ + & + & + & + & - & - & - & - & - & - & - & - & + & + & + & + \\ + & + & - & - & + & + & - & - & + & + & - & - & + & + & - & - \\ + & + & - & - & + & + & - & - & - & - & + & + & - & - & + & + \\ + & + & - & - & - & - & + & + & + & + & - & - & - & - & + & + \\ + & - & + & - & + & - & + & - & + & - & + & - & + & - & + & - \\ + & - & + & - & + & - & + & - & - & + & - & + & - & + & - & + \\ + & - & + & - & - & + & - & + & + & - & + & - & - & + & - & + \\ + & - & + & - & - & + & - & + & - & + & - & + & - & + & - & + \\ + & - & - & + & + & - & - & + & + & - & - & + & + & - & - & + \\ + & - & - & + & + & - & - & + & - & + & + & - & - & + & + & - \\ + & - & - & + & - & + & + & - & + & - & - & + & - & + & + & - \\ + & - & - & + & - & + & + & - & - & + & + & - & - & + & - & + \end{pmatrix}.$$

The matrix (5.1.b) has the submatrix  $H_1$  which we have shown in bold above. This submatrix has the property that  $H_1^t H_1 = 4I + 4A$ , for some symmetric  $(0, 1)$ -matrix  $A$  with zero diagonal. In particular, there are only two angles that exist between the columns of  $H_1$ .

The above example motivates the following definition.

## 2. Balancedly Splittable Orthogonal Designs

**5.2. Definition.** A Hadamard Matrix  $H$  of order  $n$  is *balancedly splittable* with parameters  $(n, \ell, a, b)$  if  $H$  has an  $\ell \times n$  submatrix  $H_1$  such that

$$(5.2.a) \quad H_1^t H_1 = \ell I + aA + b(J - I - A),$$

for some binary  $(0, 1)$ -matrix  $A$  with zero diagonal.

This definition and the previous example are suggestive. Notably, a connection to sets of biangular lines is inherent in the definition. These connections will be taken up in the following subsections.

\* \* \*

**5.2. Equiangular Lines.** In the previous subsection, it was intimated that balancedly splittable Hadamard matrices were related to collections of biangular lines. By a set of *lines*, we mean a collection of vectors in  $\mathbf{R}^\ell$ , for some  $\ell$ . Given a collection  $\mathcal{L}$  of lines in  $\mathbf{R}^\ell$ , define  $\Xi = \{|\langle u, v \rangle| : u, v \in \mathcal{L} \text{ and } u \neq v\}$ . If  $|\Xi| = 2$ , then we say that  $\mathcal{L}$  is a set of *biangular lines*; while if  $|\Xi| = 1$ , then we say that  $\mathcal{L}$  is a set of *equiangular lines*<sup>9</sup>).

Clearly, if  $H$  is balancedly splittable with respect to the  $\ell \times n$  submatrix  $H_1$ , then Definition 5.2 implies that the columns of  $H_1$  are at most biangular. They are equiangular precisely in the case that  $b = -a$ .

We will require the following proposition due to Delsarte et al. (1977).

**5.3. Proposition.** Let  $\mathcal{L} \subset \mathbf{R}^\ell$  be a set of lines (vectors) such that  $|\langle u, v \rangle| = a$ , for every pair of distinct lines  $u$  and  $v$  in  $\mathcal{L}$ . If  $\ell < a^{-2}$ , then

$$(5.3.a) \quad |\mathcal{L}| \leq \ell(1 - a^2)/(1 - \ell a^2).$$

Using balancedly splittable Hadamard matrices, we can construct optimal sets of equiangular lines.

**5.4. Theorem.** If there exists a balancedly splittable Hadamard matrix with parameters  $(n, \ell, a, -a)$ , then there is an optimal set of equiangular lines in  $\mathbf{R}^\ell$ .

## 2. Balancedly Splittable Orthogonal Designs

**Proof.** Suppose that  $H$  is a balancedly splittable Hadamard matrix with respect to the  $\ell \times n$  submatrix  $H_1$  with parameters  $(n, \ell, a, -a)$ . Take  $\mathcal{L}$  to be the collection of normalized columns of  $H_1$ . Note that  $a^2 \ell^2 = \ell(n - \ell)/(n - 1)$ ; then the absolute value of the inner product between distinct lines in  $\mathcal{L}$  is given by  $a = \sqrt{n - \ell}/\sqrt{\ell(n - 1)}$ ; moreover,  $\ell \leq a$ . The right-hand side of (5.3.a) reduces to  $n$ . We have, therefore, exhibited an optimal set of equiangular lines. ■

In §10, we will pursue this topic again in the more restricted setting of frames.

\* \* \*

**5.3. Constructions.** For the sake of completeness, we make Example 5.1 general.

**5.5. Proposition.** If there exists a Hadamard matrix of order  $n$ , then there exists a balancedly splittable Hadamard matrix with parameters  $(n^2, n, n, 0)$ .

**Proof.** Let  $H$  be a normalized Hadamard matrix of order  $n$ , and label the rows  $h_0 = \mathbf{1}, \dots, h_{n-1}$ . Take  $M$  to be the block matrix defined by  $M_{ij} = h_j^t h_i$ . Then  $M_{ij} M_{kj}^t = (h_j^t h_i)(h_j^t h_k)^t = h_j^t (h_i h_k^t) h_j = O$  whenever  $i \neq k$ , hence  $M$  is a Hadamard matrix of order  $n^2$  (cf Kharaghani, 1985). Take  $M_1$  to be the first block row of  $M$ . Then

$$M_1^t M_1 = \begin{pmatrix} J \\ \mathbf{1}^t h_1 \\ \vdots \\ \mathbf{1}^t h_{n-1} \end{pmatrix} \begin{pmatrix} J & h_2^t \mathbf{1} & \dots & h_{n-1}^t \mathbf{1} \end{pmatrix} = mI \otimes J,$$

and the proof is complete. ■

There are many more constructions presented in Kharaghani and Suda (2019)

## 2. Balancedly Splittable Orthogonal Designs

that the interested reader may consult. For our purposes, however, we present a novel construction. Consider the following example.

**5.6. Example.** Consider the Hadamard matrix of order 2

$$(5.6.a) \begin{pmatrix} + & + \\ + & - \end{pmatrix},$$

and label the rows  $h_0$  and  $h_1$ . Form the matrices  $c_i = h_i^t h_i$ , the so-called *auxiliary matrices*, shown in order below.

$$(5.6.b) \ c_0 = \begin{pmatrix} + & + \\ + & + \end{pmatrix}, c_1 = \begin{pmatrix} + & - \\ - & + \end{pmatrix}.$$

Take  $K = H \otimes H$ , labelling the rows as  $k_0, k_1, k_2$ , and  $k_3$ . Form the block circulant matrices  $A$  and  $B$  with first rows  $(c_0, c_1, c_1)$  and  $(c_0, c_1, -c_1)$ , shown below.

$$(5.6.c) \ A = \begin{pmatrix} + & + & + & - & + & - \\ + & + & - & + & - & + \\ + & - & + & + & + & - \\ - & + & + & + & - & + \\ + & - & + & - & + & + \\ - & + & - & + & + & + \end{pmatrix}, B = \begin{pmatrix} + & + & + & - & - & + \\ + & + & - & + & + & - \\ - & + & + & + & + & - \\ + & - & + & + & - & + \\ + & - & - & + & + & + \\ - & + & + & - & + & + \end{pmatrix}.$$

Now, form the block matrix  $F = (F_1 \dots F_7)$  by defining  $F_i = k_i^t h_0$

$$(5.6.d) \ F = \begin{pmatrix} + & + & + & + & + & + \\ - & - & + & + & - & - \\ + & + & - & - & - & - \\ - & - & - & - & + & + \end{pmatrix}.$$

Finally, take  $E = F^t$ . We then form the block matrix  $X = \begin{pmatrix} J & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$



## 2. Balancedly Splittable Orthogonal Designs

$$(5.6.e) \quad X = \begin{pmatrix} + & + & + & + & + & + & + & + & - & - & - & - & - & - \\ + & + & + & + & - & - & + & + & - & - & + & + & - & - \\ + & + & + & + & + & + & - & - & - & - & + & + & + & + \\ + & + & + & + & - & - & - & - & + & + & + & + & - & - \\ + & - & + & - & + & + & - & + & - & + & + & - & - & + \\ + & - & + & - & + & + & - & + & - & + & + & - & + & - \\ + & + & - & - & + & - & + & + & + & - & + & + & + & - \\ + & + & - & - & - & + & + & + & - & + & - & + & + & - \\ + & - & - & + & + & - & + & - & + & + & + & - & + & + \\ + & - & - & + & - & + & - & + & + & + & - & + & + & + \\ - & + & - & + & + & + & - & - & + & + & + & - & + & - \\ - & + & - & + & + & + & - & + & + & - & + & - & + & - \\ - & - & + & + & - & + & + & + & - & + & - & + & + & - \\ - & - & + & + & + & - & + & + & - & + & - & + & + & - \\ - & + & + & - & + & - & - & + & + & + & - & + & - & + \\ - & + & + & - & - & + & + & - & + & + & - & + & + & + \end{pmatrix}.$$

The matrix  $X$  is a Hadamard matrix of order 16, and evidently, it is balancedly splittable. Indeed, the submatrix shown in bold above can be used to form a balanced split. Due to the form of  $X$ , however, any one of the following can be used to form a balanced split

$$(5.6.f) \quad \begin{pmatrix} F \\ A \\ B \end{pmatrix}, \begin{pmatrix} -F \\ B \\ A \end{pmatrix}, \begin{pmatrix} E & A & B \end{pmatrix}, \begin{pmatrix} -E & B & A \end{pmatrix}.$$

In any event, the parameters of the splits are  $(16, 6, 2, -2)$ . Therefore, we have optimal sets of equiangular lines.

**5.7. Example.** Beginning with the Hadamard matrix of order 4

$$(5.7.a) \quad \begin{pmatrix} + & + & + & + \\ - & + & - & + \\ - & + & + & - \\ - & - & + & + \end{pmatrix}$$

we obtain the balancedly splittable Hadamard matrix of order 64 as constructed in A1.

## 2. Balancedly Splittable Orthogonal Designs

**5.8. Example.** Of course, we are not limited to real matrices. Beginning with the  $\text{BW}(3, 3; 3)$

$$(5.8.a) \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix},$$

where  $\omega = e^{\frac{2\pi\sqrt{-1}}{3}}$ , we can construct a balancedly splittable  $\text{BW}(36, 36; 6)$  as shown in A2.

This construction can be made perfectly general. Since this result is ultimately a special case of Theorem 6.13, we will not show it explicitly.

**5.9. Theorem.** There is a balancedly splittable Hadamard matrix of order  $4n^2$  with parameters  $(4n^2, 2n^2 - n, n, -n)$  whenever there is a Hadamard matrix of order  $n$ .

**5.10. Corollary.** If there is a Hadamard matrix of order  $n$ , then there is an optimal set of equiangular lines in  $\mathbf{R}^{2n^2-n}$ .

We mention in passing that Kharaghani and Suda (2019) showed that balancedly splittable Hadamard matrices can be used to construct various association schemes. As nothing essentially new about association schemes has been added in our study of orthogonal designs, we will not pursue this topic here and simply refer the reader to the aforementioned article.

## §6. Balancedly Splittable Orthogonal Designs

In this and the following section, the new results of Kharaghani et al. (2021) are presented. Here we define balanced splittability of orthogonal designs and give several constructions. To avoid obfuscation, we given the constructions in terms of real and complex ODs; however, the results are perfectly valid for the more general QODs.

## 2. Balancedly Splittable Orthogonal Designs

\* \* \*

**6.1. Definition.** Generalizing balanced splittability to orthogonal designs presents several difficulties. The various cases are encapsulated in the next definition.

**6.1. Definition.** Let  $X$  be a full QOD( $n; s_1, \dots, s_u$ ).  $X$  is *balancedly splittable* if there is an  $\ell \times n$  submatrix  $X_1$  where one of the following conditions holds. In what follows  $\alpha, \beta \in \{a + ib + jc + kd : a, b, c, d \in \mathbf{R}\}$ .

(6.1.a) The off-diagonal entries of  $X_1^* X_1$  are in the set

$$\{\pm \varepsilon c x_1^{m_1} \cdots x_u^{m_u} x_1^{*m'_1} \cdots x_u^{*m'_u} : m_i, m'_i \in \mathbf{N}, \varepsilon \in \{1, i, j, k\}, c \in \{\alpha, \alpha^*, \beta, \beta^*\}\}.$$

(6.1.b) The off-diagonal entries of  $X_1^* X_1$  are in the set

$$\left\{ \sum_{i=1}^u t_i |x_i|^2 : t_i \in \mathbf{N}, \sum_{i=1}^u t_i = m \right\}$$

or in the set

$$\{\pm \varepsilon c x_1^{m_1} \cdots x_u^{m_u} x_1^{*m'_1} \cdots x_u^{*m'_u} : m_i, m'_i \in \mathbf{N}, \varepsilon \in \{1, i, j, k\}, c \in \{\alpha, \alpha^*, \beta, \beta^*\}\}.$$

(6.1.c) The off-diagonal entries of  $X_1^* X_1$  are in the set

$$\{\pm \varepsilon c \sigma : \varepsilon \in \{1, i, j, k\}, c \in \{\alpha, \alpha^*, \beta, \beta^*\}\},$$

$$\text{where } \sigma = \sum_i s_i |x_i|^2 \text{ (cf. (3.10.a)).}$$

In the first case, the split is *unstable*; in the second, the split is *unfaithfully unstable*; and in the third, the split is *stable*. The term *faithful* is used to describe the first and third cases.

## 2. Balancedly Splittable Orthogonal Designs

From the definition, we see that if  $\alpha$  and  $\beta$  are the same in absolute value, the split corresponds to a set of equiangular lines. Interestingly, we will see that both conditions in (6.1.b) can hold simultaneously.

The next two subsections will present constructions for both the unfaithful, and the faithful case.

\* \* \*

**6.2. Unfaithful Constructions.** The constructions of this section are similar to those presented in Fender et al. (2018) and Pender (2020), and are applicable to real and complex orthogonal designs.

To begin, if  $W$  is a skew-symmetric  $W(q+1, q)$ , then we take  $Q$  to be its core, i.e. the submatrix obtained by deleting the first row and column. Further, we can assume that  $W = \begin{pmatrix} 0 & 1^t \\ -1 & Q \end{pmatrix}$ , hence  $JQ = QJ = O$  and  $Q^2 = J - qI$ . We recursively define the following family of matrices.

$$\mathcal{J}_m = \begin{cases} aJ_1 & \text{if } m = 0, \text{ and} \\ J_q \otimes \mathcal{A}_{m-1} & \text{if } m > 0; \end{cases}$$

$$\mathcal{A}_m = \begin{cases} bJ_1 & \text{if } m = 0, \text{ and} \\ I_q \otimes \mathcal{J}_{m-1} + Q \otimes \mathcal{A}_{m-1} & \text{if } m > 0; \end{cases}$$

where  $a$  and  $b$  are commuting indeterminants.

We require the following lemma.

### 6.2. Lemma.

$$(6.2.a) \quad \mathcal{J}_m \mathcal{A}_m^t = \mathcal{A}_m \mathcal{J}_m^t;$$

$$(6.2.b) \quad \mathcal{J}_m \mathcal{J}_m^t + q \mathcal{A}_m \mathcal{A}_m^t = (q^m a^2 + q^{m+1} b^2) I; \text{ and}$$

$$(6.2.c) \quad \mathcal{J}_1^t \mathcal{J}_1 = qa^2 J, \mathcal{A}_1^t \mathcal{A}_1 = a^2 I + b^2(qI - J), \text{ and } \mathcal{A}_1^t \mathcal{J}_1 = \mathcal{J}_1^t \mathcal{A}_1 = abJ.$$

## 2. Balancedly Splittable Orthogonal Designs

**Proof.** We have  $\mathcal{J}_0 \mathcal{A}_0^t = ab = ba = \mathcal{A}_0 \mathcal{J}_0^t$ . Assume  $\mathcal{J}_{m-1} \mathcal{A}_{m-1}^t = \mathcal{A}_{m-1} \mathcal{J}_{m-1}^t$ . Then

$$\begin{aligned} \mathcal{J}_m \mathcal{A}_m^t &= (J \otimes \mathcal{A}_{m-1})(I \otimes \mathcal{J}_{m-1} + Q \otimes \mathcal{A}_{m-1})^t \\ &= J \otimes \mathcal{A}_{m-1} \mathcal{J}_{m-1}^t + J Q^t \otimes \mathcal{A}_{m-1} \mathcal{A}_{m-1}^t \\ &= J \otimes \mathcal{J}_{m-1} \mathcal{A}_{m-1}^t + Q J \otimes \mathcal{A}_{m-1} \mathcal{A}_{m-1}^t \\ &= (I \otimes \mathcal{J}_{m-1} + Q \otimes \mathcal{A}_{m-1})(J \otimes \mathcal{A}_{m-1})^t, \end{aligned}$$

and (6.2.a) has been shown.

Clearly,  $\mathcal{J}_0 \mathcal{J}_0^t + q \mathcal{A}_0 \mathcal{A}_0^t = a^2 + qb^2$ ; so, assume  $\mathcal{J}_{m-1} \mathcal{J}_{m-1}^t + q \mathcal{A}_{m-1} \mathcal{A}_{m-1}^t = (a^2 + qb^2)I$ . Then

$$\begin{aligned} \mathcal{J}_m \mathcal{J}_m^t + q \mathcal{A}_m \mathcal{A}_m^t &= qJ \otimes \mathcal{A}_{m-1} \mathcal{A}_{m-1}^t + q(I \otimes \mathcal{J}_{m-1} \mathcal{J}_{m-1}^t - Q^2 \otimes \mathcal{A}_{m-1} \mathcal{A}_{m-1}^t) \\ &= qI \otimes (\mathcal{J}_{m-1} \mathcal{J}_{m-1}^t + q \mathcal{A}_{m-1} \mathcal{A}_{m-1}^t) \\ &= qI \otimes (q^{m-1}q^2 + q^m b^2)I \\ &= (q^m a^2 + q^{m+1} b^2)I, \end{aligned}$$

and (6.2.b) is proven.

Finally, (6.2.c) is simply a restatement of the definitions of  $\mathcal{J}_m$  and  $\mathcal{A}_m$ . ■

We can now present the first construction of the novel balancedly splittable ODs.

**6.3. Theorem.** Let  $W$ ,  $\mathcal{A}_m$ , and  $\mathcal{J}_m$  be as above. Define  $X_m = I \otimes \mathcal{J}_m + W \otimes \mathcal{A}_m$ . Then:

(6.3.a)  $X_m$  is an  $\text{OD}(q^m(q+1); q^m, q^{m+1})$ , and

(6.3.b) The matrix  $X_1$  is an unfaithful balancedly splittable  $\text{OD}(q(q+1); q, q^2)$ .

## 2. Balancedly Splittable Orthogonal Designs

**Proof.**  $X_m$  has entries from  $\{\pm a, \pm b\}$ . Observe:

$$\begin{aligned}
X_m X_m^t &= (I \otimes \mathcal{J}_m + W \otimes \mathcal{A}_m)(I \otimes \mathcal{J}_m + W \otimes \mathcal{A}_m)^t \\
&= I \otimes \mathcal{J}_m \mathcal{J}_m^t + W W^t \otimes \mathcal{A}_m \mathcal{A}_m^t \\
&= I \otimes \mathcal{J}_m \mathcal{J}_m^t + q I \otimes \mathcal{A}_m \mathcal{A}_m^t \\
&= I \otimes (\mathcal{J}_m \mathcal{J}_m^t + q \mathcal{A}_m \mathcal{A}_m^t) \\
&= I \otimes (q^m a^2 + q^{m+1} b^2) I \\
&= (q^m a^2 + q^{m+1} b^2) I,
\end{aligned}$$

which shows that  $X_m$  is an  $\text{OD}(q^m(q+1); q^m; q^{m+1})$ . It remains to prove the balanced splittability of the base case.

Take  $Y = (\mathcal{J}_1 \mathcal{A}_1 \dots \mathcal{A}_1)$ , the first block row of  $X_1$ . Then

$$\begin{aligned}
Y^t Y &= \begin{pmatrix} \mathcal{J}_1^t \mathcal{J}_1 & \mathbf{1}^t \otimes \mathcal{J}_1^t \mathcal{A}_1 \\ \mathbf{1} \otimes \mathcal{A}_1^t \mathcal{J}_1 & J \otimes \mathcal{A}_1^t \mathcal{A}_1 \end{pmatrix} \\
&= \begin{pmatrix} qa^2 J & ab \mathbf{1}^t \otimes J \\ ab \mathbf{1} \otimes J & J \otimes [(a^2 - b^2)J + qb^2 I] \end{pmatrix}.
\end{aligned}$$

Hence,  $X_1$  admits an unfaithfully balanced split. ■

**6.4. Corollary.** For every prime power  $q \equiv -1 \pmod{4}$ , and for every integer  $m > 0$ , there is an  $\text{OD}(q^m(q+1); q^m, q^{m+1})$

**Proof.** It is well known (see Ionin and Shrikhande, 2006) that there is a skew-symmetric  $W(q+1, q)$ . Apply the theorem to this matrix. ■

**6.5. Corollary.** For every prime power  $q \equiv -1 \pmod{4}$ , there is an unfaithful balancedly splittable  $\text{OD}(q(q+1); q, q^2)$ .

## 2. Balancedly Splittable Orthogonal Designs

**6.6. Example.** Using the skew-symmetric Paley weighing matrix <sup>10)</sup>  $W(4, 3)$  given by

$$(6.6.a) \quad \begin{pmatrix} 0 & + & + & + \\ - & 0 & + & - \\ - & - & 0 & + \\ - & + & - & 0 \end{pmatrix},$$

we construct the smallest case of an  $OD(12; 3, 9)$  given by the theorem

$$(6.6.b) \quad \begin{pmatrix} \mathbf{b} & \mathbf{b} & \mathbf{b} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} \\ \mathbf{b} & \mathbf{b} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} \\ \mathbf{b} & \mathbf{b} & \mathbf{b} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} \\ \bar{\mathbf{a}} & \bar{\mathbf{b}} & \mathbf{b} & \mathbf{b} & \mathbf{b} & \mathbf{b} & \mathbf{a} & \bar{\mathbf{b}} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \mathbf{b} & \mathbf{b} \\ b & \bar{a} & \bar{b} & b & b & b & \bar{b} & a & b & b & \bar{a} & \bar{b} \\ \bar{b} & b & \bar{a} & b & b & b & b & \bar{a} & \bar{b} & b & \bar{a} & \bar{b} \\ \bar{a} & \bar{b} & b & \bar{a} & \bar{b} & b & b & b & b & a & b & \bar{b} \\ b & \bar{a} & \bar{b} & b & \bar{a} & \bar{b} & b & b & b & \bar{b} & a & b \\ \bar{b} & b & \bar{a} & \bar{b} & b & \bar{a} & b & b & b & b & \bar{b} & a \\ \bar{a} & \bar{b} & b & a & b & \bar{b} & \bar{a} & \bar{b} & b & b & b & b \\ b & \bar{a} & \bar{b} & \bar{b} & a & b & b & \bar{a} & \bar{b} & b & b & b \\ \bar{b} & b & \bar{a} & b & \bar{b} & a & \bar{b} & b & \bar{a} & b & b & b \end{pmatrix},$$

where the unfaithful split is shown in bold.

Our first construction yields real ODs and is applicable in the case that we have a prime power  $q \equiv -1 \pmod{4}$ . Of course, since  $(q-1)/2$  is odd, it is known (see Ionin and Shrikhande, 2006) that there is a weighing matrix that is skew-symmetric, a property essential to the construction. If  $q \equiv 1 \pmod{4}$ , then  $(q-1)/2$  is even and the ensuing weighing matrix is symmetric. In this event, we need to appeal to complex ODs in order to apply the construction.

To apply the complex units, we make the following recursive definitions where  $W = \begin{pmatrix} 0 & \mathbf{1}^t \\ \mathbf{1} & Q \end{pmatrix}$  is a  $W(q+1, q)$  with  $Q^t = Q$ .

$$C_m = \begin{cases} aJ_1 & \text{if } m = 0, \text{ and} \\ J_q \otimes \mathcal{D}_{m-1} & \text{if } m > 0; \end{cases}$$

## 2. Balancedly Splittable Orthogonal Designs

$$\mathcal{D}_m = \begin{cases} bJ_1 & \text{if } m = 0, \text{ and} \\ I_q \otimes \mathcal{C}_{m-1} + iQ \otimes \mathcal{D}_{m-1} & \text{if } m > 0; \end{cases}$$

where again  $a$  and  $b$  are real commuting indeterminants. As above, we have the following lemma that is shown in precisely the same way as before, save one replaces transposition with conjugate transposition.

### 6.7. Lemma.

$$(6.7.a) \quad \mathcal{C}_m \mathcal{D}_m^* = \mathcal{D}_m \mathcal{C}_m^*;$$

$$(6.7.b) \quad \mathcal{C}_m \mathcal{C}_m^* + q \mathcal{D}_m \mathcal{D}_m^* = (q^m a^2 + q^{m+1} b^2) I; \text{ and}$$

$$(6.7.c) \quad \mathcal{C}_1^* \mathcal{C}_1 = qa^2 J, \mathcal{D}_1^* \mathcal{D}_1 = a^2 I + b^2 (qI - J), \text{ and } \mathcal{D}_1^* \mathcal{C}_1 = \mathcal{C}_1^* \mathcal{D}_1 = abJ.$$

**6.8. Theorem.** Let  $W$ ,  $\mathcal{C}_m$ , and  $\mathcal{D}_m$  be as above, and define  $Y_m = iI \otimes \mathcal{C}_m + W \otimes \mathcal{D}_m$ . Then:

$$(6.8.a) \quad \text{The matrix } Y_m \text{ is a COD}(q^m(q+1); q^m, q^{m+1}), \text{ and}$$

$$(6.8.b) \quad Y_1 \text{ admits an unfaithfully balanced split.}$$

**6.9. Corollary.** For every prime power  $q \equiv 1 \pmod{4}$ , and for every integer  $m > 0$ , there is a COD( $q^m(q+1); q^m, q^{m+1}$ ).

**6.10. Corollary.** For every prime power  $q \equiv 1 \pmod{4}$ , there is an unfaithful balancedly splittable COD( $q(q+1); q, q^2$ ).

We again explore the smallest case.

**6.11. Example.** Take  $q = 5$  and consider the symmetric Paley weighing matrix  $W(6, 5)$  given by

$$(6.11.a) \quad \begin{pmatrix} 0 & + & + & + & + & + \\ + & 0 & + & - & - & + \\ + & + & 0 & + & - & - \\ + & - & + & 0 & + & - \\ + & - & - & + & 0 & + \\ + & + & - & - & + & 0 \end{pmatrix}.$$



## 2. Balancedly Splittable Orthogonal Designs

Applying the construction, we obtain

[illegible]

a COD(30; 5, 25), where the unfaithful split is shown in bold.

\*\*\*

**6.3. Faithful Construction.** Here we will introduce a most useful construction of orthogonal designs admitting a faithful split. We will see later how we can use these constructed matrices in constructing other objects.

To begin, we assume the existence of a full  $\text{OD}(n; s_1, \dots, s_u)$ , say  $X$ , and label the rows of  $X$  as  $x_0, \dots, x_{n-1}$ . Further, assume that the coefficients of the indeterminants of the first row and column are  $+1$ . We need to extend the idea of an auxiliary matrix given in Example 5.6. To do this, we will follow Kharaghani and Suda (2018) in defining the auxiliary matrix of an OD thus: Let  $H$  be the Hadamard matrix obtained by setting each indeterminate of  $X$  to  $+1$ , and label the rows of  $H$  as  $h_0, \dots, h_{n-1}$ . Then the auxiliary matrices <sup>11)</sup> of  $X$  are given by  $c_i = h_i^t x_i$ . We have the following result.

## 2. Balancedly Splittable Orthogonal Designs

**6.12. Lemma.** Let  $c_i = h_i^t x_i$ , for  $i \in \{0, \dots, n-1\}$ , be the auxiliary matrices of an  $\text{OD}(n; s_1, \dots, s_u)$   $X$  where  $XX^t = \sigma I$ . Then:

$$(6.12.a) \quad \sum_i c_i c_i^t = n\sigma I_n, \text{ and}$$

$$(6.12.b) \quad c_i c_j^t = \delta_{ij} \sigma h_i^t h_i.$$

We need the simple fact that if  $(a, b)$  denotes the concatenation of sequences  $a$  and  $b$ , then  $(a, b)$  and  $(a, -b)$  is a Golay pair (see §3). Continuing to let  $c_0, \dots, c_{n-1}$  be the auxiliary matrices of the  $\text{OD}(n; s_1, \dots, s_u)$   $X$ , then  $a = (c_0, c_1, \dots, c_{n-1}, c_{n-1}, \dots, c_1)$  and  $b = (c_0, c_1, \dots, c_{n-1}, -c_{n-1}, \dots, -c_1)$  form a complementary pair. Let  $A$  and  $B$  be the block-circulant matrices with first rows  $a$  and  $b$ , respectively.

Now, take  $\tilde{X} = \begin{pmatrix} + & + \\ + & - \end{pmatrix} \otimes X$  and  $\tilde{H} = \begin{pmatrix} + & + \\ + & - \end{pmatrix} \otimes H$ , and label the block rows  $\tilde{x}_0, \dots, \tilde{x}_{2n-1}$  and  $\tilde{h}_0, \dots, \tilde{h}_{2n-1}$ . Define  $G = \tilde{h}_0^t \tilde{x}_0$ , and define the block matrices  $E^t = (E_1^t \dots E_{2n-1}^t)^t$  and  $F = (F_1 \dots F_{2n-1})$  by  $E_i = h_0^t \tilde{x}_i$  and  $F_i = \tilde{h}_i^t x_0$ .

As before, we then take  $Z = \begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$ . The next result then follows.

**6.13. Theorem.** If there is an  $\text{OD}(n; s_1, \dots, s_u)$ , then the block matrix  $Z$  is an  $\text{OD}(4n^2; 4ns_1, \dots, 4ns_u)$ .

**Proof.** The proof amounts to checking the block entries of  $ZZ^t$ .

To begin,  $GE_i^t = (\tilde{h}_0^t \tilde{x}_0)(h_0^t \tilde{x}_i)^t = \tilde{h}_0^t (\tilde{x}_0 \tilde{x}_i^t) h_0 = O$ , hence  $GE^t = EG^t = O$ . Then  $F_i c_j^t = (\tilde{h}_i^t x_0)(h_j^t x_j)^t = \tilde{h}_i^t (x_0 x_j^t) h_j = \delta_{0j} \sigma \tilde{h}_i^t h_0$  so that

$$FA^t = FB^t = \sigma \begin{pmatrix} \tilde{h}_1 \\ \vdots \\ \tilde{h}_{2n-1} \end{pmatrix}^t (\mathbf{1}_{2n-1} \otimes h_0).$$

We have, therefore, that  $FA^t - FB^t = O$ . Then the inner product between the first and second, and the first and third, block rows of  $Z$  vanish.

## 2. Balancedly Splittable Orthogonal Designs

Next,  $E_i E_j^t = (h_0^t \tilde{x}_i)(h_0^t \tilde{x}_j)^t = h_0^t(\tilde{x}_i \tilde{x}_j^t)h_0 = \delta_{ij} 2\sigma J_n$ . It follows that  $EE^t = (E_i E_j^t) = (\delta_{ij} 2\sigma J_n) = 2\sigma(I_{2n-1} \otimes J_n)$ .

We need to examine the product  $AB^t$  and in order to do that, we need to examine the cross-product correlations (see §3). To begin, the product between the first block row and column of  $A$  and  $B^t$  is given by  $c_0 c_0^t + \sum_{i=1}^{n-1} c_i c_i^t - \sum_{i=1}^{n-1} c_i c_i^t = \sigma J_n$ . Next, let  $a$  and  $b$  be two sequences of length  $2n - 1$  defined by

$$a_i = \begin{cases} c_i & \text{if } 0 \leq i < n, \text{ and} \\ c_{2n-i-1} & \text{if } n \leq i < 2n - 1, \end{cases}$$

$$b_i = \begin{cases} c_0 & \text{if } i = 0, \\ -c_i & \text{if } 0 < i < n, \text{ and} \\ c_{2n-i-1} & \text{if } n \leq i < 2n - 1. \end{cases}$$

For  $j \in \{1, \dots, 2n-2\}$ , we have  $C_j(a, b) = \sum_{i=0}^{2n-2} a_i b_{i+j}^t = \sum_{i=0}^{n-1} a_i b_{i+j}^t + \sum_{i=n}^{2n-2} a_i b_{i+j}^t$ , where precisely one of the right-hand sums is nonzero by (6.12.b). For  $j \in \{1, \dots, n-1\}$ , we have that

$$\begin{aligned} \sum_{i=0}^{n-1} a_i b_{i+j}^t &= a_0 b_j^t + \sum_{i=1}^{n-j-1} a_i b_{i+j}^t + \sum_{i=n-j}^{n-1} a_i b_{i+j}^t \\ &= c_0 c_j^t - \sum_{i=1}^{n-j-1} c_i c_{i+j}^t + \sum_{i=n-j}^{n-1} c_i c_{2n-j-i-1}^t \\ &= \sum_{i=0}^{j-1} c_{n-j+i} c_{n-i-1}^t, \end{aligned}$$

and

$$\sum_{i=n}^{2n-2} a_i b_{i+j}^t = \sum_{i=n}^{2n-j-2} a_i b_{i+j}^t + \sum_{i=2n-j-1}^{2n-2} a_i b_{i+j}^t$$

## 2. Balancedly Splittable Orthogonal Designs

$$\begin{aligned}
&= \sum_{i=0}^{n-j-2} c_{n-i-1} c_{n-j-i-1}^t + c_j c_0^t - \sum_{i=1}^{j-1} c_{j-i} c_i^t \\
&= - \sum_{i=1}^{j-1} c_{j-i} c_i^t.
\end{aligned}$$

We have shown that, for  $j \in \{1, \dots, n-1\}$ ,

$$\begin{aligned}
C_j(a, b) &= \sum_{i=0}^{j-1} c_{n-j+i} c_{n-i-1}^t - \sum_{i=1}^{j-1} c_{j-i} c_i^t \\
&= \begin{cases} \sigma h_{n-i-1}^t h_{n-i-1} & \text{if } j-1 = 2i, \text{ for some } i; \text{ and} \\ -\sigma h_i^t h_i & \text{if } j = 2i, \text{ for some } i \end{cases}
\end{aligned}$$

Similarly, we find that  $C_{2n-j-1}(a, b) = -C_j(a, b)$ , for  $j \in \{1, \dots, n-1\}$ , and  $C_j(b, a) = -C_j(a, b)$ , for all  $j \in \{1, \dots, 2n-2\}$ .

Putting things together, we have shown that

$$\begin{aligned}
AB^t &= \text{circ}(\sigma J, C_{2n-2}(a, b), \dots, C_1(a, b)) \\
&= \text{circ}(\sigma J, -C_1(a, b), \dots, -C_{n-1}(a, b), C_{n-1}(a, b), \dots, C_1(a, b)) \\
&= \sigma \text{circ}(J, -h_{n-1}^t h_{n-1}, \dots, -h_{n/2}^t h_{n/2}, \\
&\quad h_{n/2}^t h_{n/2}, \dots, h_{n-1}^t h_{n-1}), \text{ and} \\
BA^t &= \sigma \text{circ}(J, h_{n-1}^t h_{n-1}, \dots, h_{n/2}^t h_{n/2}, \\
&\quad -h_{n/2}^t h_{n/2}, \dots, -h_{n-1}^t h_{n-1}).
\end{aligned}$$

It follows that the product between the second and third block rows vanishes, i.e.

$$-EE^t + AB^t + BA^t = O.$$

It remains to evaluate the block diagonal entries of  $ZZ^t$ . The first block row

## 2. Balancedly Splittable Orthogonal Designs

gives

$$\begin{aligned}
 GG^t + 2 \sum_{i=1}^{2n-1} F_i F_i^t &= 2\sigma J + 2\sigma \sum_{i=1}^{2n-1} \tilde{h}_i^t \tilde{h}_i \\
 &= 2\sigma J + 2\sigma \left( \sum_{i=0}^{2n-1} \tilde{h}_i^t \tilde{h}_i - \tilde{h}_0^t \tilde{h}_0 \right) \\
 &= 2\sigma J + 2\sigma (2nI - J) \\
 &= 4n\sigma I.
 \end{aligned}$$

By a similar argument about the sequences  $a$  and  $b$  given above, we find that

$$\begin{aligned}
 AA^t &= \sigma \text{circ}(2nI - J, h_{n-1}^t h_{n-1}, \dots, h_{n/2}^t h_{n/2}, \\
 &\quad h_{n/2}^t h_{n/2}, \dots, h_{n-1}^t h_{n-1}), \text{ and} \\
 BB^t &= \sigma \text{circ}(2nI - J, -h_{n-1}^t h_{n-1}, \dots, -h_{n/2}^t h_{n/2}, \\
 &\quad -h_{n/2}^t h_{n/2}, \dots, -h_{n-1}^t h_{n-1}).
 \end{aligned}$$

Thus,  $EE^t + AA^t + BB^t = 4\sigma I$ . The proof is complete.  $\blacksquare$

We have shown that  $Z$  is an OD. It remains to show that  $Z$  admits a faithful split. This is accomplished with the next result.

**6.14. Theorem.** Assume the OD  $Z$  of the previous theorem. Then:

(6.14.a) The submatrices  $\begin{pmatrix} F \\ A \\ B \end{pmatrix}$  and  $\begin{pmatrix} -F \\ B \\ A \end{pmatrix}$  yield stable splits, and

(6.14.b) The submatrices  $\begin{pmatrix} E & A & B \end{pmatrix}$  and  $\begin{pmatrix} -E & B & A \end{pmatrix}$  yield unstable splits.

**Proof.** It suffices to show the result for one matrix of each class. Note that

$$\begin{pmatrix} F \\ A \\ B \end{pmatrix} (F^t \ A^t \ B^t) = \begin{pmatrix} FF^t & FA^t & FB^t \\ AF^t & AA^t & AB^t \\ BF^t & BA^t & BB^t \end{pmatrix}.$$

## 2. Balancedly Splittable Orthogonal Designs

It follows by the proof of the previous theorem that

$$\begin{aligned}
FF^t &= 2n\sigma I - \sigma J, \\
FA^t &= FB^t = \sigma \sum_{i=1}^{2n-1} \tilde{h}_i^t h_0, \\
AA^t &= \sigma \text{circ}(2nI - J, h_{n-1}^t h_{n-1}, \dots, h_{n/2}^t h_{n/2}, \\
&\quad h_{n/2}^t h_{n/2}, \dots, h_{n-1}^t h_{n-1}), \text{ and} \\
BB^t &= \sigma \text{circ}(2nI - J, -h_{n-1}^t h_{n-1}, \dots, -h_{n/2}^t h_{n/2}, \\
&\quad -h_{n/2}^t h_{n/2}, \dots, -h_{n-1}^t h_{n-1}).
\end{aligned}$$

Therefore, the product between distinct rows is  $\pm\sigma$ , which shows (6.14.a).

Next,

$$\begin{pmatrix} E^t \\ A^t \\ B^t \end{pmatrix} (E \ A \ B) = \begin{pmatrix} E^t E & E^t A & E^t B \\ A^t E & A^t A & A^t B \\ B^t E & B^t A & B^t B \end{pmatrix}.$$

However,  $E^t E = n \sum_{i=1}^{2n-1} \tilde{x}_i^t \tilde{x}_0$  has off-diagonal entries in the set

$$\{\pm x_1^{m_1} \dots x_u^{m_u} x_1^{*m'_1} \dots x_u^{*m'_u} : m_i, m'_i \in \mathbf{N}\},$$

which shows (6.14.b).  $\blacksquare$

**6.15. Example.** Applying the construction to the OD(2; 1, 1)

$$(6.15.a) \quad \begin{pmatrix} a & b \\ b & \bar{a} \end{pmatrix},$$

we obtain the balancedly splittble OD(16; 8, 8)

## 2. Balancedly Splittable Orthogonal Designs

$$(6.15.b) \quad \begin{pmatrix} a & b & a & b & \mathbf{a} & \mathbf{b} & \mathbf{a} & \mathbf{b} & \mathbf{a} & \mathbf{b} & \bar{a} & \bar{b} & \bar{a} & \bar{b} & \bar{a} & \bar{b} \\ a & b & a & b & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & a & b & \bar{a} & \bar{b} & a & b \\ a & b & a & b & \mathbf{a} & \mathbf{b} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \bar{a} & \bar{b} & a & b & a & b \\ a & b & a & b & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & a & b & a & b & \bar{a} & \bar{b} \\ b & \bar{a} & b & \bar{a} & \mathbf{a} & \mathbf{b} & \mathbf{b} & \bar{\mathbf{a}} & \mathbf{b} & \bar{\mathbf{a}} & a & b & b & \bar{a} & \bar{b} & a \\ b & \bar{a} & b & \bar{a} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \bar{\mathbf{b}} & \mathbf{a} & a & b & \bar{b} & a & b & \bar{a} \\ a & b & \bar{a} & \bar{b} & \mathbf{b} & \bar{\mathbf{a}} & \mathbf{a} & \mathbf{b} & \mathbf{b} & \bar{\mathbf{a}} & \bar{b} & a & a & b & b & \bar{a} \\ a & b & \bar{a} & \bar{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & b & \bar{a} & a & b & \bar{b} & a \\ b & \bar{a} & \bar{b} & a & \mathbf{b} & \bar{\mathbf{a}} & \mathbf{b} & \bar{\mathbf{a}} & \mathbf{a} & \mathbf{b} & b & \bar{a} & \bar{b} & a & a & b \\ b & \bar{a} & \bar{b} & a & \bar{\mathbf{b}} & \mathbf{a} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{a} & \mathbf{b} & \bar{b} & a & b & \bar{a} & a & b \\ \bar{b} & a & \bar{b} & a & \mathbf{a} & \mathbf{b} & \mathbf{b} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \mathbf{a} & a & b & b & \bar{a} & b & \bar{a} \\ \bar{b} & a & \bar{b} & a & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{a}} & a & b & \bar{b} & a & \bar{b} & a \\ \bar{a} & \bar{b} & a & b & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{a} & \mathbf{b} & \mathbf{b} & \bar{\mathbf{a}} & \bar{b} & \bar{a} & a & b & b & \bar{a} \\ \bar{a} & \bar{b} & a & b & \mathbf{b} & \bar{\mathbf{a}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{b}} & \mathbf{a} & \bar{b} & a & a & b & \bar{b} & a \\ \bar{b} & a & b & \bar{a} & \mathbf{b} & \bar{\mathbf{a}} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{a} & \mathbf{b} & b & \bar{a} & b & \bar{a} & a & b \\ \bar{b} & a & b & \bar{a} & \bar{\mathbf{b}} & \mathbf{a} & \mathbf{b} & \bar{\mathbf{a}} & \mathbf{a} & \mathbf{b} & \bar{b} & a & \bar{b} & a & a & b \end{pmatrix},$$

where a stable vertical split is shown in bold.

Finally, we see at once how Theorem 5.9 is a consequence of Theorem 6.13 as we can simply set the indeterminants to +1 to obtain the result.

**6.16. Example.** Beginning with the OD(4; 1, 1, 1, 1)

$$(6.16.a) \quad \begin{pmatrix} a & b & c & d \\ \bar{b} & a & \bar{d} & c \\ \bar{c} & d & a & \bar{b} \\ \bar{d} & \bar{c} & b & a \end{pmatrix},$$

we obtain the OD(64; 16, 16, 16, 16) constructed in A3.

**6.17. Example.** As before, we are not limited to real matrices. Consider the COD(6; 1, 5)

$$(6.17.a) \quad \begin{pmatrix} ia & b & b & b & b & b \\ b & ia & b & \bar{b} & \bar{b} & b \\ b & b & ia & b & \bar{b} & \bar{b} \\ b & \bar{b} & b & ia & b & \bar{b} \\ b & \bar{b} & \bar{b} & b & ia & b \\ b & b & \bar{b} & \bar{b} & b & ia \end{pmatrix},$$

where  $a$  and  $b$  are real, commuting indeterminants. Using the method, one can construct a COD(144; 24, 120), which we do not show in this essay due to its size.

## 2. Balancedly Splittable Orthogonal Designs

### §7. Related Configurations

In this section, the balancedly splittable orthogonal designs of the previous section are applied in the construction of related objects. We will explore quasi-symmetric balanced incomplete block designs, equiangular tight frames, and unbiased orthogonal designs.

\* \* \*

**7.1. Quasi-Symmetric BIBDs.** Symmetric designs are characterized by the fact that the blocks of the design intersect in a constant number of points. The “next best” designs are those which have two cardinalities for the intersections between distinct blocks.

**7.1. Definition.** A balanced incomplete block design is *quasi-symmetric* if there exist two cardinalities that exist for the intersections between pairs of distinct blocks. The cardinalities of the intersections are called the *intersection numbers* of the design <sup>12</sup>).

These beautiful objects are studied extensively in Shrikhande and Sane (1991). Ionin and Shrikhande (2006) also contains a study of these objects, especially as it pertains to SRGs and association schemes.

**7.2. Example.** The following is the incidence matrix of a quasi-symmetric BIBD(6, 15, 5, 2, 1)

$$(7.2.a) \quad \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

with block intersection numbers 0 and 2.



## 2. Balancedly Splittable Orthogonal Designs

Using the balancedly splittable Hadamard matrices constructed in Theorem 6.13, we can construct quasi-symmetric BIBDs. For the proposition, we assume the notations of §5.3.

**7.3. Proposition.** If there is a Hadamard matrix of order  $n$ , there is a quasi-symmetric BIBD $(2n^2 - n, 4n^2 - 1, 2n^2 - n - 1, n^2 - n, n^2 - n - 1)$  with intersection numbers  $(n^2 - n)/2$  and  $(n^2 - 2n)/2$ .

**Proof.** If there exists a normalized Hadamard matrix of order  $n$ , there exists a balancedly splittable Hadamard matrix of order  $4n^2$  by Theorem 5.9 (or by Theorem 6.13 upon setting the indeterminants to unity if there is a full OD of order  $n$ ). It suffices to show the result for one of the four splits obtained by the theorem; in particular, we will show the result for the submatrix  $X = \begin{pmatrix} -F \\ B \\ A \end{pmatrix}$ . Form the matrix  $\tilde{Y} = (1/2)(J - X)$ , and take  $Y$  to be the matrix formed by omitting the first row of  $\tilde{Y}$  consisting of all zeros.

Since  $H$  is assumed to be normalized, each row of  $H$  and  $\begin{pmatrix} + & + \\ + & - \end{pmatrix} \otimes H$ , other than the first, of course, contain  $n/2$  and  $n^2/2 - 1$ s, respectively. By construction, then, the first row of  $F$  consists of all ones, while the remaining rows have  $n^2 - n + 1$ s; thus, the same rows of  $-F$  have  $n^2 - n - 1$ s. Similarly,  $A$  and  $B$  have  $n(2n - 2)/2 = n^2 - n - 1$ s in each row. It also follows that there must be  $2n^2 - 2n + n - 1 = 2n^2 - n - 1 - 1$ s in each column of  $X$ .

Finally, the index of the design follows by considering the Menon design<sup>13</sup>) induced by the regular Hadamard matrix of the construction.

It remains to consider the block intersection numbers of the design. Consider the first row of  $X$  along with any other two distinct rows. Without loss of general-

## 2. Balancedly Splittable Orthogonal Designs

ity, we can assume the following situation.

$$\begin{array}{cccc}
 \overbrace{\begin{array}{ccc} - & \dots & - \\ + & \dots & + \\ + & \dots & + \end{array}}^a & 
 \overbrace{\begin{array}{ccc} - & \dots & - \\ + & \dots & + \\ - & \dots & - \end{array}}^b & 
 \overbrace{\begin{array}{ccc} - & \dots & - \\ - & \dots & - \\ + & \dots & + \end{array}}^c & 
 \overbrace{\begin{array}{ccc} - & \dots & - \\ - & \dots & - \\ - & \dots & - \end{array}}^d
 \end{array}$$

This yields the two linear systems of equations

$$\begin{aligned}
 a + b + c + d &= 2n^2 - n, \\
 -a - b + c + d &= -n, \\
 -a + b - c + d &= -n, \text{ and} \\
 a - b - c + d &= \pm n.
 \end{aligned}$$

Solving for  $d$  gives  $2d = n^2 - n$  or  $n^2 - 2n$ , which proves the result.  $\blacksquare$

**7.4. Example.** The quasi-symmetric BIBD(6, 15, 5, 2, 1) given by (7.2.a) is obtained by the splittable Hadamard matrix (5.6.e).

The quasi-symmetric designs of the previous result are not new, however. They represent a subset of the designs obtained in Bracken et al. (2006) viz. McGuire (1997).

\* \* \*

**7.2. Equiangular Tight Frames.** In our earliest collegiate education, we learn that given some linear space, the “nice” bases are those which are orthonormal. As Han et al. (2007) points out, however, the property of orthonormality can be too restrictive in many applications. For instance, if a signal is interfered with in transmission, then the lost information cannot be recovered.

By contrast, a frame is an overdetermined <sup>14)</sup> spanning set of vectors. In this way, calculations and applications involving these objects can at times be simpli-

## 2. Balancedly Splittable Orthogonal Designs

fied and data loss deterred. Frames have theoretical applications as well, as shown in the seminal paper of Paley and Wiener (1987) using different language, however.

We will not have the opportunity to explore the theory of frames in any depth; rather, we direct the reader to the standard references of Christensen (2016), Waldron (2018), and Young (2001).

**7.5. Definition.** Let  $H$  be a Hilbert space <sup>15)</sup> with inner product  $\langle \cdot, \cdot \rangle$ , and let  $\{f_i\} \subset H$ . Then  $\{f_i\}$  is a *frame* if there exists two constants  $A$  and  $B$  such that

$$(7.5.a) \quad A\|f\|^2 \leq \sum_i |\langle f, f_i \rangle|^2 \leq B\|f\|^2, \text{ for all } f \in H.$$

$A$  and  $B$  are called frame bounds. If  $\#\{f_i\} < \infty$ , it is a *finite frame*. If  $A = B$ , the frame is said to be *tight*. If  $A = B = 1$ , then  $\{f_i\}$  is said to be a *Parseval frame*.

The frame bounds of a frame are not unique. The *optimal upper frame bound* is the infimum of the collection of all upper bounds of the frame. Similarly, the *optimal lower frame bound* is the supremum of the collection of all lower bounds of the frame.

It also follows from the definition that  $\overline{\text{span}}\{f_i\} = H$ .

**7.6. Example.** If  $\{e_i\}$  is an orthonormal basis, then  $\{e_0, e_0, e_1, e_1, \dots\}$ , where each element is repeated twice, is a tight frame with  $A = 2$ .

**7.7. Example.** If  $\{e_i\}$  is an orthonormal basis, then  $\{e_0, 2^{-\frac{1}{2}}e_1, 2^{-\frac{1}{2}}e_1, 3^{-\frac{1}{2}}e_2, 3^{-\frac{1}{2}}e_2, 3^{-\frac{1}{2}}e_2\}$ , where the element  $(n+1)^{-\frac{1}{2}}e_n$  is repeated  $n+1$  times, is a frame with  $A = 1$ .

From this point on, we will assume that we are working with a finite dimensional Hilbert space of dimension  $k$ . Our frames will, therefore, always be finite, say of order  $n$ . Just as for the standard bases of a space, we can relate matrices to finite frames.

## 2. Balancedly Splittable Orthogonal Designs

**7.8. Definition.** Let  $\{f_0, \dots, f_{n-1}\}$  be a finite frame in a finite dimensional Hilbert space  $H$ . The matrix  $\Theta = \begin{pmatrix} f_0^* \\ \vdots \\ f_{n-1}^* \end{pmatrix}$  is the *analysis operator* of the frame, and  $\Theta^*$  is called the *synthesis operator* of the frame.

The following is then a restatement of the definitions.

**7.9. Proposition.** Let  $x_0, \dots, x_{n-1}$  be vectos in a Hilbert space  $H$  of dimension  $n$ , and define the  $k \times n$  matrix  $T = (x_0 \dots x_{n-1})$ . Then:

(7.9.a)  $\{x_0, \dots, x_{n-1}\}$  is a frame if and only if  $\text{rank}(T) = n$ ,

(7.9.b)  $\{x_0, \dots, x_{n-1}\}$  is a tight frame with bound  $A$  if and only if  $TT^* = \sqrt{A}I$ .

In what follows, we will take  $\mathbf{H} = \{a_1 + a_i i + a_j j + a_k k : a_1, a_i, a_j, a_k \in \mathbf{R}\}$ , i.e. the  $\mathbf{R}$ -algebra of quaternions<sup>16</sup>), and we will consider frames over  $\mathbf{H}^k$ . Furthermore, for  $q = z + wj$  where  $z, w \in \mathbf{C}$ , we define  $\text{Co}_1(q) = z$  and  $\text{Co}_2(q) = w^*$ . Finally, define  $[\cdot]_{\mathbf{C}} : \mathbf{H}^d \rightarrow \mathbf{C}^{2d}$  by  $[q]_{\mathbf{C}} = \begin{pmatrix} \text{Co}_1(q) \\ \text{Co}_2(q) \end{pmatrix} = \begin{pmatrix} z \\ w^* \end{pmatrix}$ .

The following is Theorem 3.2 of Waldron (2020).

**7.10. Theorem.** (7.10.a) Tight frames for  $\mathbf{C}^{2k}$  correspond to tight frames for  $\mathbf{H}^k$ , and

(7.10.b) Tight frames for  $\mathbf{H}^k$  correspond to tight frames for  $\mathbf{C}^k$  if and only if

$$\sum_{i,j} |\text{Co}_1(\langle f_i, f_j \rangle)|^2 = \sum_{i,j} |\text{Co}_2(\langle f_i, f_j \rangle)|^2.$$

In light of this result, it is of interest to find frames over  $\mathbf{H}^k$  for which they are not equivalent to frames over  $\mathbf{C}^{2d}$ , i.e. there can be no reduction in the ring of scalars. We present two examples of such frames.

**7.11. Example.** The following is a QOD(2; 1, 1)

$$(7.11.a) \begin{pmatrix} \bar{a} & bi \\ \bar{b}j & ak \end{pmatrix},$$

where  $a$  and  $b$  are real variables. We then arrive at the following QOD(16; 8, 8)

## 2. Balancedly Splittable Orthogonal Designs

$$(7.11.b) \quad \begin{pmatrix} \bar{a} & bi & \bar{a} & bi & \bar{a} & bi & \bar{a} & bi & \bar{a} & bi & a & \bar{bi} & a & \bar{bi} & a & \bar{bi} \\ \bar{a} & bi & \bar{a} & bi & a & \bar{bi} & \bar{a} & bi & a & \bar{bi} & \bar{a} & bi & a & \bar{bi} & \bar{a} & bi \\ \bar{a} & bi & \bar{a} & bi & \bar{a} & bi & a & \bar{bi} & a & \bar{bi} & \bar{a} & bi & \bar{a} & bi & \bar{a} & bi \\ \bar{a} & bi & \bar{a} & bi & a & \bar{bi} & a & \bar{bi} & \bar{a} & bi & \bar{a} & bi & \bar{a} & bi & a & \bar{bi} \\ \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi & \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi & \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{bj} & \mathbf{\bar{a}k} \\ \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{bj} & \mathbf{\bar{a}k} & \bar{a} & bi & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{\bar{b}j} & \mathbf{ak} \\ \bar{a} & bi & a & \bar{bi} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi & \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{bj} & \mathbf{\bar{a}k} & \bar{a} & bi & \mathbf{\bar{b}j} & \mathbf{ak} \\ \bar{a} & bi & a & \bar{bi} & \mathbf{bj} & \mathbf{\bar{a}k} & \bar{a} & bi & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi & \mathbf{bj} & \mathbf{\bar{a}k} \\ \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi & \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{bj} & \mathbf{\bar{a}k} & \bar{a} & bi \\ \mathbf{\bar{b}j} & \mathbf{ak} & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{bj} & \mathbf{\bar{a}k} & \bar{a} & bi & \mathbf{bj} & \mathbf{\bar{a}k} & \mathbf{\bar{b}j} & \mathbf{ak} & \bar{a} & bi \\ bj & \bar{ak} & bj & \bar{ak} & \bar{a} & bi & \bar{bj} & ak & bj & \bar{ak} & \bar{a} & bi & \bar{bj} & ak & \bar{bj} & ak \\ bj & \bar{ak} & bj & \bar{ak} & \bar{a} & bi & bj & \bar{ak} & \bar{bj} & ak & \bar{a} & bi & bj & \bar{ak} & bj & \bar{ak} \\ a & \bar{bi} & \bar{a} & bi & bj & \bar{ak} & \bar{a} & bi & \bar{bj} & ak & \bar{bj} & ak & \bar{a} & bi & \bar{bj} & ak \\ a & \bar{bi} & \bar{a} & bi & \bar{bj} & ak & \bar{a} & bi & bj & \bar{ak} & bj & \bar{ak} & \bar{a} & bi & bj & \bar{ak} \\ bj & \bar{ak} & \bar{bj} & ak & \bar{bj} & ak & bj & \bar{ak} & \bar{a} & bi & \bar{bj} & ak & \bar{bj} & ak & \bar{a} & bi \\ bj & \bar{ak} & \bar{bj} & ak & bj & \bar{ak} & \bar{bj} & ak & \bar{a} & bi & bj & \bar{ak} & bj & \bar{ak} & \bar{a} & bi \end{pmatrix}.$$

Taking the horizontal frame shown in bold as the synthesis operator of a frame over

$\mathbf{H}^6$  after setting  $a = b = 1$ , we find that  $(1/2)H^*H$  is given by

$$(7.11.c) \quad \begin{pmatrix} 3 & i & -i & \bar{j} & \bar{k} & 1 & \bar{i} & \bar{j} & \bar{k} & \bar{j} & \bar{k} & 1 & \bar{i} & \bar{j} & \bar{k} \\ \bar{i} & 3 & \bar{i} & -k & \bar{j} & i & 1 & k & \bar{j} & \bar{k} & \bar{j} & i & 1 & k & \bar{j} \\ -i & 3 & i & \bar{j} & \bar{k} & -i & j & k & \bar{j} & \bar{k} & -i & j & k \\ \bar{i} & -\bar{i} & 3 & k & \bar{j} & \bar{i} & -\bar{k} & j & k & \bar{j} & \bar{i} & -\bar{k} & j \\ j & \bar{k} & j & \bar{k} & 3 & i & 1 & i & 1 & i & 1 & \bar{i} & -\bar{i} & 1 & i \\ k & j & k & j & \bar{i} & 3 & \bar{i} & 1 & \bar{i} & 1 & i & 1 & i & -\bar{i} & 1 \\ 1 & \bar{i} & -i & 1 & i & 3 & i & 1 & i & 1 & i & 1 & \bar{i} & -\bar{i} \\ i & 1 & \bar{i} & -\bar{i} & 1 & \bar{i} & 3 & \bar{i} & 1 & \bar{i} & 1 & i & 1 & i & - \\ j & \bar{k} & \bar{j} & k & 1 & i & 1 & i & 3 & i & -\bar{i} & 1 & i & 1 & \bar{i} \\ k & j & \bar{k} & \bar{j} & \bar{i} & 1 & \bar{i} & 1 & \bar{i} & 3 & i & -\bar{i} & 1 & i & 1 \\ j & \bar{k} & j & \bar{k} & 1 & \bar{i} & 1 & i & -\bar{i} & 3 & i & -\bar{i} & -\bar{i} \\ k & j & k & j & i & 1 & \bar{i} & 1 & i & -\bar{i} & 3 & i & -i & - \\ 1 & \bar{i} & -i & -\bar{i} & 1 & \bar{i} & 1 & i & -\bar{i} & 3 & i & -\bar{i} \\ i & 1 & \bar{i} & -i & -i & 1 & \bar{i} & 1 & i & -\bar{i} & 3 & i & - \\ j & \bar{k} & \bar{j} & k & 1 & i & -\bar{i} & 1 & \bar{i} & -\bar{i} & -\bar{i} & 3 & i \\ k & j & \bar{k} & \bar{j} & \bar{i} & 1 & i & -i & 1 & i & -i & -\bar{i} & 3 \end{pmatrix}.$$

It follows that  $320 = \sum_{i,j} |\text{Co}_1(\langle f_i, f_j \rangle)|^2 \neq \sum_{i,j} |\text{Co}_2(\langle f_i, f_j \rangle)|^2 = 64$  so that the frame is not reducible to a frame over  $\mathbf{C}^{12}$ .

**7.12. Example.** Beginning with the QOD(6; 1, 5)

## 2. Balancedly Splittable Orthogonal Designs

$$(7.12.a) \quad \begin{pmatrix} a & b & b & b & b & b \\ b & \bar{a} & bk & \overline{bk} & \overline{bk} & bk \\ b & bk & \bar{a} & bk & \overline{bk} & \overline{bk} \\ b & \overline{bk} & bk & \bar{a} & bk & \overline{bk} \\ b & \overline{bk} & \overline{bk} & bk & \bar{a} & bk \\ b & bk & \overline{bk} & \overline{bk} & bk & \bar{a} \end{pmatrix},$$

we apply the construction to obtain a QOD(144; 24, 120). Taking  $H$  to again be the first horizontal frame of the splittable QOD, we find that  $29056 = \sum_{i,j} |\text{Co}_1(\langle f_i, f_j \rangle)|^2 \neq \sum_{i,j} |\text{Co}_2(\langle f_i, f_j \rangle)|^2 = 8960$ . It follows that we have constructed another quaternion frame not reducible to a complex frame.

\* \* \*

**7.3. Unbiased Orthogonal Designs.** Let  $A$  and  $B$  be two orthonormal bases of the Hilbert space  $\mathbf{C}^k$ . The bases are said to be *mutually unbiased* in the event that  $|\langle a, b \rangle| = k^{-\frac{1}{2}}$ , for every  $a \in A$  and  $b \in B$ , and where  $\langle a, b \rangle$  denotes the usual sesquilinear inner product between  $a$  and  $b$ .

The reader will recognize unbiased bases as a subtype of equiangular lines. These objects are fundamental in many applications such as quantum key distribution (see Boykin et al., 2005).

Unitary Hadamard matrices, i.e. a matrices with entries from the unit circle in the complex plane whose rows are pairwise orthogonal, have received much attention with their connection to unbiased bases. Two unitary Hadamard matrices  $H$  and  $K$  of order  $n$  are *unbiased* if  $HK^* = \sqrt{n}L$ , for some unitary Hadamard matrix  $L$ . Recently, these ideas were studied for the case of unitary weighing matrices<sup>17</sup> in Best et al. (2015). The case of unbiased real Hadamard matrices were studied in Holzmann et al. (2010), and the case of unbiased quaternary complex Hadamard in Best and Kharaghani (2010).

We have seen that extending concepts from matrices of concrete values to matrices of indeterminants usually presents us with subtle difficulties, and extending

## 2. Balancedly Splittable Orthogonal Designs

unbiasedness is no different. In Kharaghani and Suda (2018), unbiased orthogonal designs are presented thus.

**7.13. Definition.** Let  $X_1$  and  $X_2$  be two instances of an  $\text{OD}(n; s_1, \dots, s_u)$  in the indeterminants  $x_1, \dots, x_u$ . Then  $X_1$  and  $X_1$  are *unbiased* with parameter  $\alpha \in \mathbf{R}_+$  if there is a  $(-1, 0, 1)$ -matrix  $W$  such that

$$(7.13.a) \quad X_1 X_2^t = \left( \alpha^{-\frac{1}{2}} \sum_i s_i x_i^t \right) W.$$

We use the stable construction presented in the previous section to construct pairs of unbiased orthogonal designs. Note that this is essentially a generalization of a method presented in Kharaghani and Suda (2019). In what follows, we take  $Z$  to be as in §9.3.

**7.14. Proposition.** Let

$$V = \begin{pmatrix} -F \\ B \\ A \end{pmatrix}, \text{ and } U = \begin{pmatrix} G & F \\ E & A \\ -E & B \end{pmatrix}$$

so that  $Z = \begin{pmatrix} U & V \end{pmatrix}$ , and take  $Y = \begin{pmatrix} U & -V \end{pmatrix}$ . Then  $Y$  is an OD of the same order and type as  $Z$ . In particular,  $ZY^t = 2\sigma K$ , where  $K$  is a Hadamard matrix.

**Proof.** That  $Y$  is also an OD is clear. Then  $ZY^t = (UU^t - VV^t)$ . We claim that  $K = (1/2\sigma)(UU^t - VV^t)$  is a Hadamard matrix. Indeed, since the vertical frames of  $Z$  constitute a stable split, and since  $Z$  is an OD, we find that  $K$  has entries from the set  $\{-1, 1\}$ . Furthermore,

$$\begin{aligned} KK^t &= \frac{1}{4\sigma^2} (UU^t UU^t + VV^t VV^t) \\ &= \frac{n}{\sigma} (UU^t + VV^t) \\ &= 4n^2 I, \end{aligned}$$

## 2. Balancedly Splittable Orthogonal Designs

and the proof is complete. ■

## Notes

9. Recall the angle between two vectors in  $\mathbf{R}^\ell$  is given by  $\alpha = \arccos(\langle u, v \rangle / \|u\| \|v\|)$ . It follows that we only need to consider  $|\langle u, v \rangle|$  whenever we want to deduce the number of angles that exist between a collection of vectors in  $\mathbf{R}^\ell$ .
10. The classic construction due to Paley goes as follows. Let  $q$  be odd, and take  $\text{GF}(q) = \{a_0 = 0, a_1, \dots, a_{q-1}\}$ . If  $\eta$  is the quadratic character on  $\text{GF}(q)$ , define  $Q$  by  $Q_{ij} = \eta(a_j - a_i)$ . Finally, form the matrix  $W = \begin{pmatrix} 0 & \mathbf{1}^t \\ (-1)^{(q-1)/2} \mathbf{1} & Q \end{pmatrix}$ . Then  $W$  is a  $W(q+1, q)$ . If  $q \equiv -1 \pmod{4}$ , then  $W^t = -W$ ; while if  $q \equiv 1 \pmod{4}$ , then  $W^t = W$ . See Hall (1986) for more detail.
11. In defining the auxiliary matrix of a full OD, we limit ourselves to using the Hadamard matrix obtained from the OD by setting the indeterminants equal to  $+1$ . This is done that the formation of the auxiliary matrices is unique.
12. Quasi-symmetric designs are often studied in the more general context of  $t$ -designs (see §1.1). This is the approach taken in Ionin and Shrikhande (2006) and Shrikhande and Sane (1991).
13. It isn't difficult to see that a regular Hadamard matrix  $H$  must have square order  $4u^2$ . Moreover, it follows that  $(1/2)(J - H)$  is a symmetric BIBD( $4u^2, 2h^2 - h, h^2 - h$ ). The converse can also be seen to be true. The reader may consult Stinson (2004) for a standard treatment of these objects. Ionin and Shrikhande (2006) studies these objects and their applications in constructing other configurations.
14. Note the similarity to the redundancy introduced in constructing the error-correcting codes of §2.
15. Recall that a Hilbert space is a complete normed linear space where the norm is induced by an inner product. Halmos (1982) and Halmos (1998) are standard references on the subject of Hilbert spaces.
16. There is some ambiguity in defining frames over  $\mathbf{H}^k$ . As  $\mathbf{H}$  is a skew-field, we can take  $\mathbf{H}^k$  to be either a left- or right-module. This discussion is taken up in detail in Waldron (2020), where the right-module is always used.
17. There are different ways of defining unbiased weighing matrices. Classically, two unitary weighing matrices  $W(n, k)$ , say  $H$  and  $K$ , are *unbiased* if  $HK^* = \sqrt{k}L$ , for some unitary weighing matrix  $L$  of weight  $k$ . In Nozaki and Suda (2015), the weighing matrices  $H$  and  $K$  above are *quasi-unbiased* with parameters  $(n, k, \ell, a)$  if  $a^{-\frac{1}{2}}HK^*$  is a  $W(n, \ell)$ . It is this idea of quasi-unbiasedness that is most easily extended to orthogonal designs.





# 3

## **A New Family of Balanced Weighing Matrices and Association Schemes**

This chapter serves as a particular application of a more general construction to be given in the following chapter. Here a new family of balanced weighing matrices are constructed, and an equivalence to certain association schemes is developed. The work shown here is a result of Kharaghani et al. (2022).

### **§8. Generalized Kronecker Product**

In this section, we briefly review the Kronecker product and a few of its properties and applications, namely, we will see its applications to regular Hadamard matrices. Following this, we will consider a simple generalization of the Kronecker product in order to allow particular bijections to act on a matrix.

\* \* \*

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

**8.1. Definitions.** Let  $A$  be an  $n \times m$  matrix, and let  $B$  be an  $s \times t$  matrix, each with entries from some field. The *Kronecker product* of  $A$  by  $B$  is the  $ns \times mt$  matrix given by  $A \otimes B = (A_{ij}B)$ . This noncommutative matrix product has the following immediate properties wherever the sizes of the matrices make sense.

**8.1. Lemma.** Let  $A, B, C$  and  $D$  be matrices over some field  $F$ , and let  $\lambda \in F$ . Then:

$$(8.1.a) \quad \lambda A \otimes B = A \otimes \lambda B = \lambda(A \otimes B),$$

$$(8.1.b) \quad A \otimes B = (A \times I)(I \otimes B) = (I \otimes B)(A \otimes I),$$

$$(8.1.c) \quad (A + B) \otimes C = (A \otimes C) + (B \otimes C),$$

$$(8.1.d) \quad A \otimes (B + C) = (A \otimes B) + (A \otimes C),$$

$$(8.1.e) \quad (A \otimes B)(C \otimes D) = AC \otimes BD,$$

$$(8.1.f) \quad (A \otimes B)^t = A^t \otimes B^t, \text{ and}$$

$$(8.1.g) \quad \text{if } A \text{ is } n \times n \text{ and } B \text{ is } m \times m, \text{ then } \det(A \otimes B) = [\det(A)]^m [\det(B)]^n.$$

There are a number of immediate applications of the Kronecker product to what we have done so far. For instance, if  $H$  is a  $W(n_1, k_1)$  and  $K$  a  $W(n_2, k_2)$ , then  $H \otimes K$  is a  $W(n_1 n_2, k_1 k_2)$ .

**8.2. Example.** Let  $H$  be the Hadamard matrix of order 4 given by

$$(8.2.a) \quad \begin{pmatrix} - & + & + & + \\ + & - & + & + \\ + & + & - & + \\ + & + & + & - \end{pmatrix}.$$

Then  $H \otimes H$  is the Hadamard matrix of 16 given by

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

$$(8.2.b) \quad \begin{pmatrix} + & - & - & - & - & + & + & + & - & + & + & + & - & + & + & + \\ - & + & - & - & + & - & + & + & + & - & + & + & + & - & + & + \\ - & - & + & - & + & + & - & + & + & + & - & + & + & + & - & + \\ - & - & - & + & + & + & + & - & + & + & + & - & + & + & + & - \\ - & + & + & + & + & - & - & - & - & + & + & + & - & + & + & + \\ + & - & + & + & - & + & - & - & + & - & + & + & + & - & + & + \\ + & + & - & + & - & - & + & - & + & + & - & + & + & + & - & + \\ + & + & + & - & - & - & - & + & + & + & + & - & + & + & + & - \\ - & + & + & + & - & + & + & + & + & - & - & - & - & + & + & + \\ + & - & + & + & + & - & + & + & - & + & - & - & + & - & + & + \\ + & + & - & + & + & + & - & + & - & - & + & - & + & + & - & + \\ + & + & + & - & + & + & + & - & - & - & - & + & + & + & + & - \\ - & + & + & + & - & + & + & + & - & + & + & + & + & - & - & - \\ + & - & + & + & + & - & + & + & + & - & + & + & - & + & - & - \\ + & + & - & + & + & + & - & + & + & + & - & + & - & - & + & - \\ + & + & + & - & + & + & + & - & + & + & + & - & - & - & - & + \end{pmatrix},$$

where the block structure is evident <sup>18</sup>).

As one further application of the standard Kronecker product before we move on, we will consider regular Hadamard matrices.

**8.3. Definition.** A Hadamard matrix is *row regular* (or *column regular*) if it's rows (resp. columns) have a constant sum. A Hadamard matrix is *regular* if it is both row and column regular.

It isn't difficult to see that a Hadamard matrix is row regular or column regular if and only if it is regular. Moreover, if the constant row (column) sum is  $s$ , then the order of the matrix is  $s^2$  (see Stinson, 2004, Chapter 4). By the definition of the standard Kronecker product, we also see that if  $H$  and  $K$  are regular Hadamard matrices, then so is  $H \otimes K$ .

Example 8.2 has evinced the existence of a regular  $W(4, 4)$ . It is also known that there is a regular  $W(36, 36)$  (see Stinson, 2004, Chapter 4). Also, if there is a Hadamard matrix of order  $n$ , then there is a symmetric, regular, Hadamard matrix with constant block diagonal of order  $n^2$  (see Colbourn and Dinitz, 2007, Part V). We then have the following result.

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

**8.4. Proposition.** If there exist Hadamard matrices of orders  $n_1, \dots, n_k$ , then there is a regular Hadamard matrix of order  $4^a 9^b n_1^2 \cdots n_k^2$ , for  $a, b \in \mathbf{Z}_+$  and  $a \geq b$ .

Having convinced ourselves of the utility of the standard Kronecker product, we generalize in the following way. Let  $\mathcal{M}$  be a collection of  $n \times m$  matrices with entries from some commutative ring  $R$ , and let  $\Xi$  be an  $s \times t$  matrix over the collection of endofunctions of  $\mathcal{M}$ . For each  $A \in \mathcal{M}$ , the  $ns \times mt$  matrix  $\Xi \otimes A$  over  $R$  is defined as  $(\Xi_{ij}(A))$ . If certain properties of  $A$  are left invariant under the elements of  $\Xi$ , then these properties will be reflected in the block structure of  $\Xi \otimes A$ .

With the above ideas in mind, if we have a collection  $\mathcal{M}$  of objects, we desire a subset of the collection of endofunctions of  $\mathcal{M}$  that are property preserving. If  $\mathcal{M}$  is a collection of incidence structures, then a set of bijections of  $\mathcal{M}$  that preserve incidence is called a *group of symmetries* of  $\mathcal{M}$ . If  $\Xi$  is a BGW over a group of symmetries, then  $\Xi \otimes A$  has both inter-block regularities and intra-block regularities.

\* \* \*

**8.2. A First Application: Block designs.** In the previous subsection, we began a brief discussion of groups of symmetries of incidence structures. As an example, let  $\mathbf{D} = (X, \mathcal{B})$  be a BIBD( $v, b, r, k, \lambda$ ), and let  $\mathcal{B} = \bigcup_{i=1}^m \mathcal{B}_i$  be a partition of the blocks of  $\mathbf{D}$ . If  $G_i$  acts on  $\mathcal{B}_i$ , then,  $(X, \bigcup_{i=1}^m \mathcal{B}_i g_i)$ , for  $g_i \in G_i$ , is again a BIBD( $v, b, r, k, \lambda$ ). Let  $A = A(\mathbf{D})$  be the incidence matrix of  $\mathbf{D}$ , and assume that  $A = (A(X, \mathcal{B}_1) \cdots A(X, \mathcal{B}_m))$ . Then  $G = \prod_i G_i$  acts naturally on the columns of  $A$ . If in addition  $G_i$  is sharply transitive on  $\mathcal{B}_i$ , and if each point  $x \in X$  appears  $r_i$  times in  $\mathcal{B}_i$ , then

$$|G|^{-1} \sum_{g \in G} Ag = (r_1 b_1^{-1} J_{v \times b_1} \cdots r_m b_m^{-1} J_{v \times b_m}),$$

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

where  $b_i = |\mathcal{B}_i|$ . Therefore, if  $r_i b_j = r_j b_i$ , for all  $i, j \in \{1, \dots, m\}$ , then  $\sum_{g \in G} Ag$  is an integer multiple of  $J_{v \times b}$  since  $b_i \mid |G|$ . Since  $G$  acts on the columns of  $A$ , we also have that  $Xg(Yg)^t = XY^t$ , for each  $X, Y \in AG$ .

This motivates the following. If  $G$  is a group of symmetries of a collection  $\mathcal{M}$  of  $\text{BIBD}(v, b, r, k, \lambda)$ s such that  $\sum_{g \in G} Xg = \alpha_X J$ , for some  $\alpha_X \in \mathbb{N}$  and all  $X \in \mathcal{M}$ ; and if  $Xg(Yg)^t = XY^t$ , for all  $X, Y \in \mathcal{M}$ ; then we say that  $G$  is an *admissible* group of symmetries of  $\mathbf{D}$ . We then have the following.

**8.5. Theorem.** Let  $\mathcal{M}$  be a collection of  $\text{BIBD}(v, b, r, k, \lambda)$ s, and let  $G$  be an admissible group of symmetries of  $\mathcal{M}$ . If  $\Xi$  is a  $\text{BGW}(w, \ell, \mu; G)$  such that  $kr\mu = v\lambda\ell$ , then  $\Xi \otimes X$  is a  $\text{BIBD}(vw, bw, r\ell, k\ell, \lambda\ell)$ , for any  $X \in \mathcal{M}$ .

**Proof.** Straightforward calculation. See Theorem 2.4 of Ionin (2001) for details. ■

**8.6. Corollary.** If  $X$  is quasi-residual, then so is  $\Xi \otimes X$ .

**8.7. Example.** Let  $q = p^n$  be a prime power, and let  $H$  be a  $\text{GH}(q, 1)$  over  $\text{EA}(q)$  where the elements have the usual representation of  $(0, 1)$ -matrices. Then it isn't difficult to see that  $A = (I_q \otimes \mathbf{1}_q \ H)$  is a  $\text{BIBD}(q^2, q + q^2, 1 + q, q, 1)$ . Importantly, the columns are placed into  $1 + q$  consecutive disjoint groups of  $q$  columns each such that each point appears precisely once in every group<sup>19</sup>). Let  $G$  be the group which cyclically permutes the blocks in each group so that  $|G| = q$ .

If  $1 + q$  is a prime power, then there is a  $\text{BGW}(2 + q, 1 + q, q; C_q)$ , say  $\Xi$ , where the group elements cyclically permute the blocks of each partition class in the above design. The parametric conditions of the theorem are met, hence  $\Xi \otimes A$  is a  $\text{BIBD}(q^2(2 + q), (q + q^2)(2 + q), (1 + q)^2, q(1 + q), 1 + q)$ . More generally, we can take  $\Xi$  to be any  $\text{BGW}((p^{n+1} - 1)/(p - 1), p^n, p^n - p^{n-1}; C_q)$ , where  $p = 1 + q$ .

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

\* \* \*

**8.3. A Second Application: Bhaskar Rao designs.** It isn't difficult to extend the ideas of Theorem 8.5 to the more general Bhaskar Rao designs. In order to accomplish this, however, we need to extend the idea of admissible groups of symmetries.

Let  $\mathcal{M}$  be a collection of  $\text{GBRD}(v, b, r, k, \lambda; H)$ s. Then a group of bijections  $G$  of  $\mathcal{M}$  is an admissible group of symmetries if (a)  $\sum_{g \in G} Xg = \alpha_X HJ$ , for some  $\alpha_X \in \mathbb{N}$  and every  $X \in \mathcal{M}$ , and (b)  $Xg(Yg)^* = XY^*$ , for every  $X, Y \in \mathcal{M}$ .

The following is then a simple generalization of Theorem 8.5

**8.8. Theorem.** Let  $\mathcal{M}$  be a collection of  $\text{GBRD}(v, b, r, k, \lambda; H)$ s, and let  $G$  be an admissible group of symmetries of  $\mathcal{M}$ . If  $\Xi$  is a  $\text{BGW}(w, \ell, \mu; G)$  such that  $kr\mu = v\lambda\ell$ , then  $\Xi \otimes X$  is a  $\text{GBRD}(vw, bw, r\ell, k\ell, \lambda\ell; H)$ , for any  $X \in \mathcal{M}$ .

**8.9. Example.** The following example was noted in Pender (2020). The following  $\text{BGW}(15, 7, 3; C_3)$  was found by computational means in Gibbons and Mathon (1987)

$$(8.9.a) \quad \begin{pmatrix} 0 & 3 & 3 & 3 & 3 & 3 & 3 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 3 & 2 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 2 & 2 & 3 \\ 0 & 0 & 3 & 2 & 0 & 1 & 0 & 0 & 3 & 0 & 0 & 2 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 2 & 0 & 1 & 0 & 2 & 3 & 0 & 0 & 2 & 0 & 2 \\ 0 & 0 & 0 & 0 & 3 & 2 & 0 & 1 & 2 & 2 & 3 & 0 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 & 0 & 3 & 2 & 0 & 0 & 2 & 2 & 3 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 & 0 & 0 & 3 & 2 & 2 & 0 & 2 & 2 & 3 & 0 & 0 \\ 0 & 2 & 0 & 1 & 0 & 0 & 0 & 3 & 0 & 2 & 0 & 2 & 2 & 3 & 0 \\ 3 & 3 & 1 & 0 & 2 & 0 & 0 & 0 & 3 & 2 & 0 & 1 & 0 & 0 & 0 \\ 3 & 0 & 3 & 1 & 0 & 2 & 0 & 0 & 0 & 3 & 2 & 0 & 1 & 0 & 0 \\ 3 & 0 & 0 & 3 & 1 & 0 & 2 & 0 & 0 & 0 & 3 & 2 & 0 & 1 & 0 \\ 3 & 0 & 0 & 0 & 3 & 1 & 0 & 2 & 0 & 0 & 0 & 3 & 2 & 0 & 1 \\ 3 & 2 & 0 & 0 & 0 & 3 & 1 & 0 & 1 & 0 & 0 & 0 & 3 & 2 & 0 \\ 3 & 0 & 2 & 0 & 0 & 0 & 3 & 1 & 0 & 1 & 0 & 0 & 0 & 3 & 2 \\ 3 & 1 & 0 & 2 & 0 & 0 & 0 & 3 & 2 & 0 & 1 & 0 & 0 & 0 & 3 \end{pmatrix},$$

where the nonzero entries are logarithms of some generator of  $C_3$ . Note that the core of (8.9.a) is composed of 4 circulant matrices. Taking  $R$  to be the residual part of (8.9.a), and letting  $g = \begin{pmatrix} O & I_7 \\ \omega I_7 & O \end{pmatrix}$ , we see that  $G = \langle g \rangle$  forms an admissible

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

group of symmetries for  $R$ . For any  $n \in N$ , there is a  $\text{BGW}((7^{n+1} - 1)/6, 7^n, 6 \cdot 7^{n-1}; G)$  which satisfy the parametric conditions of Theorem 8.8. Therefore, for any  $n > 0$ , there is a  $\text{GBRD}(4(7^{n+1} - 1)/3, 2 \cdot 7^{n+1}, 7^{n+1}, 4 \cdot 7^n, 3 \cdot 7^n; C_3)$ .

This concludes our introduction to the generalized Kronecker product.

## §9. A New Family of Balanced Weighing Matrices

Here the generalized Kronecker product of the previous subsection will be put to use in constructing a new family of balanced weighing matrices. Additionally, the simplex codes appear again and are applied in the construction. The construction presented here is indicative of a general method to be presented in the following chapter.

\* \* \*

**9.1. Lemmata.** In §2.2, we introduced the linear simplex code  $\mathcal{S}_{q,n}$ . There it was shown that the code had constant weight  $q^{n-1}$ ; in particular, it follows that it is equidistant with constant Hamming distance  $q^{n-1}$  since the code is linear.

In Rajkundlia (1983), and later reproduced in Ionin (2001) using the language of BGW matrices, generalized Hadamard matrices  $\text{GH}(q, q^{n-1})$  were used recursively in conjunction with the classical parameter  $\text{BGW}((q^n - 1)/(q - 1), q^{n-1}, q^{n-1} - q^{n-2}; \text{GF}(q)^*)$ s in order to construct certain designs. It turns out that the  $\text{GH}(q, q^{n-1})$  used in the construction can be replaced by  $\mathcal{S}_{q,n}$ , and so simplify the construction.

In order to apply the linear code  $\mathcal{S}_{q,n}$ , we will require the following lemma.

**9.1. Lemma.** Let  $\text{GF}(q) = \{a_0 = 0, a_1, \dots, a_{q-1}\}$ , and let  $n > 1$ . Then there exist disjoint  $(0, 1)$ -matrices  $A_{a_1}, \dots, A_{a_{q-1}}$  of dimensions  $q^n \times (q^n - 1)/(q - 1)$



### 3. A New Family of Balanced Weighing Matrices and Association Schemes

such that  $\mathcal{S}_{q,n} = \sum_{\alpha \in \text{GF}(q)^*} \alpha A_\alpha$ . If we define  $A_0 = J - \sum_{\alpha \in \text{GF}(q)^*} A_\alpha$ , then the following hold.

$$(9.1.a) \quad \sum_{\alpha \in \text{GF}(q)} A_\alpha A_\alpha^t = \frac{q^{n-1}-1}{q-1} J + q^{n-1} I, \text{ and}$$

$$(9.1.b) \quad \sum_{\substack{\alpha, \beta \in \text{GF}(q) \\ \alpha \neq \beta}} A_\alpha A_\beta^t = q^{n-1} (J - I).$$

**Proof.** Labeling the rows of  $\mathcal{S}_{q,n}$  by  $r_0, \dots, r_{q^n-1}$ , and taking  $A = \mathcal{S}_{q,n}$ , we then have that

$$\begin{aligned} \left( \sum_{\alpha \in \text{GF}(q)} A_\alpha A_\alpha^t \right)_{ij} &= \sum_{\alpha \in \text{GF}(q)} (A_\alpha A_\alpha^t)_{ij} \\ &= \sum_{\alpha \in \text{GF}(q)} \sum_{\ell=0}^{\frac{q(q^{n-1}-1)}{q-1}} (A_\alpha)_{i\ell} (A_\alpha)_{j\ell} \\ &= \sum_{\alpha \in \text{GF}(q)} \# \{ \ell \in \{0, \dots, \frac{q(q^{n-1}-1)}{q-1}\} : A_{i\ell} = A_{j\ell} = \alpha \} \\ &= \# \{ \ell \in \{0, \dots, \frac{q(q^{n-1}-1)}{q-1}\} : A_{i\ell} = A_{j\ell} \} \\ &= \frac{q^n - 1}{q - 1} - \text{dist}(r_i, r_j), \end{aligned}$$

which shows (9.1.a).

Since  $\sum_{\alpha \in \text{GF}(q)} A_\alpha = J$ , it follows that  $\sum_{\alpha, \beta} A_\alpha A_\beta^t = (\sum_{\alpha} A_\alpha) (\sum_{\beta} A_\beta)^t = \frac{q^n-1}{q-1} J$ , and (9.1.b) has been proven. ■

If  $W$  is a BGW( $v, k, \lambda; C_n$ ) over some cyclic group  $C_n = \{1, g, \dots, g^{n-1}\}$  of order  $n$ , then there are  $n$  disjoint  $(0, 1)$ -matrices  $W_0, \dots, W_{n-1}$  such that  $W = W_0 + gW_1 + \dots + g^{n-1}W_{n-1}$ . We call  $W_0$  and  $W_1$  the *decomposition matrices* of the weighing matrix. Because  $W$  is a BGW matrix, we have the following lemma.

#### 9.2. Lemma.

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

$$(9.2.a) \sum_{i,j} g^{i-j} W_i W_j^t = \sum_{i,j} g^{i-j} W_j^t W_i = kI + \frac{\lambda}{n} \left( \sum_i g_i \right) (J - I),$$

$$(9.2.b) \sum_i W_i W_i^t = \sum_i W_i^t W_i = kI + \frac{\lambda}{n} (J - I), \text{ and}$$

$$(9.2.c) \sum_i W_i W_{i+j}^t = \sum_i W_{i+j}^t W_i = \frac{\lambda}{n} (J - I), \text{ for } j \in \{1, \dots, n-1\}.$$

**Proof.** (9.2.a) is simply a restatement of the fact that both  $W$  and  $W^*$  are  $\text{BGW}(v, k, \lambda; C_n)$ s (see Ionin and Shrikhande, 2006). (9.2.b) follows by noting that there are  $k$  nonzero entries in every row of  $W$ , and that 1 appears  $\lambda/n$  times in the conjugate inner product between distinct rows. Similarly, (9.2.c) follows by noting that each nonidentity element of the group appears  $\lambda/n$  times in the conjugate inner product between distinct rows of  $W$ , and that  $i \neq i+j$ , for each  $i$  whenever  $j \not\equiv 0 \pmod{n}$ . ■

Now, consider the balanced  $W(19, 9)$  shown to exist by computational means in de Launey and Sarvate (1984).

$$W_{19} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & + & + & + & + & + & + & + & + \\ 0 & - & 0 & 0 & + & 0 & + & + & + & 0 & 0 & 0 & 0 & + & 0 & - & + & - & 0 \\ 0 & 0 & - & 0 & + & + & 0 & 0 & + & + & 0 & 0 & 0 & - & + & 0 & 0 & + & - \\ 0 & 0 & 0 & - & 0 & + & + & + & 0 & + & 0 & 0 & 0 & 0 & - & + & - & 0 & + \\ 0 & + & + & 0 & - & 0 & 0 & + & 0 & + & + & - & 0 & 0 & 0 & 0 & + & 0 & - \\ 0 & 0 & + & + & 0 & - & 0 & + & + & 0 & 0 & + & - & 0 & 0 & 0 & - & + & 0 \\ 0 & + & 0 & + & 0 & 0 & - & 0 & + & + & - & 0 & + & 0 & 0 & 0 & 0 & - & + \\ 0 & + & 0 & + & + & + & 0 & - & 0 & 0 & + & 0 & - & + & - & 0 & 0 & 0 & 0 \\ 0 & + & + & 0 & 0 & + & + & 0 & - & 0 & - & + & 0 & 0 & + & - & 0 & 0 & 0 \\ 0 & 0 & + & + & + & 0 & + & 0 & 0 & - & 0 & - & + & - & 0 & + & 0 & 0 & 0 \\ + & 0 & 0 & 0 & + & 0 & - & + & - & 0 & 0 & - & - & 0 & + & 0 & 0 & 0 & + \\ + & 0 & 0 & 0 & - & + & 0 & 0 & + & - & - & 0 & - & 0 & 0 & + & + & 0 & 0 \\ + & 0 & 0 & 0 & 0 & - & + & - & 0 & + & - & - & 0 & + & 0 & 0 & 0 & + & 0 \\ + & + & - & 0 & 0 & 0 & 0 & + & 0 & - & 0 & 0 & + & 0 & - & - & 0 & + & 0 \\ + & 0 & + & - & 0 & 0 & 0 & - & + & 0 & + & 0 & 0 & - & 0 & - & 0 & 0 & + \\ + & - & 0 & + & 0 & 0 & 0 & 0 & - & + & 0 & + & 0 & - & - & 0 & + & 0 & 0 \\ + & + & 0 & - & + & - & 0 & 0 & 0 & 0 & 0 & + & 0 & 0 & 0 & + & 0 & - & - \\ + & - & + & 0 & 0 & + & - & 0 & 0 & 0 & 0 & 0 & + & + & 0 & 0 & - & 0 & - \\ + & 0 & - & + & - & 0 & + & 0 & 0 & 0 & + & 0 & 0 & 0 & + & 0 & - & - & 0 \end{pmatrix}.$$

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

Take  $R_1$  and  $D$  to be the residual and derived parts, respectively, of  $W_{19}$ . Define the matrix  $|R_1|$  by  $|R_1|_{ij} = |R_{1ij}|$ . Then  $|R_1|$  is the incidence matrix of a residual BIBD(10, 18, 9, 5, 4), hence  $|R_2| = J - |R_1|$  is a BIBD with the same parameters. Moreover,  $|R_2|$  is residual since  $|R_2|$  together with  $|D|$  also forms a symmetric design. We therefore seek a signing of  $|R_2|$  over  $\{-1, 1\}$ . This search was conducted via Maple, and the following signing was produced.

$$R_2 = \begin{pmatrix} + & + & + & + & + & + & + & + & + & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & - & - & 0 & + & 0 & 0 & 0 & + & - & + & + & 0 & + & 0 & 0 & 0 & + \\ - & 0 & - & 0 & 0 & + & + & 0 & 0 & + & - & + & 0 & 0 & + & + & 0 & 0 \\ - & - & 0 & + & 0 & 0 & 0 & + & 0 & + & + & - & + & 0 & 0 & 0 & + & 0 \\ 0 & 0 & + & 0 & - & - & 0 & + & 0 & 0 & 0 & + & - & + & + & 0 & + & 0 \\ + & 0 & 0 & - & 0 & - & 0 & 0 & + & + & 0 & 0 & + & - & + & 0 & 0 & + \\ 0 & + & 0 & - & - & 0 & + & 0 & 0 & 0 & + & 0 & + & + & - & + & 0 & 0 \\ 0 & + & 0 & 0 & 0 & + & 0 & - & - & 0 & + & 0 & 0 & 0 & + & - & + & + \\ 0 & 0 & + & + & 0 & 0 & - & 0 & - & 0 & 0 & + & + & 0 & 0 & + & - & + \\ + & 0 & 0 & 0 & + & 0 & - & - & 0 & + & 0 & 0 & 0 & + & 0 & + & + & - \end{pmatrix}.$$

Remarkably,  $R_2$  together with  $D$  also forms a balanced  $W(19, 9)$ . The matrices  $R_1$ ,  $R_2$ , and  $D$  then satisfy several properties.

#### 9.3. Lemma.

$$(9.3.a) \quad R_1 R_1^t = R_2 R_2^t = I, \quad R_1 R_2^t = R_2 R_1^t;$$

$$(9.3.b) \quad D D^t = 9I - J;$$

$$(9.3.c) \quad R_1 D^t = R_2 D^t = O;$$

$$(9.3.d) \quad |R_1| |R_1|^t = |R_2| |R_2|^t = 5I + 4J, \quad |R_1| |R_2|^t = |R_2| |R_1|^t = 5(J - I);$$

and

$$(9.3.e) \quad |D| |D|^t = 5I + 3J, \quad |R_1| |D|^t = |R_2| |D|^t = 4J.$$

**Proof.** Restatement of the facts that  $R_1$ ,  $R_2$ , and  $D$  form balanced  $W(19, 4)$  weighing matrices. ■

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

\* \* \*

**9.2. Construction.** Having the lemmata of the previous subsection at our disposal, we are ready to present the construction of a new family of balanced weighing matrices. We desire to apply BGWs in the construction of these matrices, so we need an admissible group of symmetries. Take  $\mathcal{M} = \{R_1, R_2\}$ ; then it isn't difficult to see that  $-R_2 \mapsto -R_1 \mapsto R_2 \mapsto R_1 \mapsto -R_2$  is an admissible cyclic group of symmetries of order 4 for  $\mathcal{M}$ —though, this will be derived explicitly below.

Let  $n > 1$ , and take  $\Xi$  to be a BGW $((9^n - 1)/8, 9^{n-1}, 9^{n-1} - 9^{n-2}; C_4)$ . We claim that  $\Xi \otimes R_1$  is the residual part of a balanced W $([9(9^n - 1)/4] + 1, 9^n)$ . Note there are disjoint  $(0, 1)$ -matrices  $\Xi_0, \Xi_1, \Xi_2, \Xi_3$  such that  $\Xi = \Xi_0 + g\Xi_1 + g^2\Xi_2 + g^3\Xi_3$  if  $C_4 = \{e, g, g^2, g^3\}$ . Then  $\Xi \otimes R_1 = \Xi_0 \otimes R_1 - \Xi_1 \otimes R_2 - \Xi_2 \otimes R_1 + \Xi_3 \otimes R_2$ .

Next, let  $\mathcal{S}_{9,n} = \sum_{\alpha \in \text{GF}(9)^*} \alpha A_\alpha$ , and define  $A_0 = J - \sum_{\alpha \in \text{GF}(9)^*} A_\alpha$ . Finally, take  $\Theta = \sum_{\alpha \in \text{GF}(9)} A_\alpha \otimes D$ . It will be shown that  $\Theta$  is the derived part of a balanced W $([9(9^n - 1)/4] + 1, 9^n)$ .

We require the following lemma.

#### 9.4. Lemma.

$$(9.4.a) \quad (\Xi \otimes R_1)(\Xi \otimes R_1)^t = 9^n I;$$

$$(9.4.b) \quad \Theta \Theta^* = 9^n I - J;$$

$$(9.4.c) \quad (\Xi \otimes R_1) \Theta^t = \Theta (\Xi \otimes R_1)^t = O;$$

$$(9.4.d) \quad |\Xi \otimes R_1| |\Xi \otimes R_1|^t = 5 \cdot 9^n I + 4 \cdot 9^n J;$$

$$(9.4.e) \quad |\Theta| |\Theta|^t = 5 \cdot 9^n I + (4 \cdot 9^n - 1) J; \text{ and}$$

$$(9.4.f) \quad |\Xi \otimes R_1| |\Theta|^t = |\Theta| |\Xi \otimes R_1|^t = 4 \cdot 9^n J.$$

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

**Proof.** By Lemma 9.2 and (9.3.a),

$$\begin{aligned}
(\Xi \otimes R_1)(\Xi \otimes R_1) &= 9\Xi_0\Xi_0^t \otimes I - \Xi_0\Xi_1^t \otimes R_1R_2^t - 9\Xi_0\Xi_2^t \otimes I + \Xi_0\Xi_3^t \otimes R_1R_2^t \\
&\quad - \Xi_1\Xi_0^t \otimes R_2R_1^t + 9\Xi_1\Xi_1^t \otimes I + \Xi_1\Xi_2^t \otimes R_2R_1^t - 9\Xi_1\Xi_3^t \otimes I \\
&\quad - 9\Xi_2\Xi_0^t \otimes I + \Xi_2\Xi_1^t \otimes R_1R_2^t + 9\Xi_2\Xi_2^t \otimes I - \Xi_2\Xi_3^t \otimes R_1R_2^t \\
&\quad + \Xi_3\Xi_0^t \otimes R_2R_1^t - 9\Xi_3\Xi_1^t \otimes I - \Xi_3\Xi_2^t \otimes R_2R_1^t + 9\Xi_3\Xi_3^t \otimes I \\
&= 9 \sum_i (\Xi_i\Xi_i^t - \Xi_i\Xi_{i+2}^t) \otimes I - \sum_i (\Xi_i\Xi_{i+1} - \Xi_i\Xi_{i+3}) \otimes R_1R_2^t \\
&= 9^n I,
\end{aligned}$$

and (9.4.a) is shown.

Next, by Lemma 9.1 and (9.3.b), and upon indexing the rows of  $D$  by elements of  $\text{GF}(9)$ ,

$$\begin{aligned}
\Theta\Theta^t &= \sum_{\alpha, \beta \in \text{GF}(9)} A_\alpha A_\beta^t \otimes r_\alpha r_\beta^t \\
&= \sum_{\alpha \in \text{GF}(9)} A_\alpha A_\alpha^t \otimes r_\alpha r_\alpha^t + \sum_{\alpha \neq \beta} A_\alpha A_\beta^t \otimes r_\alpha r_\beta^t \\
&= 8 \sum_{\alpha \in \text{GF}(9)} A_\alpha A_\alpha^t - \sum_{\alpha \neq \beta} A_\alpha A_\beta^t \\
&= (9^{n-1} - 1)J + 8 \cdot 9^{n-1}I - 9^{n-1}(J - I) \\
&= 9^n I - J,
\end{aligned}$$

which shows (9.4.b).

By Lemma (9.3.c),

$$\begin{aligned}
(\Xi \otimes R_1)\Theta^t &= \sum_{\alpha \in \text{GF}(9)} (\Xi_0 A_\alpha^t \otimes R_1 r_\alpha^t - \Xi_1 A_\alpha^t \otimes R_2 r_\alpha^t - \Xi_2 A_\alpha^t \otimes R_2 r_\alpha^t + \Xi_3 A_\alpha^t \otimes R_2 r_\alpha^t) \\
&= O,
\end{aligned}$$

and (9.4.c) has been shown.

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

Since  $|\Xi \otimes R_1| = (\Xi_0 + \Xi_2) \otimes |R_1| + (\Xi_1 + \Xi_3) \otimes |R_2|$ , (9.4.d) is shown similarly to (9.4.a).

(9.4.e) is shown just as (9.4.b) after noting that  $|\Theta| = \sum_{\alpha \in \text{GF}(9)} A_\alpha \otimes |r_\alpha|$ .

Finally, (9.4.f) is shown precisely as in (9.4.c). ■

We are now ready to present the main construction.

**9.5. Theorem.** Given  $\Xi \otimes R_1$  and  $\Theta$  defined above,

$$(9.5.a) \quad \begin{pmatrix} \mathbf{0} & \Xi \otimes R_1 \\ 1 & \Theta \end{pmatrix}$$

is a balanced  $W([9(9^n - 1)/4] + 1, 9^n)$ .

**Proof.** By the lemma,  $(\Xi \otimes R_1)(\Xi \otimes R_1)^t = 9^n I$ ,  $\Theta \Theta^t = 9^n I - J$ , and  $(\Xi \otimes R_1)\Theta^t = O$ ; thus, (9.5.a) is a weighing matrix with the appropriate parameters. It remains to show it is balanced. But the lemma again gives  $|\Xi \otimes R_1||\Xi \otimes R_1|^t = 5 \cdot 9^n I + 4 \cdot 9^n J$ ,  $|\Theta||\Theta|^t = 5 \cdot 9^n I + (4 \cdot 9^n - 1)J$ , and  $|\Xi \otimes R_1|\Theta^t = 4 \cdot 9^n J$ . We have then shown that (9.5.a) is balanced, and the proof is complete. ■

## §10. Weighing Matrices and Association Schemes

In Brouwer et al. (1989) it is shown that symmetric designs are equivalent to certain 3-class association schemes. It is a natural question to ask whether or not a balanced weighing matrix could be related to this scheme, particularly an augmentation. This is the goal of this subsection.

\* \* \*

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

**10.1. Adjacency Matrices.** Assume the existence of a balanced  $W(v, k)$ , say  $W = W_0 - W_1$ , where  $W_0$  and  $W_1$  are disjoint  $(0, 1)$ -matrices, and define the following matrices where  $P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

$$\begin{aligned} A_0 &= I_{4v}, \\ A_1 &= I_2 \otimes P \otimes I_v, \\ A_2 &= I_2 \otimes J_2 \otimes (J_v - I_v), \\ A_3 &= \begin{pmatrix} O & I_2 \otimes W_0 + P \otimes W_1 \\ I_2 \otimes W_0^t + P \otimes W_1^t & O \end{pmatrix}, \\ A_4 &= \begin{pmatrix} O & I_2 \otimes W_1 + P \otimes W_0 \\ I_2 \otimes W_1^t + P \otimes W_0^t & O \end{pmatrix}, \\ A_5 &= \begin{pmatrix} O & J_2 \otimes (J_v - W_0 - W_1) \\ J_2 \otimes (J_v - W_0^t - W_1^t) & O \end{pmatrix}. \end{aligned}$$

We claim that  $\{A_0, A_1, A_2, A_3, A_4, A_5\}$  form a 5-class symmetric association scheme. Clearly, (4.2.a), (4.2.b), and (4.2.f) are satisfied. It remains to show closure under multiplication.

First,  $A_1^2 = (I_2 \otimes P \otimes I_v)^2 = I_2 \otimes P^2 \otimes I_v = I = A_0$ . Next, since

$$\begin{aligned} [J_2 \otimes (J_v - I_v)]^2 &= 2J_2 \otimes [(v-1)I_v + (v-2)(J_v - I_v)] \\ &= 2(v-1)J_2 \otimes I_v + 2(v-2)J_2 \otimes (J_v - I_v) \\ &= 2(v-1)(I_2 + P) \otimes I_v + 2(v-2)J_2 \otimes (J_v - I_v), \end{aligned}$$

it follows that  $A_2^2 = 2(v-1)(A_0 + A_1) + 2(v-2)A_2$ . By (9.2.b) and (9.2.c), we

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

find

$$\begin{aligned}
(I_2 \otimes W_0 + P \otimes W_1)(I_2 \otimes W_0^t + P \otimes W_1^t) &= I_2 \otimes (W_0 W_0^t + W_1 W_1^t) \\
&\quad + P \otimes (W_0 W_1^t + W_1 W_0^t) \\
&= I_2 \otimes [kI_v + \frac{\lambda}{2}(J_v - I_v)] + P \otimes \frac{\lambda}{2}(J_v - I_v) \\
&= kI_{2v} + \frac{\lambda}{2}J_2 \otimes (J_v - I_v),
\end{aligned}$$

where  $\lambda = k(k-1)/(v-1)$ . Therefore,  $A_3^3 = A_4^2 = kA_0 + \frac{\lambda}{2}A_2$ . Finally,

$$\begin{aligned}
J_2^2 \otimes [J_v - W_0 - W_1](J_v - W_0^t - W_1^t) &= J_2^2 \otimes (J_v^2 - 2J_v(W_0 + W_1) \\
&\quad + W_0 W_0^t + W_1 W_1^t + W_0 W_1^t + W_1 W_0^t) \\
&= J_2 \otimes [2(v-k)I_v + 2(v-2k+\lambda)(J_v - I_v)],
\end{aligned}$$

hence  $A_5^2 = 2(v-k)(A_0 + A_1) + 2(v-2k+\lambda)A_1$ . We next show that  $A_i A_j \in$

$\langle A_0, \dots, A_5 \rangle$  whenever  $i \neq j$ .

Note  $(P \otimes I_v)[J_2 \otimes (J_v - I_v)] = J_2 \otimes (J_v - I_v)$  so that  $A_1 A_2 = A_2 A_1 = A_2$ .

Then

$$\begin{aligned}
(I_2 \otimes W_0 + P \otimes W_1)(I_2 \otimes W_1^t + P \otimes W_0^t) &= I_2 \otimes (W_0 W_1^t + W_1 W_0^t) + P \otimes (W_0 W_0^t + W_1 W_1^t) \\
&= I_2 \otimes \frac{\lambda}{2}(J_v - I_v) + P \otimes [kI_v + \frac{\lambda}{2}(J_v - I_v)] \\
&= \frac{\lambda}{2}J_2 \otimes (J_v - I_v) + kP \otimes I_v
\end{aligned}$$

so that  $A_3 A_4 = A_4 A_3 = kA_1 + \frac{\lambda}{2}A_2$ . Next,  $(P \otimes I_v)(I_2 \otimes W_0 + P \otimes W_1) = I_2 \otimes W_1 + P \otimes W_0$ , hence  $A_1 A_3 = A_3 A_1 = A_4$ ; and similary,  $A_1 A_4 = A_4 A_1 = A_3$ . Then  $(P \otimes I_v)[J_2 \otimes (J_v - W_0 - W_1)] = J_2 \otimes (J_v - W_0 - W_1)$  and  $A_1 A_5 = A_5 A_1 = A_5$ . Now,

$$\begin{aligned}
(I_2 \otimes W_0 + P \otimes W_1)[J_2 \otimes (J_v - W_0^t - W_1^t)] &= J_2 \otimes [(W_0 + W_1)(J_v - (W_0 + W_1)^t)] \\
&= (k - \lambda)J_2 \otimes (J_v - I_v)
\end{aligned}$$



### 3. A New Family of Balanced Weighing Matrices and Association Schemes

so that  $A_3A_5 = A_5A_3 = (k - \lambda)A_2$ . Similarly,  $A_4A_5 = A_5A_4 = (k - \lambda)A_2$ . Since

$$\begin{aligned} [J_2 \otimes (J_v - I_v)](I_2 \otimes W_0 + P \otimes W_1) &= J_2 \otimes (J_v - I_v)(W_0 + W_1) \\ &= (k - 1)J_2 \otimes J_v + J_2 \otimes (J_v - W_0 - W_1), \end{aligned}$$

it follows that  $A_2A_3 = A_3A_2 = (k - 1)(A_3 + A_4 + A_5) + kA_5$ . Similarly,  $A_2A_4 = A_4A_2 = (k - 1)(A_3 + A_4 + A_5) + kA_5$ . Finally,

$$[J_2 \otimes (J_v - I_v)][J_2 \otimes (J_v - W_0 - W_1)] = J_2 \otimes [2(v - k)J_v - 2(J_v - W_0 - W_1)],$$

and  $A_2A_5 = A_5A_2 = 2(v - k)(A_3 + A_4 + A_5) - 2A_5$ .

We have shown the following result.

**10.1. Theorem.** If there is a balanced  $W(v, k)$ , then there is a 5-class symmetric association scheme.

\* \* \*

**10.2. Character Tables.** Our work from the previous subsection shows that the third intersection matrix is given by

$$B_3 = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & k-1 & k-1 & k \\ k & 0 & \frac{k(k-1)}{2(v-1)} & 0 & 0 & 0 \\ 0 & k & \frac{k(k-1)}{2(v-1)} & 0 & 0 & 0 \\ 0 & 0 & \frac{k(v-k)}{v-1} & 0 & 0 & 0 \end{pmatrix}.$$

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

It can be shown that  $B_3^t$  has the six distinct eigenvalues  $\pm k, \pm\sqrt{k}$ , and  $\pm\sqrt{k(v-k)/(v-1)}$  with corresponding eigenvectors

$$\mathbf{1}_6, \begin{pmatrix} 1 \\ 1 \\ 1 \\ -1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 0 \\ \frac{1}{\sqrt{k}} \\ -\frac{1}{\sqrt{k}} \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 0 \\ -\frac{1}{\sqrt{k}} \\ \frac{1}{\sqrt{k}} \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ \frac{-1}{v-1} \\ \frac{v-1}{v-k} \\ \frac{\sqrt{k(v-1)(v-k)}}{\sqrt{k(v-1)(v-k)}} \\ \frac{\sqrt{v-k}}{\sqrt{k(v-1)}} \\ \frac{-k}{\sqrt{k(v-1)(v-k)}} \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ \frac{-1}{k-v} \\ \frac{v-1}{k-v} \\ \frac{\sqrt{k(v-1)(v-k)}}{\sqrt{k(v-1)(v-k)}} \\ -\sqrt{\frac{v-k}{k(v-1)}} \\ \frac{k}{\sqrt{k(v-1)(v-k)}} \end{pmatrix}.$$

The valencies of the scheme are  $k_0 = k_1 = 1$ ,  $k_2 = 2(v-1)$ ,  $k_3 = k_4 = k$ , and  $k_5 = 2(v-k)$ . Define  $\Delta_k = \text{diag}(k_0, \dots, k_5)$ . Then  $v_i^t = (\Delta_k u_i)^t$  are the standardized left eigenvectors of  $B_3^t$ . The vectors  $v_i^t$  form the rows of the first character table  $P$ .

Next, the multiplicities of the scheme are given by  $m_i = 4v/\langle u_i, v_i \rangle$  and evaluate to 1, 1,  $v$ ,  $v$ ,  $v-1$ , and  $v-1$ . Then  $m_i u_i$  are the columns of the second character table  $Q$ .

Summing up, we have the following result.

**10.2. Theorem.** The 5-class symmetric association scheme of Theorem 10.1 has the character tables

$$P = \begin{matrix} & A_0 & A_1 & A_2 & A_3 & A_4 & A_5 \\ \begin{matrix} E_0 \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \end{matrix} & \begin{pmatrix} 1 & 1 & 2(v-1) & k & k & 2(v-k) \\ 1 & -1 & 0 & \sqrt{k} & -\sqrt{k} & 0 \\ 1 & -1 & 0 & -\sqrt{k} & \sqrt{k} & 0 \\ 1 & 1 & 2(v-1) & -k & -k & 2(k-v) \\ 1 & 1 & -2 & -\sqrt{\frac{k(v-k)}{v-1}} & -\sqrt{\frac{k(v-k)}{v-1}} & 2\sqrt{\frac{k(v-k)}{v-1}} \\ 1 & 1 & -2 & \sqrt{\frac{k(v-k)}{v-1}} & \sqrt{\frac{k(v-k)}{v-1}} & -2\sqrt{\frac{k(v-k)}{v-1}} \end{pmatrix} \end{matrix}$$

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

$$Q = \begin{matrix} & E_0 & E_1 & E_2 & E_3 & E_4 & E_5 \\ \begin{matrix} A_0 \\ A_1 \\ A_2 \\ A_3 \\ A_4 \\ A_5 \end{matrix} & \begin{pmatrix} 1 & v & v & 1 & v-1 & v-1 \\ 1 & -v & -v & 1 & v-1 & v-1 \\ 1 & 0 & 0 & 1 & -1 & -1 \\ 1 & \frac{v}{\sqrt{k}} & -\frac{v}{\sqrt{k}} & -1 & -\sqrt{\frac{(v-1)(v-k)}{k}} & \sqrt{\frac{(v-1)(v-k)}{k}} \\ 1 & -\frac{v}{\sqrt{k}} & \frac{v}{\sqrt{k}} & -1 & -\sqrt{\frac{(v-1)(v-k)}{k}} & \sqrt{\frac{(v-1)(v-k)}{k}} \\ 1 & 0 & 0 & -1 & \sqrt{\frac{k(v-1)}{v-k}} & -\sqrt{\frac{k(v-1)}{v-k}} \end{pmatrix} \end{pmatrix}.$$

Interestingly, the converse holds as well.

**10.3. Theorem.** If there is a 5-class symmetric association scheme with the character tables given in the statement of Theorem 10.2, then there is a balanced  $W(v, k)$ .

**Proof.** Let  $\tilde{A}_0, \dots, \tilde{A}_5$  be the adjacency matrices of the scheme. Then  $\tilde{A}_1 + \tilde{A}_2$  has eigenvalues  $2v - 1$  and  $-1$  with multiplicities 2 and  $4v - 2$ , respectively. It follows (see note 21) that  $\tilde{A}_1 + \tilde{A}_2 \sim I_2 \otimes A(K_{2v})$ . By the eigenvalues of  $\tilde{A}_1$ , it is the adjacency matrix of  $2v$  disjoint 2-cliques, hence  $\tilde{A}_1 \sim I_2 \otimes P \otimes I_v$  and  $\tilde{A}_2 \sim I_2 \otimes J_2 \otimes (J_v - I_v)$ .

It follows that

$$A_3 = \begin{pmatrix} O & O & X_1 & X_2 \\ O & O & X_3 & X_4 \\ X_1^t & X_3^t & O & O \\ X_2^t & X_4^t & O & O \end{pmatrix} \text{ and } A_4 = \begin{pmatrix} O & O & Y_1 & Y_2 \\ O & O & Y_3 & Y_4 \\ Y_1^t & Y_3^t & O & O \\ Y_2^t & Y_4^t & O & O \end{pmatrix}.$$

It can be shown (see Bannai and Ito, 1984, Theorem 3.6.2, for example) that the eigenvalues of the scheme imply that  $A_1 A_3 = A_4$  so that

$$\begin{pmatrix} O & O & X_3 & X_4 \\ O & O & X_1 & X_2 \\ X_2^t & X_4^t & O & O \\ X_1^t & X_3^t & O & O \end{pmatrix} = \begin{pmatrix} O & O & Y_1 & Y_2 \\ O & O & Y_3 & Y_4 \\ Y_1^t & Y_3^t & O & O \\ Y_2^t & Y_4^t & O & O \end{pmatrix}.$$

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

Therefore, there are  $(0, 1)$ -matrices  $W_0$  and  $W_1$  such that

$$A_3 = \begin{pmatrix} O & O & W_0 & W_1 \\ O & O & W_1 & W_0 \\ W_0^t & W_1^t & O & O \\ W_1^t & W_0^t & O & O \end{pmatrix} \text{ and } A_4 = \begin{pmatrix} O & O & W_1 & W_0 \\ O & O & W_0 & W_1 \\ W_1^t & W_0^t & O & O \\ W_0^t & W_1^t & O & O \end{pmatrix}.$$

Appealing to the character tables again, it can be shown that

$$\begin{aligned} A_3^2 &= kA_0 + \frac{k(k-1)}{2(v-1)}A_2, \\ A_3A_4 &= A_4A_3 = kA_1 + \frac{k(k-1)}{2(v-1)}A_2, \text{ and} \\ A_4^2 &= kA_0 + \frac{k(k-1)}{2(v-1)}A_2. \end{aligned}$$

Therefore,  $(A_3 - A_4)^2 = 2k(A_0 - A_1)$  and  $(A_3 + A_4)^2 = 2k(A_0 + A_1) + \frac{2k(k-1)}{v-1}A_2$ , from which it follows that

$$\begin{aligned} (W_0 - W_1)(W_0 - W_1)^t &= kI, \text{ and} \\ (W_0 + W_1)(W_0 + W_1)^t &= kI + \lambda(J - I), \end{aligned}$$

where  $\lambda = k(k-1)/(v-1)$ .

We then have that  $W_0 - W_1$  is the required matrix. ■

In light of the equivalence, we make the following definition.

**10.4. Definition.** A symmetric 5-class association scheme is a *weighing scheme* if it has the character tables given in Theorem 10.2.

## Notes

18. Seberry (2017), Seberry and Yamada (1992), and Wallis et al. (1972) all contain a wealth of constructions for weighing matrices and orthogonal designs using the Kronecker product.

### 3. A New Family of Balanced Weighing Matrices and Association Schemes

19. Such a decomposition of the block set is called a *parallelism*. Beth et al. (1999) studies partitions of the blocks of an incidence structure in great detail.

# 4

## A Unified Construction of Weighing Matrices

In this final chapter of results, we present a general method to construct certain configurations using classical parameter BGW matrices and the generalized simplex codes. This method is similar to that given by Rajkundlia (1983) and Ionin (2001), but it can be seen that it is much simpler to apply than the recursive methods of the aforementioned seminal articles. Following the presentation of this construction technique, we find an equivalence between arbitrary BGW matrices over a finite abelian group and certain commutative association schemes. This result generalizes that obtained over the course of the previous chapter.

### §11. A General Method

In this section, the general method intimated above is derived and used in the construction of weighing matrices and symmetric designs. Resulting parameters of configurations so constructible are then tabulated.

#### 4. A Unified Construction of Weighing Matrices

\* \* \*

**11.1. A First Application: Weighing matrices.** It is a peculiarity of the construction that the application of the method to weighing matrices is predicated upon the weight of the matrix being a prime power. Indeed, as in the previous chapter, the generalized simplex codes are applied to the derived part of the matrix by substituting the rows of the derived part for the letters of the code. The simplex code, as the reader may remember, has a prime power number of letters.

To begin, let  $W = \begin{pmatrix} 0 & R \\ 1 & D \end{pmatrix}$  be a  $W(v, q)$  weighing matrix in normal form, where  $q$  is some prime power. Then it is easy to see that  $RR^t = kI$  and  $DD^t = kI - J$ . We say that  $W$  is the *seed matrix* of the construction.

Let  $\mathcal{S}_{q,n} = \sum_{\alpha \in \text{GF}(q)^*} \alpha A_\alpha$ , and define  $A_0 = J - \sum_{\alpha \in \text{GF}(q)^*} A_\alpha$ . Index the rows of  $D$  by the elements of  $\text{GF}(q)$ , and take  $\mathcal{D} = \sum_{\alpha \in \text{GF}(q)} A_\alpha \otimes r_\alpha$ . Next, by our previous work, there is a  $W((q^n - 1)/(q - 1), q^{n-1})$ , say  $H$ , for every  $n > 1$ . We define  $\mathcal{R} = H \otimes R$ .

We claim that  $\mathcal{W} = \begin{pmatrix} 0 & \mathcal{R} \\ 1 & \mathcal{D} \end{pmatrix}$  is again a weighing matrix. Indeed,  $\mathcal{R}\mathcal{R}^t = (H \otimes R)(H \otimes R)^t = HH^t \otimes RR^t = q^n I$ . Further, by Lemma 9.1,

$$\begin{aligned} \mathcal{D}\mathcal{D}^t &= \left( \sum_{\alpha} A_\alpha \otimes r_\alpha \right) \left( \sum_{\alpha} A_\alpha \otimes r_\alpha \right) \\ &= \sum_{\alpha} A_\alpha A_\alpha^t \otimes r_\alpha r_\alpha^t + \sum_{\alpha \neq \beta} A_\alpha A_\beta^t \otimes r_\alpha r_\beta^t \\ &= (q - 1) \sum_{\alpha} A_\alpha A_\alpha^t - \sum_{\alpha \neq \beta} A_\alpha A_\beta^t \\ &= (q - 1) \left( \frac{q^{n-1} - 1}{q - 1} J + q^{n-1} I \right) - q^{n-1} (J - I) \\ &= q^n I - J. \end{aligned}$$

#### 4. A Unified Construction of Weighing Matrices

Finally,

$$\begin{aligned}
 \mathcal{RD}^t &= (H \otimes R) \left( \sum_{\alpha} A_{\alpha} \otimes r_{\alpha} \right)^t \\
 &= \sum_{\alpha} H A_{\alpha}^t \otimes R r_{\alpha}^t \\
 &= \sum_{\alpha} H A_{\alpha}^t \otimes O \\
 &= O.
 \end{aligned}$$

It follows that  $\mathcal{W}$  is a weighing matrix. We record this result below.

**11.1. Theorem.** If there exists a  $W(v, q)$  weighing matrix of prime power weight, then there is a weighing matrix with parameters

$$(11.1.a) \quad \left( \frac{(v-1)(q^n-1)}{q-1} + 1, q^n \right).$$

**11.2. Example.** Take as a seed matrix the  $W(8, 5)$  shown below

$$(11.2.a) \quad \begin{pmatrix} 0 & + & 0 & 0 & - & - & + & + \\ 0 & 0 & + & 0 & - & + & - & + \\ 0 & 0 & 0 & + & - & + & + & - \\ + & + & + & + & + & 0 & 0 & 0 \\ + & + & - & - & 0 & + & 0 & 0 \\ + & - & + & - & 0 & 0 & + & 0 \\ + & - & - & + & 0 & 0 & 0 & + \\ + & 0 & 0 & 0 & - & - & - & - \end{pmatrix}.$$

Applying the construction, we find a  $W(43, 25)$



#### 4. A Unified Construction of Weighing Matrices

(11.2.b)

Evidently, this resolves the question of existence of a  $W(43, 25)$  given in Part V of Colbourn and Dinitz (2007).

To further evince the utility of the method, we include a table of constructible parameters given a seed weighing matrix known to exist.

Table 4.1: Small parameter consequential order/weight pairs.

$Seed(v, k)$	$Succident(v', k')$	$Seed(v, k)$	$Succident(v', k')$
$(6, 5)^\ddagger$ :	$(31, 25)^\ddagger, (156, 125)^\ddagger, (781, 625)^\ddagger$	$(16, 3)$ :	$(69, 9), (196, 27), (601, 81)$
$(8, 5)$ :	$(43, 25)^\dagger, (218, 125)$	$(16, 5)$ :	$(91, 25), (466, 125)$
$(8, 7)^\ddagger$ :	$(57, 49)^\ddagger, (400, 343)^\ddagger$	$(16, 7)$ :	$(121, 49), (856, 343)$
$(10, 5)$ :	$(55, 25), (280, 125)$	$(16, 9)$ :	$(151, 81)$
$(10, 9)^\ddagger$ :	$(91, 81)^\ddagger, (820, 729)^\ddagger$	$(16, 11)$ :	$(181, 121)^\dagger$
$(12, 5)$ :	$(67, 25), (342, 125)$	$(16, 13)$ :	$(211, 169)$
$(12, 7)$ :	$(89, 49)^\dagger, (628, 343)$	$(18, 13)$ :	$(239, 169)$

#### 4. A Unified Construction of Weighing Matrices

Continuation of Table 4.1.

<i>Seed</i> ( $v, k$ )	<i>Succident</i> ( $v', k'$ )	<i>Seed</i> ( $v, k$ )	<i>Succident</i> ( $v', k'$ )
(12, 9):	(111, 81) <sup>†</sup>	(19, 9) <sup>‡</sup> :	(181, 81) <sup>*</sup>
(13, 9) <sup>‡</sup> :	(121, 81)	(20, 7):	(153, 49)
(14, 9):	(131, 81)	(20, 13):	(267, 169) <sup>†</sup>
(14, 13) <sup>‡</sup> :	(183, 169) <sup>‡</sup>		

\* Note that the  $W(181, 81)$  constructible from the balanced seed  $W(19, 9)$  can be made to be balanced as shown in the previous chapter. It is not, however, a consequence of the construction of this chapter that the succident matrix is balanced.

<sup>†</sup> Denotes previously unknown order weight pairs.

<sup>‡</sup> Denotes a balanced weighing matrix.

\* \* \*

**11.2. A Second Application: Block designs.** The construction is perfectly amenable to certain symmetric designs. For precisely the same reasoning given in the previous subsection, we require that the design have a prime power block size. Furthermore, we require that the parameters of the residual of the design be invariant under complementation. Explicitly, if  $A = \begin{pmatrix} 0 & R \\ 1 & D \end{pmatrix}$  is the incidence matrix of the given design, then  $J - R$  must have the same parameters as  $R$ .

As shown in Hall (1986), there is a  $W(2q + 2, 2q + 2)$  if and only if there is a symmetric  $BIBD(2q + 1, q, (q - 1)/2)$ , called a *Hadamard design*<sup>20</sup>. Accordingly, the residual of this design is a  $BIBD(q + 1, 2q, q, (q + 1)/2, (q - 1)/2)$ , the parameters of which are invariant under complementation. Also, the derived design is a  $BIBD(q, 2q, q - 1, (q - 1)/2, (q - 3)/2)$ .

Let  $H = H_0 - H_1$  be a balanced  $W((q^n - 1)/(q - 1), q^{n-1})$ , and let  $\{A_\alpha\}_{\alpha \in GF(q)}$  be as above. If  $A = \begin{pmatrix} 0 & R \\ 1 & D \end{pmatrix}$  is a Hadamard design with parameters  $(2q + 1, q, (q - 1)/2)$ , then we form  $\mathcal{R} = H_0 \otimes R + H_1 \otimes (J - R)$ . Indexing the rows of  $D$  by

#### 4. A Unified Construction of Weighing Matrices

the elements of  $\text{GF}(q)$ , we then form  $\mathcal{D} = \sum_{\alpha \in \text{GF}(q)} A_\alpha \otimes r_\alpha$ .

We claim that  $\mathcal{A} = \begin{pmatrix} 0 & \mathcal{R} \\ 1 & \mathcal{D} \end{pmatrix}$  is a symmetric design. To show this, we proceed as before. Using Lemma 9.2, we find

$$\begin{aligned}
\mathcal{R}\mathcal{R}^t &= [H_0 \otimes R + H_1 \otimes (J_{q+1,2q} - R)][H_0 \otimes R + H_1 \otimes (J_{q+1,2q} - R)]^t \\
&= (H_0 H_0^t + H_1 H_1^t) \otimes \left( \frac{q+1}{2} I_{q+1} + \frac{q-1}{2} J_{q+1} \right) \\
&\quad + \frac{q+1}{2} (H_0 H_1^t + H_1 H_0^t) \otimes (J_{q+1} - I_{q+1}) \\
&= \frac{q^{n-2}}{4} [(q+1) I_{\frac{q^n-1}{q-1}} + (q-1) J_{\frac{q^n-1}{q-1}}] \otimes [(q+1) I_{q+1} + (q-1) J_{q+1}] \\
&\quad + \frac{q^{n-2}(q^2-1)}{4} (J_{\frac{q^n-1}{q-1}} - I_{\frac{q^n-1}{q-1}}) \otimes (J_{q+1} - I_{q+1}) \\
&= \frac{q^{n-2}}{4} \left[ 2q(q+1) I_{\frac{(q+1)(q^n-1)}{q-1}} + 2q(q-1) J_{\frac{(q+1)(q^n-1)}{q-1}} \right] \\
&= q^n I_{\frac{(q+1)(q^n-1)}{q-1}} + \frac{q^n - q^{n-1}}{2} (J_{\frac{(q+1)(q^n-1)}{q-1}} - I_{\frac{(q+1)(q^n-1)}{q-1}}),
\end{aligned}$$

whence  $\mathcal{R}$  is a quasi-residual BIBD with parameters

$$\left( \frac{(q+1)(q^n-1)}{q-1}, \frac{2q^{n+1}-2q}{q-1}, q^n, \frac{q^n+q^{n-1}}{2}, \frac{q^n-q^{n-1}}{2} \right).$$

Next, by Lemma 9.1

$$\begin{aligned}
\mathcal{D}\mathcal{D}^t &= \left( \sum_{\alpha} A_\alpha \otimes r_\alpha \right) \left( \sum_{\alpha} A_\alpha \otimes r_\alpha \right)^t \\
&= \sum_{\alpha} A_\alpha A_\alpha^t \otimes r_\alpha r_\alpha^t + \sum_{\alpha \neq \beta} A_\alpha A_\beta^t \otimes r_\alpha r_\beta^t \\
&= (q-1) \sum_{\alpha} A_\alpha A_\alpha^t + \frac{q-3}{2} \sum_{\alpha \neq \beta} A_\alpha A_\beta^t \\
&= (q-1) \left( \frac{q^{n-1}-1}{q-1} J - q^{n-1} I \right) + \frac{q^{n-1}(q-3)}{2} (J - I) \\
&= (q^n-1) I + \frac{q^n - q^{n-1} - 2}{2} (J - I),
\end{aligned}$$

#### 4. A Unified Construction of Weighing Matrices

and  $\mathcal{D}$  is a quasi-derived BIBD with parameters

$$\left( q^{n+1}, \frac{2q^{n+1} - 2q}{q-1}, q^n - 1, \frac{q^n - q^{n-1}}{2}, \frac{q^n - q^{n-1} - 2}{2} \right).$$

Finally,

$$\begin{aligned} \mathcal{RD}^t &= [H_0 \otimes R + H_1 \otimes (J - R)] \left( \sum_{\alpha} A_{\alpha} \otimes r_{\alpha} \right)^t \\ &= \sum_{\alpha} H_0 A_{\alpha}^t \otimes R r_{\alpha}^t + \sum_{\alpha} H_1 A_{\alpha}^t \otimes (J - R) r_{\alpha}^t \\ &= \frac{q-1}{2} \sum_{\alpha} H_0 A_{\alpha}^t \otimes \mathbf{1} + \frac{q-1}{2} \sum_{\alpha} H_1 A_{\alpha}^t \otimes \mathbf{1} \\ &= \frac{q-1}{2} (H_0 + H_1) \sum_{\alpha} A_{\alpha}^t \otimes \mathbf{1} \\ &= \frac{q-1}{2} (H_0 + H_1) J \\ &= \frac{q^n - q^{n-1}}{2} J. \end{aligned}$$

We have shown that  $\mathcal{A} = \begin{pmatrix} \mathcal{R} \\ \mathcal{D} \end{pmatrix}$  is a symmetric BIBD. We record this result below.

**11.3. Theorem.** If there is a symmetric BIBD( $2q+1, q, (q-1)/2$ ), then there is a symmetric BIBD with parameters

$$(11.3.a) \quad \left( \frac{2q^{n+1} - 2q}{q-1} + 1, q^n, \frac{q^n - q^{n-1}}{2} \right).$$

**11.4. Example.** Consider the Hadamard design

$$(11.4.a) \quad \begin{pmatrix} 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

with parameters  $(7, 3, 1)$ . The result of the first iteration of the construction is the symmetric BIBD( $25, 9, 3$ )

#### 4. A Unified Construction of Weighing Matrices

$$(11.4.b) \quad \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix}.$$

## §12. BGW Matrices and Association Schemes

In this section we present perhaps the most striking result of this essay. Here the equivalence between BGW matrices and commutative association schemes is described in detail.

\* \* \*

**12.1. Adjacency Matrices.** We must first extend Lemma 9.2 to the case of arbitrary BGW matrices. Let  $G = \{g_0 = 1, g_1, \dots, g_{n-1}\}$  be an abelian group, and let  $W = \sum_{g \in G} gW_g$  be a  $\text{BGW}(v, k, \lambda; G)$ . We then have the following.

**12.1. Lemma.**

$$(12.1.a) \quad \sum_{g,h} gh^{-1}W_gW_h^t = \sum_{g,h} h^{-1}gW_h^tW_g = kI + \frac{\lambda}{n} \left( \sum_g g \right) (J - I),$$

#### 4. A Unified Construction of Weighing Matrices

$$(12.1.b) \quad \sum_g W_g W_g^t = \sum_g W_g^t W_g = kI + \frac{\lambda}{n}(J - I), \text{ and}$$

$$(12.1.c) \quad \sum_g W_g W_{gh^{-1}}^t = \sum_g W_{gh^{-1}}^t W_g = \frac{\lambda}{n}(J - I) \text{ whenever } h \neq 1.$$

**Proof.** Restatement of the fact that  $W = \sum_g g W_g$  is a BGW( $v, k, \lambda; G$ ). ■

Let  $\{U_g : g \in G\}$  be the usual regular linear representation of  $G$ , that is,  $U_g = (\delta(g_i^{-1} g g_j))$  where  $\delta(h) = 1$  if  $h = 1$  and 0 otherwise. Consider the following family of matrices.

$$\begin{aligned} A_{0,g} &= I_2 \otimes U_g \otimes I_v, \text{ for } g \in G, \\ A_1 &= I_2 \otimes J_n \otimes (J_v - I_v), \\ A_{2,g} &= \begin{pmatrix} O & \sum_{h \in G} (U_h \otimes W_{gh}) \\ \sum_{h \in G} (U_h \otimes W_{g^{-1}h^{-1}}^t) & O \end{pmatrix}, \text{ for } g \in G, \text{ and} \\ A_3 &= \begin{pmatrix} O & J_n \otimes (J_v - \sum_{h \in G} W_h) \\ J_n \otimes (J_v - \sum_{h \in G} W_h^t) & O \end{pmatrix}. \end{aligned}$$

In showing that  $\{A_1, A_3\} \cup \{A_{0,g}, A_{2,g} : g \in G\}$  is an association scheme, we proceed precisely as before. First,  $A_1 + A_3 + \sum_{g \in G} (A_{0,g} + A_{2,g}) = J$ . Second,  $A_{0,g}^t = A_{0,g^{-1}}, A_1^t = A_1, A_{2,g}^t = A_{2,g^{-1}}$ , and  $A_3^t = A_3$ .

We next show closure under multiplication. We have  $A_{0,g} A_{0,h} = A_{0,gh} = A_{0,hg} = A_{0,h} A_{0,g}$ . Then  $A_{0,g} A_1 = A_1 A_{0,g} = A_1$  and  $A_{0,g} A_3 = A_3 A_{0,g} = A_3$ . Since  $\sum_j (U_{gj} \otimes W_{hj}) = \sum_j (U_j \otimes W_{g^{-1}hj})$ , it follows that  $A_{0,g} A_{2,h} = A_{2,h} A_{0,g} = A_{2,g^{-1}h}$ . As  $[J_n \otimes (J_v - I_v)]^2 = n(v-1) \sum_g (U_g \otimes I_v) + n(v-$

#### 4. A Unified Construction of Weighing Matrices

2)  $J_n \otimes (J_v - I_v)$ , we have  $A_1^2 = n(v-1) \sum_g A_{0,g} + n(v-2)A_1$ . Next,

$$\begin{aligned} [J_n \otimes (J_v - I_v)] \sum_h (U_h \otimes W_{gh}) &= J_n \otimes (J_v - I_v) \sum_h W_h \\ &= (k-1)J_n \otimes J_v + J_n \otimes (J_v - \sum_h W_h), \end{aligned}$$

hence  $A_1 A_{2,g} = A_{2,g} A_1 = (k-1) \sum_{g \in G} A_{2,g} + kA_3$ . We have

$$\begin{aligned} [J_n \otimes (J_v - I_v)] \left[ J_n \otimes (J_v - \sum_h W_h) \right] &= nJ_n \otimes (J_v - I_v) \left( J_v - \sum_h W_h \right) \\ &= n(v-k)J_n \otimes J_v - nJ_n \otimes \left( J_v - \sum_h W_h \right) \end{aligned}$$

so that  $A_1 A_3 = A_3 A_1 = n(v-k) \sum_{g \in G} A_{2,g} + n(v-k-1)A_3$ . Then

$$\begin{aligned} \sum_h (U_h \otimes W_{gh}) \left[ J_n \otimes (J_v - \sum_j W_j^t) \right] &= J_n \otimes \left( \sum_h W_h J_v - \sum_{h,j} W_h W_j^t \right) \\ &= (k-\lambda)J_n \otimes (J_v - I_v), \end{aligned}$$

and  $A_{2,g} A_3 = A_3 A_{2,g} = (k-\lambda)A_1$ . Since

$$\begin{aligned} \left( J_v - \sum_{h \in G} W_h \right) \left( J_v - \sum_{g \in G} W_g^t \right) &= vJ_v - 2kJ_v + kI_v + \lambda(J_v - I_v) \\ &= (v-2k+\lambda)(J_v - I_v) + (v-k)I_v, \end{aligned}$$

one finds that  $A_3^2 = n(v-2k+\lambda)A_1 + n(v-k) \sum_{g \in G} A_{0,g}$ . Finally, for fixed  $g, h \in G$ , we note that

$$\sum_{\alpha, \beta} (U_{\alpha\beta} \otimes W_{g\alpha} W_{h^{-1}\beta^{-1}}^t) = \sum_{\gamma} \left( U_{\gamma} \otimes \sum_{\alpha\beta=\gamma} W_{g\alpha} W_{h^{-1}\beta^{-1}}^t \right)$$

#### 4. A Unified Construction of Weighing Matrices

$$\begin{aligned}
&= \sum_{\gamma \neq g^{-1}h^{-1}} \left( U_\gamma \otimes \sum_{\alpha\beta=\gamma} W_{g\alpha} W_{h^{-1}\beta^{-1}}^t \right) \\
&\quad + U_{g^{-1}h^{-1}} \otimes \sum_{\gamma} W_\gamma W_\gamma^t \\
&= \sum_{\gamma \neq g^{-1}h^{-1}} \left[ U_\gamma \otimes \frac{\lambda}{n} (J_v - I_v) \right] \\
&\quad + U_{g^{-1}h^{-1}} \otimes \left[ kI_v + \frac{\lambda}{n} (J_v - I_v) \right] \\
&= \frac{\lambda}{n} J_n \otimes (J_v - I_v) + kU_{g^{-1}h^{-1}} \otimes I_v
\end{aligned}$$

so that  $A_{2,g}A_{2,h} = A_{2,h}A_{2,g} = \frac{\lambda}{n}A_1 + kA_{0,g^{-1}h^{-1}}$ .

We have shown the following result.

**12.2. Theorem.** If  $G$  is a finite abelian group, and if there is a BGW( $v, k, \lambda; G$ ), then there is either a commutative  $2n$ - or  $(2n + 1)$ -class association scheme predicated upon whether or not  $v = k$ .

\* \* \*

**12.2. Character Tables.** As before, we desire to give explicitly the character tables of the scheme. The difference between the schemes arising from an arbitrary BGW vs. a balanced weighing matrix, is that the number of classes is no longer fixed. Therefore, we desire to exhibit a general form for the primitive idempotents of the scheme in terms of the adjacency matrices.

To this end, let  $\hat{G} = \{\chi_g : g \in G\}$  be the collection of irreducible characters<sup>21)</sup> of the group  $G$ , i.e. the dual group of  $G$ . Following Kharaghani and Suda (2021), we make the following definitions for each  $g \in G$ :

$$F_{0,g} = \sum_{h \in G} \chi_g(h) A_{0,h}, \quad \text{and} \quad F_{2,g} = \sum_{h \in G} \chi_g(h) A_{2,h}.$$



#### 4. A Unified Construction of Weighing Matrices

Using the intersection numbers derived above, and using the generalized orthogonality relation of characters (see Isaacs, 2006, Theorem 2.13), we have the following lemma.

##### 12.3. Lemma.

$$(12.3.a) \quad F_{0,g}F_{0,h} = n\delta_{gh}F_{0,g},$$

$$(12.3.b) \quad F_{0,g}F_{2,h} = F_{2,h}F_{0,g} = n\delta_{g^{-1}h}F_{2,h}, \text{ and}$$

$$(12.3.c) \quad F_{2,g}F_{2,h} = n\lambda\delta_{gh}\delta_{g1}A_1 + nk\delta_{gh}F_{0,g^{-1}}.$$

**Proof.** We have

$$\begin{aligned} F_{0,g}F_{0,h} &= \sum_{\alpha,\beta} \chi_g(\alpha)\chi_h(\beta)A_{0,\alpha\beta} \\ &= n \sum_{\gamma} \left( \sum_{\alpha\beta=\gamma} \chi_g(\alpha)\chi_h(\beta) \right) A_{0,\gamma} \\ &= n \sum_{\gamma} \left( \sum_{\beta} \chi_g(\beta^{-1}\gamma)\chi_h(\beta) \right) A_{0,\gamma} \\ &= n\delta_{gh} \sum_{\gamma} \chi_g(\gamma)A_{0,\gamma} \\ &= n\delta_{gh}F_{0,g}, \end{aligned}$$

which shows (12.3.a). Next,

$$\begin{aligned} F_{0,g}F_{2,h} &= \sum_{\alpha,\beta} \chi_g(\alpha)\chi_h(\beta)A_{0,g}A_{2,h} \\ &= n \sum_{\gamma} \left( \sum_{\alpha\beta=\gamma} \chi_{g^{-1}}(\alpha)\chi_h(\beta) \right) A_{2,\gamma} \\ &= n\delta_{g^{-1}h} \sum_{\gamma} \chi_h(\gamma)A_{2,\gamma} \end{aligned}$$

#### 4. A Unified Construction of Weighing Matrices

$$= n\delta_{g^{-1}h}F_{2,h}.$$

Since the scheme is commutative,  $F_{2,h}F_{0,g} = F_{0,g}F_{2,h}$ , which shows (12.3.b).

Finally,

$$\begin{aligned} F_{2,g}F_{2,h} &= \sum_{\alpha,\beta} \chi_g(\alpha)\chi_h(\beta) \left( \frac{\lambda}{n}A_1 + kA_{0,\alpha^{-1}\beta^{-1}} \right) \\ &= n \sum_{\gamma} \left( \sum_{\alpha\beta=\gamma} \chi_{g^{-1}}(\alpha)\chi_{h^{-1}}(\beta) \right) \left( \frac{\lambda}{n}A_1 + kA_{0,\gamma} \right) \\ &= n \sum_{\gamma} \left( \sum_{\beta} \chi_{g^{-1}}(\beta^{-1}\gamma)\chi_{h^{-1}}(\beta) \right) \left( \frac{\lambda}{n}A_1 + kA_{0,\gamma} \right) \\ &= n\delta_{gh} \sum_{\gamma} \chi_{g^{-1}}(\gamma) \left( \frac{\lambda}{n}A_1 + kA_{0,\gamma} \right) \\ &= n\lambda\delta_{gh}\delta_{g1}A_1 + nk\delta_{gh}F_{0,g^{-1}}, \end{aligned}$$

showing (12.3.c). This completes the proof.  $\blacksquare$

We are now ready to give the idempotents of the scheme. There are two cases to consider, namely, whether or not  $v = k$ . In the latter case, we have

$$\begin{aligned} E_0 &= \frac{1}{2nv}(F_{0,0} + F_{2,0} + A_1 + A_3), \\ E_1 &= \frac{1}{2nv}(F_{0,0} - F_{2,0} + A_1 - A_3), \\ E_{2,1} &= \frac{1}{2nv} \left( (v-1)F_{0,0} + \sqrt{\frac{(v-1)(v-k)}{k}}F_{2,0} - A_1 - \sqrt{\frac{k(v-1)}{v-k}}A_3 \right), \\ E_{2,2} &= \frac{1}{2nv} \left( (v-1)F_{0,0} - \sqrt{\frac{(v-1)(v-k)}{k}}F_{2,0} - A_1 + \sqrt{\frac{k(v-1)}{v-k}}A_3 \right), \\ E_{3,g} &= \frac{1}{2nv} \left( vF_{0,g} + \frac{v}{\sqrt{k}}F_{2,g} \right), \text{ for } g \in G/\{1\}, \text{ and} \\ E_{4,g} &= \frac{1}{2nv} \left( vF_{0,g} - \frac{v}{\sqrt{k}}F_{2,g} \right), \text{ for } g \in G/\{1\}. \end{aligned}$$

#### 4. A Unified Construction of Weighing Matrices

In the case that  $v = k$ , then  $A_3 = O$  and we replace  $E_{2,1}$  and  $E_{2,2}$  with  $E_2 = E_{2,1} + E_{2,2}$ . The following lemma is immediate.

**12.4. Lemma.** Let  $\mathcal{E} = \{E_0, E_1, E_{2,1}, E_{2,2}, E_{3,g}, E_{4,g} : g \in G/\{1\}\}$ . Then:

$$(12.4.a) \quad EF = \delta_{EF}E, \text{ for all } E, F \in \mathcal{E};$$

$$(12.4.b) \quad I = \sum_{E \in \mathcal{E}} E; \text{ and}$$

$$(12.4.c) \quad E^* \in \mathcal{E}, \text{ for each } E \in \mathcal{E}.$$

**Proof.** Straightforward but tedious calculation. ■

Using these lemmata, we have the following result.

**12.5. Theorem.** The commutative association scheme given in Theorem 12.2 has the first and second character tables

$$P = \begin{matrix} & A_{0,h} & A_1 & A_{2,h} \\ \begin{matrix} E_0 \\ E_1 \\ E_2 \\ E_{3,g} \\ E_{4,g} \end{matrix} & \begin{pmatrix} 1 & n(v-1) & k \\ 1 & n(v-1) & -k \\ -n & 0 & \\ \chi_{g^{-1}}(h) & 0 & \sqrt{k}\chi_{g^{-1}}(h) \\ \chi_{g^{-1}}(h) & 0 & \chi_{g^{-1}}(h) \end{pmatrix} \end{matrix},$$

$$Q = \begin{matrix} & E_0 & E_1 & E_2 & E_{3,g} & E_{4,g} \\ \begin{matrix} A_{0,h} \\ A_1 \\ A_{2,h} \end{matrix} & \begin{pmatrix} 1 & 1 & 2(v-1) & v\chi_g(h) & v\chi_g(h) \\ 1 & 1 & -2 & 0 & 0 \\ 1 & -1 & 0 & \frac{v}{\sqrt{k}}\chi_g(h) & -\frac{v}{\sqrt{k}}\chi_g(h) \end{pmatrix} \end{matrix}$$

#### 4. A Unified Construction of Weighing Matrices

in the case that  $v = k$  and

$$P = \begin{matrix} & A_{0,h} & A_1 & A_{2,h} & A_3 \\ \begin{matrix} E_0 \\ E_1 \\ E_{2,1} \\ E_{2,2} \\ E_{3,g} \\ E_{3,g} \end{matrix} & \begin{pmatrix} 1 & n(v-1) & k & n(v-k) \\ 1 & n(v-1) & -k & n(k-v) \\ 1 & -n & \sqrt{\frac{k(v-k)}{v-1}} & -n\sqrt{\frac{k(v-k)}{v-1}} \\ 1 & -n & -\sqrt{\frac{k(v-k)}{v-1}} & n\sqrt{\frac{k(v-k)}{v-1}} \\ -\chi_{g^{-1}}(h) & 0 & \sqrt{k}\chi_{g^{-1}}(h) & 0 \\ \chi_{g^{-1}}(h) & 0 & -\sqrt{k}\chi_{g^{-1}}(h) & 0 \end{pmatrix} \end{matrix},$$

$$Q = \begin{matrix} & E_0 & E_1 & E_{2,1} & E_{2,2} & E_{3,g} & E_{4,g} \\ \begin{matrix} A_{0,h} \\ A_1 \\ A_{2,h} \\ A_3 \end{matrix} & \begin{pmatrix} 1 & 1 & v-1 & v-1 & v\chi_g(h) & v\chi_g(h) \\ 1 & 1 & -1 & -1 & 0 & 0 \\ 1 & -1 & \sqrt{\frac{(v-1)(v-k)}{v-k}} & -\sqrt{\frac{(v-1)(v-k)}{v-k}} & \frac{v}{\sqrt{k}}\chi_g(h) & -\frac{v}{\sqrt{k}}\chi_g(h) \\ 1 & 1 & -\sqrt{\frac{k(v-1)}{v-k}} & \sqrt{\frac{k(v-1)}{v-k}} & 0 & 0 \end{pmatrix} \end{matrix}$$

in the case that  $v > k$ .

As before, the converse also holds.

**12.6. Theorem.** If there is a commutative scheme with the character tables given in Theorem 12.5, then there is a  $\text{BGW}(v, k, \lambda; G)$ , where  $G$  is an abelian group isomorphic to  $\{\chi_g\}$ .

**Proof.** The derivation of this fact is the same mutatis mutandis as that for Theorem 10.3 and is, therefore, omitted. ■

In light of this equivalence, we make the following final definition.

**12.7. Definition.** The commutative association schemes with character tables given in Theorem 12.5 are called *generalized weighing schemes*.

#### 4. A Unified Construction of Weighing Matrices

Since the largest—indeed, the most important—families of generalized matrices are those over a group which is either cyclic or elementary abelian, it would be beneficial to conclude with a brief discussion of the irreducible characters of these groups in an effort to make the preceeding results more concrete.

To this end, recall the irreducible characters of the cyclic group  $C_p \simeq \{1, g, \dots, g^{p-1}\}$  of order  $p$  are given by  $\chi_{g^i}(g^j) = e^{\frac{2\pi\sqrt{-1}ij}{p}}$ . This suffices for the case in which the group is cyclic.

Let  $p$  be a prime, and let  $C_p$  be as above. If  $q = p^n$ , then  $\text{EA}(q) \simeq \underbrace{C_p \otimes \dots \otimes C_p}_n = \{g^{m_0} \otimes \dots \otimes g^{m_{n-1}} : 0 \leq m_0, \dots, m_{n-1} < p\}$ . By Theorem 4.21 of Isaacs (2006), it follows that

$$\begin{aligned} \chi_{g^{m_0} \otimes \dots \otimes g^{m_{n-1}}}(g^{k_0} \otimes \dots \otimes g^{k_{n-1}}) &= \prod_i \chi_{g^{m_i}}(g^{k_i}) \\ &= \prod_i e^{\frac{2\pi\sqrt{-1}m_i k_i}{p}} \\ &= e^{\frac{2\pi\sqrt{-1}}{p} \sum_i m_i k_i}. \end{aligned}$$

This concludes our study of weighing matrices and related configurations.

### Notes

20. Let  $C$  be the matrix obtained upon deleting the first row and column of a normalized Hadamard matrix of order  $4n$ . Then  $(1/2)(J + C)$  is a symmetric BIBD( $4n - 1, 2n - 1, n - 1$ ). See Hall (1986) for a proof of this result and related discussions.
21. Recall that a linear representation of a group  $G$  is a homomorphism  $\varrho : G \rightarrow \text{GL}(V)$ , where  $V$  is some linear space. The character of the representation is then  $\chi(g) = \text{Tr}(\varrho(g))$ , for each  $g \in G$ . Isaacs (2006) is the classical reference for the character theory of finite groups.

# Appendix

### A1. A Balancedly Splittable $W(64, 64)$

Beginning with (5.7.a), a Hadamard matrix of order 4, we first construct the auxiliary matrices as shown.

$$c_0 = \begin{pmatrix} + & + & + & + \\ + & + & + & + \\ + & + & + & + \\ + & + & + & + \end{pmatrix}, c_1 = \begin{pmatrix} + & - & + & - \\ - & + & - & + \\ + & - & + & - \\ - & + & - & + \end{pmatrix}, c_2 = \begin{pmatrix} + & - & - & + \\ - & + & + & - \\ - & + & + & - \\ + & - & - & + \end{pmatrix}, c_3 = \begin{pmatrix} + & + & - & - \\ + & + & - & - \\ - & - & + & + \\ - & - & + & + \end{pmatrix}.$$

The block-circulant matrix  $A$  has first row  $(c_0, c_1, c_2, c_3, c_3, c_2, c_1)$  and is given by

[illegible]

Similarly, the block-circulant matrix  $B$  has first row  $(c_0, c_1, c_2, c_3, -c_3, -c_2, -c_1)$

## Appendix

and is given by

[illegible]

It remains to construct the block matrix  $F$ . Label the 8 rows of the Hadamard matrix formed by the Kronecker product of  $\begin{pmatrix} + & + \\ + & - \end{pmatrix}$  with (5.7.a) by  $k_0, \dots, k_7$ . We then form the following  $F_i = k_i^t h_0$ , for  $i \in \{1, 2, 3, 4, 5, 6, 7\}$ .

[illegible]

## Appendix

We then use these to form

$$F = \left( \begin{array}{c|c|c|c|c|c|c} - & - & - & - & - & - & - \\ + & + & + & + & + & + & + \\ - & - & - & - & - & - & - \\ + & + & + & + & + & + & + \\ - & - & - & - & - & - & - \\ + & + & + & + & + & + & + \\ - & - & - & - & - & - & - \\ + & + & + & + & + & + & + \end{array} \right).$$

Taking  $E = F^t$ , we find that  $\begin{pmatrix} J_8 & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$  is the required balancedly splittable Hadamard matrix of order 64, which is too large to display here.

### A2. A Balancedly Splittable BW(36, 36; 6)

Beginning with (5.8.a), a Butson Hadamard matrix of order 3 over the complex 3-rd roots of unity, we can construct a balancedly splittable BW(36, 36; 6).

First, we rewrite (5.8.a) as

$$H = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 4 & 2 \\ 0 & 2 & 4 \end{pmatrix},$$

where the entries of the matrix are the logarithms of  $-e^{\frac{2\pi\sqrt{-1}}{3}}$ .

The auxiliary matrices of  $H$  are then given by

$$c_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, c_1 = \begin{pmatrix} 0 & 4 & 2 \\ 2 & 0 & 4 \\ 4 & 2 & 0 \end{pmatrix}, c_2 = \begin{pmatrix} 0 & 2 & 4 \\ 4 & 0 & 2 \\ 2 & 4 & 0 \end{pmatrix}.$$



## Appendix

The block-circulant matrix  $A$  has first row  $(c_0, c_1, c_2, c_2, c_1)$  and is given by

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 & 2 & 4 & 0 & 4 & 2 \\ 0 & 0 & 0 & 2 & 0 & 4 & 4 & 0 & 2 & 4 & 0 & 2 & 2 & 0 & 4 \\ 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 & 2 & 4 & 0 & 4 & 2 & 0 \\ \hline 0 & 4 & 2 & 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 & 2 & 4 \\ 2 & 0 & 4 & 0 & 0 & 0 & 2 & 0 & 4 & 4 & 0 & 2 & 4 & 0 & 2 \\ 4 & 2 & 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 & 2 & 4 & 0 \\ \hline 0 & 2 & 4 & 0 & 4 & 2 & 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 \\ 4 & 0 & 2 & 2 & 0 & 4 & 0 & 0 & 0 & 2 & 0 & 4 & 4 & 0 & 2 \\ 2 & 4 & 0 & 4 & 2 & 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 \\ \hline 0 & 2 & 4 & 0 & 2 & 4 & 0 & 4 & 2 & 0 & 0 & 0 & 0 & 4 & 2 \\ 4 & 0 & 2 & 4 & 0 & 2 & 2 & 0 & 4 & 0 & 0 & 0 & 2 & 0 & 4 \\ 2 & 4 & 0 & 2 & 4 & 0 & 4 & 2 & 0 & 0 & 0 & 0 & 4 & 2 & 0 \\ \hline 0 & 4 & 2 & 0 & 2 & 4 & 0 & 2 & 4 & 0 & 4 & 2 & 0 & 0 & 0 \\ 2 & 0 & 4 & 4 & 0 & 2 & 4 & 0 & 2 & 2 & 0 & 4 & 0 & 0 & 0 \\ 4 & 2 & 0 & 2 & 4 & 0 & 2 & 4 & 0 & 4 & 2 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

The block-circulant matrix  $B$  has first row  $(c_0, c_1, c_2, -c_2, -c_1)$  and is given by

$$B = \begin{pmatrix} 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 3 & 5 & 1 & 3 & 1 & 5 \\ 0 & 0 & 0 & 2 & 0 & 4 & 4 & 0 & 2 & 1 & 3 & 5 & 5 & 3 & 1 \\ 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 & 5 & 1 & 3 & 1 & 5 & 3 \\ \hline 3 & 1 & 5 & 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 3 & 5 & 1 \\ 5 & 3 & 1 & 0 & 0 & 0 & 2 & 0 & 4 & 4 & 0 & 2 & 1 & 3 & 5 \\ 1 & 5 & 3 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 & 5 & 1 & 3 \\ \hline 3 & 5 & 1 & 3 & 1 & 5 & 0 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 \\ 1 & 3 & 5 & 5 & 3 & 1 & 0 & 0 & 0 & 2 & 0 & 4 & 4 & 0 & 2 \\ 5 & 1 & 3 & 1 & 5 & 3 & 0 & 0 & 0 & 4 & 2 & 0 & 2 & 4 & 0 \\ \hline 0 & 2 & 4 & 3 & 5 & 1 & 3 & 1 & 5 & 0 & 0 & 0 & 0 & 4 & 2 \\ 4 & 0 & 2 & 1 & 3 & 5 & 5 & 3 & 1 & 0 & 0 & 0 & 2 & 0 & 4 \\ 2 & 4 & 0 & 5 & 1 & 3 & 1 & 5 & 3 & 0 & 0 & 0 & 4 & 2 & 0 \\ \hline 0 & 4 & 2 & 0 & 2 & 4 & 3 & 5 & 1 & 3 & 1 & 5 & 0 & 0 & 0 \\ 2 & 0 & 4 & 4 & 0 & 2 & 1 & 3 & 5 & 5 & 3 & 1 & 0 & 0 & 0 \\ 4 & 2 & 0 & 2 & 4 & 0 & 5 & 1 & 3 & 1 & 5 & 3 & 0 & 0 & 0 \end{pmatrix}.$$

As before, we take  $K = \begin{pmatrix} + & + \\ + & - \end{pmatrix} \otimes H$  and form  $F_i = k_i^* h_0$ , for  $i \in \{1, 2, 3, 4, 5\}$ .

$$\begin{aligned} F_1 &= \begin{pmatrix} 0 & 0 & 0 \\ 2 & 2 & 2 \\ 4 & 4 & 4 \\ 0 & 0 & 0 \\ 2 & 2 & 2 \\ 4 & 4 & 4 \end{pmatrix}, & F_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 4 & 4 & 4 \\ 2 & 2 & 2 \\ 0 & 0 & 0 \\ 4 & 4 & 4 \\ 2 & 2 & 2 \end{pmatrix}, & F_3 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \end{pmatrix}, \\ F_4 &= \begin{pmatrix} 0 & 0 & 0 \\ 2 & 2 & 2 \\ 4 & 4 & 4 \\ 3 & 3 & 3 \\ 5 & 5 & 5 \\ 1 & 1 & 1 \end{pmatrix}, & F_5 &= \begin{pmatrix} 0 & 0 & 0 \\ 4 & 4 & 4 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \\ 1 & 1 & 1 \\ 5 & 5 & 5 \end{pmatrix}. \end{aligned}$$

## Appendix

Then

$$F = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 4 & 4 & 4 & 0 & 0 & 0 & 2 & 2 & 2 \\ 4 & 4 & 4 & 2 & 2 & 2 & 0 & 0 & 0 & 4 & 4 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 3 & 3 & 3 & 3 & 3 & 3 \\ 2 & 2 & 2 & 4 & 4 & 4 & 3 & 3 & 3 & 5 & 5 & 5 \\ 4 & 4 & 4 & 2 & 2 & 2 & 3 & 3 & 3 & 1 & 1 & 1 \end{pmatrix},$$

and  $E = F^*$ .

Finally, we form the matrix  $\begin{pmatrix} J_6 & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$  given by

0000000	000000000000000000	333333333333333333
0000000	2224444000222444	5551111333555111
0000000	444222000444222	111555333111555
0000000	0000003333333333	3333333000000000
0000000	222444333555111	5551111000222444
0000000	444222333111555	1115555000444222
042042	000042024024042	000042024351315
042042	000204402402204	000204402135531
042042	000420240240420	000420240513153
024024	0420000402024024	315000042024351
024024	204000204402402	531000204402135
024024	420000420240240	153000420240513
000333	024042000042024	351315000042024
000333	402204000204402	135531000204402
000333	240420000420240	513153000420240
042315	024024042000042	024351315000042
042315	402402204000204	402135531000204
042315	240240420000420	240513153000420
024351	042024024042000	042024351315000
024351	204402402204000	204402135531000
024351	42024024020000	420240513153000
315315	000042024351315	000042024024042
315315	000204402135531	000204402402204
315315	000420240513153	000420240240420
351351	315000042024351	042000042024024
351351	531000204402135	204000204402402
351351	153000420240513	420000420240240
333000	351315000042024	024042000042024
333000	135531000204402	402204000204402
333000	513153000420240	240420000420240
315042	024351315000042	024024042000042
315042	402135531000204	402402204000204
315042	240513153000420	240240420000420
351024	042024351315000	042024024042000
351024	204402135531000	204402402204000
351024	420240513153000	420240240420000

which is a balancedly splittable  $\text{BW}(36, 36; 6)$ .

## Appendix

### A3. A Balancedly Splittable $\text{OD}(64; 16, 16, 16, 16)$

As before, we start with (6.16.a). The corresponding Hadamard matrix is (5.7.a), so we form the the auxiliary matrices given by

$$c_0 = \begin{pmatrix} a & b & c & d \\ a & b & c & d \\ a & b & c & d \\ a & b & c & d \end{pmatrix}, c_1 = \begin{pmatrix} b & \bar{a} & d & \bar{c} \\ \bar{b} & a & \bar{d} & c \\ b & \bar{a} & d & \bar{c} \\ \bar{b} & a & \bar{d} & c \end{pmatrix}, c_2 = \begin{pmatrix} c & \bar{d} & \bar{a} & b \\ \bar{c} & d & a & \bar{b} \\ \bar{c} & d & a & \bar{b} \\ c & \bar{d} & \bar{a} & b \end{pmatrix}, c_3 = \begin{pmatrix} d & c & \bar{b} & \bar{a} \\ d & c & \bar{b} & \bar{a} \\ \bar{d} & \bar{c} & b & a \\ \bar{d} & \bar{c} & b & a \end{pmatrix}.$$

The block-circulant matrix  $A$  has first row  $(c_0, c_1, c_2, c_3, c_3, c_2, c_1)$  and is given by

[illegible]

## Appendix

The block-circulant matrix  $B$  has first row  $(c_0, c_1, c_2, c_3, -c_3, -c_2, -c_1)$  and is given by

[illegible]

Taking  $Y$  to be the Kronecker product of  $\begin{pmatrix} + & + \\ + & - \end{pmatrix}$  with  $X$  and  $K$  the corresponding Hadamard matrix, we form the matrices  $F_i = k_i^t x_0$ , for  $i \in \{1, 2, 3, 4, 5, 6, 7\}$ .

$$F_1 = \begin{pmatrix} \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \end{pmatrix}, \quad F_2 = \begin{pmatrix} \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \end{pmatrix}, \quad F_3 = \begin{pmatrix} \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ a & b & c & d \end{pmatrix}, \quad F_4 = \begin{pmatrix} a & b & c & d \\ a & b & c & d \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \end{pmatrix},$$

## Appendix

$$F_5 = \begin{pmatrix} \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \end{pmatrix}, \quad F_6 = \begin{pmatrix} \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \end{pmatrix}, \quad F_7 = \begin{pmatrix} \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ a & b & c & d \\ a & b & c & d \\ a & b & c & d \\ a & b & c & d \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \\ \bar{a} & \bar{b} & \bar{c} & \bar{d} \end{pmatrix}.$$

The matrix  $F$  is then given by

$$F = \begin{pmatrix} \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} \\ a b c d & a b c d & a b c d & a b c d & a b c d & a b c d & a b c d \\ \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & a b c d & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & a b c d \\ a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & a b c d & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d \\ \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & a b c d & a b c d \\ a b c d & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d \\ \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} \\ a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} & \bar{a} \bar{b} \bar{c} \bar{d} & a b c d & \bar{a} \bar{b} \bar{c} \bar{d} \end{pmatrix}$$

We next form the matrices  $E_i = h_0^t y_i$ , for  $i \in \{1, 2, 3, 4, 5, 6, 7\}$ .

$$\begin{aligned}
E_1 &= \begin{pmatrix} \bar{b} a \bar{d} c \bar{b} a \bar{d} c \\ \bar{b} a \bar{d} c \bar{b} a \bar{d} c \\ \bar{b} a \bar{d} c \bar{b} a \bar{d} c \\ \bar{b} a \bar{d} c \bar{b} a \bar{d} c \end{pmatrix}, & E_2 &= \begin{pmatrix} \bar{c} d a \bar{b} \bar{c} d a \bar{b} \\ \bar{c} d a \bar{b} \bar{c} d a \bar{b} \\ \bar{c} d a \bar{b} \bar{c} d a \bar{b} \\ \bar{c} d a \bar{b} \bar{c} d a \bar{b} \end{pmatrix}, & E_3 &= \begin{pmatrix} \bar{d} \bar{c} b a \bar{d} \bar{c} b a \\ \bar{d} \bar{c} b a \bar{d} \bar{c} b a \\ \bar{d} \bar{c} b a \bar{d} \bar{c} b a \\ \bar{d} \bar{c} b a \bar{d} \bar{c} b a \end{pmatrix}, \\
E_4 &= \begin{pmatrix} a b c d \bar{a} \bar{b} \bar{c} \bar{d} \\ a b c d \bar{a} \bar{b} \bar{c} \bar{d} \\ a b c d \bar{a} \bar{b} \bar{c} \bar{d} \\ a b c d \bar{a} \bar{b} \bar{c} \bar{d} \end{pmatrix}, & E_5 &= \begin{pmatrix} \bar{b} a \bar{d} c b \bar{a} d \bar{c} \\ \bar{b} a \bar{d} c b \bar{a} d \bar{c} \\ \bar{b} a \bar{d} c b \bar{a} d \bar{c} \\ \bar{b} a \bar{d} c b \bar{a} d \bar{c} \end{pmatrix}, & E_6 &= \begin{pmatrix} \bar{c} d a \bar{b} c \bar{d} \bar{a} b \\ \bar{c} d a \bar{b} c \bar{d} \bar{a} b \\ \bar{c} d a \bar{b} c \bar{d} \bar{a} b \\ \bar{c} d a \bar{b} c \bar{d} \bar{a} b \end{pmatrix}, \\
E_7 &= \begin{pmatrix} \bar{d} \bar{c} b a d c \bar{b} \bar{a} \\ \bar{d} \bar{c} b a d c \bar{b} \bar{a} \\ \bar{d} \bar{c} b a d c \bar{b} \bar{a} \\ \bar{d} \bar{c} b a d c \bar{b} \bar{a} \end{pmatrix}.
\end{aligned}$$

## Appendix

The matrix  $E^t$  is then given by

$$E^t = \begin{pmatrix} \bar{b} \bar{b} \bar{b} \bar{b} & \bar{c} \bar{c} \bar{c} \bar{c} & \bar{d} \bar{d} \bar{d} \bar{d} & a a a a & \bar{b} \bar{b} \bar{b} \bar{b} & \bar{c} \bar{c} \bar{c} \bar{c} & \bar{d} \bar{d} \bar{d} \bar{d} \\ a a a a & d d d d & \bar{c} \bar{c} \bar{c} \bar{c} & b b b b & a a a a & d d d d & \bar{c} \bar{c} \bar{c} \bar{c} \\ \bar{d} \bar{d} \bar{d} \bar{d} & a a a a & b b b b & c c c c & \bar{d} \bar{d} \bar{d} \bar{d} & a a a a & b b b b \\ c c c c & \bar{b} \bar{b} \bar{b} \bar{b} & a a a a & d d d d & c c c c & \bar{b} \bar{b} \bar{b} \bar{b} & a a a a \\ \bar{b} \bar{b} \bar{b} \bar{b} & \bar{c} \bar{c} \bar{c} \bar{c} & \bar{d} \bar{d} \bar{d} \bar{d} & \bar{a} \bar{a} \bar{a} \bar{a} & b b b b & c c c c & d d d d \\ a a a a & d d d d & \bar{c} \bar{c} \bar{c} \bar{c} & \bar{b} \bar{b} \bar{b} \bar{b} & \bar{a} \bar{a} \bar{a} \bar{a} & \bar{d} \bar{d} \bar{d} \bar{d} & c c c c \\ \bar{d} \bar{d} \bar{d} \bar{d} & a a a a & b b b b & \bar{c} \bar{c} \bar{c} \bar{c} & d d d d & \bar{a} \bar{a} \bar{a} \bar{a} & \bar{b} \bar{b} \bar{b} \bar{b} \\ c c c c & \bar{b} \bar{b} \bar{b} \bar{b} & a a a a & \bar{d} \bar{d} \bar{d} \bar{d} & \bar{c} \bar{c} \bar{c} \bar{c} & b b b b & \bar{a} \bar{a} \bar{a} \bar{a} \end{pmatrix}.$$

Finally, we form

[illegible]

whereupon the matrix  $\begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$  is the required balancedly splittable OD(64; 16, 16, 16, 16).

## Appendix

# Bibliography

- Bannai, E., Bannai, E., Ito, T., and Tanaka, R. (2021). *Algebraic combinatorics*. De Gruyter.
- Bannai, E. and Ito, T. (1984). *Algebraic combinatorics. I. The Benjamin/Cummings Publishing Co., Inc., Menlo Park, CA. Association schemes*.
- Batten, L. M. (1997). *Combinatorics of finite geometries*. Cambridge University Press, Cambridge, second edition.
- Best, D. and Kharaghani, H. (2010). Unbiased complex Hadamard matrices and bases. *Cryptogr. Commun.*, 2(2):199–209.
- Best, D., Kharaghani, H., and Ramp, H. (2015). Mutually unbiased weighing matrices. *Des. Codes Cryptogr.*, 76(2):237–256.
- Beth, T., Jungnickel, D., and Lenz, H. (1999). *Design theory. Vols. I and II*, volume 69 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition.
- Boykin, O. P., Sitharam, M., Tarifi, M., and Wocjan, P. (2005). Real mutually unbiased bases.
- Bracken, C., McGuire, G., and Ward, H. (2006). New quasi-symmetric designs constructed using mutually orthogonal Latin squares and Hadamard matrices. *Des. Codes Cryptogr.*, 41(2):195–198.
- Brouwer, A. E., Cohen, A. M., and Neumaier, A. (1989). *Distance-regular graphs*, volume 18 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin.
- Cameron, P. J. (1994). *Combinatorics: topics, techniques, algorithms*. Cambridge University Press, Cambridge.
- Cameron, P. J. (1999). *Permutation groups*, volume 45 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge.
- Cameron, P. J. and van Lint, J. H. (1991). *Designs, graphs, codes and their links*, volume 22 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge.



## Bibliography

- Christensen, O. (2016). *An introduction to frames and Riesz bases*. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, [Cham], second edition.
- Colbourn, C. J. and Dinitz, J. H., editors (2007). *Handbook of combinatorial designs*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, second edition.
- de Launey, W. and Sarvate, D. G. (1984). Nonexistence of certain GBRDs. *Ars Combin.*, 18:5–20.
- Delsarte, P. (1973). An algebraic approach to the association schemes of coding theory. *Philips Res. Rep. Suppl.*, (10):vi+97.
- Delsarte, P., Goethals, J. M., and Seidel, J. J. (1977). Spherical codes and designs. *Geometriae Dedicata*, 6(3):363–388.
- Dembowski, P. (1997). *Finite geometries*. Classics in Mathematics. Springer-Verlag, Berlin. Reprint of the 1968 original.
- Fender, K., Kharaghani, H., and Suda, S. (2018). On a class of quaternary complex Hadamard matrices. *Discrete Math.*, 341(2):421–426.
- Fisher, R. A. (1940). An examination of the different possible solutions of a problem in incomplete blocks. *Ann. Eugenics*, 10:52–75.
- Gibbons, P. B. and Mathon, R. A. (1987). Group signings of symmetric balanced incomplete block designs. In *Proceedings of the Singapore conference on combinatorial mathematics and computing (Singapore, 1986)*, volume 23, pages 123–134.
- Hall, Jr., M. (1986). *Combinatorial theory*. Wiley-Interscience Series in Discrete Mathematics. John Wiley & Sons, Inc., New York, second edition. A Wiley-Interscience Publication.
- Halmos, P. R. (1982). *A Hilbert space problem book*, volume 17 of *Encyclopedia of Mathematics and its Applications*. Springer-Verlag, New York-Berlin, second edition.
- Halmos, P. R. (1998). *Introduction to Hilbert space and the theory of spectral multiplicity*. AMS Chelsea Publishing, Providence, RI. Reprint of the second (1957) edition.
- Hamming, R. W. (1950). Error detecting and error correcting codes. *Bell System Tech. J.*, 29:147–160.
- Han, D., Kornelson, K., Larson, D., and Weber, E. (2007). *Frames for undergraduates*, volume 40 of *Student Mathematical Library*. American Mathematical Society, Providence, RI.

## Bibliography

- Hill, R. (1986). *A first course in coding theory*. Oxford Applied Mathematics and Computing Science Series. The Clarendon Press, Oxford University Press, New York.
- Holzmann, W. H., Kharaghani, H., and Orrick, W. (2010). On the real unbiased Hadamard matrices. In *Combinatorics and graphs*, volume 531 of *Contemp. Math.*, pages 243–250. Amer. Math. Soc., Providence, RI.
- Huffman, W. C. and Pless, V. (2003). *Fundamentals of error-correcting codes*. Cambridge University Press, Cambridge.
- Ionin, Y. J. (2001). Applying balanced generalized weighing matrices to construct block designs. *Electron. J. Combin.*, 8(1):Research Paper 12, 15.
- Ionin, Y. J. and Kharaghani, H. (2003a). Doubly regular digraphs and symmetric designs. *J. Combin. Theory Ser. A*, 101(1):35–48.
- Ionin, Y. J. and Kharaghani, H. (2003b). New families of strongly regular graphs. *J. Combin. Des.*, 11(3):208–217.
- Ionin, Y. J. and Shrikhande, M. S. (2006). *Combinatorics of symmetric designs*, volume 5 of *New Mathematical Monographs*. Cambridge University Press, Cambridge.
- Isaacs, I. M. (2006). *Character theory of finite groups*. AMS Chelsea Publishing, Providence, RI. Corrected reprint of the 1976 original [Academic Press, New York; MR0460423].
- Kharaghani, H. (1985). New class of weighing matrices. *Ars Combin.*, 19:69–72.
- Kharaghani, H., Pender, T., and Suda, S. (2021). Balancedly splittable orthogonal designs and equiangular tight frames. *Des. Codes Cryptogr.*, 89(9):2033–2050.
- Kharaghani, H., Pender, T., and Suda, S. (2022). A family of balanced generalized weighing matrices. *Combinatorica*, to appear.
- Kharaghani, H. and Suda, S. (2018). Unbiased orthogonal designs. *Des. Codes Cryptogr.*, 86(7):1573–1588.
- Kharaghani, H. and Suda, S. (2019). Balancedly splittable Hadamard matrices. *Discrete Math.*, 342(2):546–561.
- Kharaghani, H. and Suda, S. (2021). Divisible design digraphs and association schemes. *Finite Fields Appl.*, 69:Paper No. 101763, 17.
- Lam, T. Y. and Leung, K. H. (2000). On vanishing sums of roots of unity. *J. Algebra*, 224(1):91–109.

## Bibliography

- MacWilliams, F. J. and Sloane, N. J. A. (1977). *The theory of error-correcting codes. I and II*. North-Holland Publishing Co., Amsterdam-New York-Oxford. North-Holland Mathematical Library, Vol. 16.
- McGuire, G. (1997). Quasi-symmetric designs and codes meeting the Grey-Rankin bound. *J. Combin. Theory Ser. A*, 78(2):280–291.
- Nozaki, H. and Suda, S. (2015). Weighing matrices and spherical codes. *J. Algebraic Combin.*, 42(1):283–291.
- Paley, R. E. A. C. and Wiener, N. (1987). *Fourier transforms in the complex domain*, volume 19 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI. Reprint of the 1934 original.
- Pender, T. S. (2020). Balanced group matrices. honors thesis.
- Rajkundlia, D. P. (1983). Some techniques for constructing infinite families of BIBDs. *Discrete Math.*, 44(1):61–96.
- Seberry, J. (2017). *Orthogonal designs*. Springer, Cham. Hadamard matrices, quadratic forms and algebras, Revised and updated edition of the 1979 original [MR0534614].
- Seberry, J. and Yamada, M. (1992). Hadamard matrices, sequences, and block designs. In *Contemporary design theory*, Wiley-Intersci. Ser. Discrete Math. Optim., pages 431–560. Wiley, New York.
- Shrikhande, M. S. and Sane, S. S. (1991). *Quasi-symmetric designs*, volume 164 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge.
- Stinson, D. R. (2004). *Combinatorial designs*. Springer-Verlag, New York. Constructions and analysis, With a foreword by Charles J. Colbourn.
- Waldron, S. (2020). Tight frames over the quaternions and equiangular lines.
- Waldron, S. F. D. (2018). *An introduction to finite tight frames*. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, New York.
- Wallis, W. D., Street, A. P., and Wallis, J. S. (1972). *Combinatorics: Room squares, sum-free sets, Hadamard matrices*. Lecture Notes in Mathematics, Vol. 292. Springer-Verlag, Berlin-New York.
- Young, R. M. (2001). *An introduction to nonharmonic Fourier series*. Academic Press, Inc., San Diego, CA, first edition.

# Index

- adjacency algebra, 18
- admissible group of symmetries, 55, 56
- analysis operator, 44
- aperiodic autocorrelation, 14
- association scheme, 16
- auxiliary matrices, 25, 34
- balanced generalized weighing matrix, 11
  - normal form, 12
- balanced incomplete block design, 2
  - automorphism group, 5
  - complement design, 4
  - derived design, 4
  - incidence matrix of, 3
  - isomorphisms of, 5
  - normal form, 5
  - residual design, 4
  - symmetric design, 3
- balancedly splittable Hadamard matrix, 23
- balancedly splittable orthogonal designs, 28
- Butson weighing matrix, 9
- c-GBRD, 12
- character tables, 19
- circulant matrix, 13
  - complementary, 14
- complementary sequences, 14
- complex orthogonal design, 13
- conference matrix, 9
- conjugate transpose, 10
- cross-correlation, 15
- decomposition matrices, 58
- dual-eigenvalues of a scheme, 18
- eigenvalues of a scheme, 18
- elementary abelian group, 11
- equiangular lines, 23
- Fisher's inequality, 3
- frame, 44
  - bounds of, 44
- generalized Bhaskar Rao design, 10
- generalized Hadamard matrix, 11
- generalized weighing schemes, 85
- group of symmetries, 54
- Hadamard design, 75
- Hadamard matrix, 9
- Hamming code, 8
- Hamming distance, 7
- Hamming weight, 7
- intersection matrix, 19
- intersection numbers, 16
- Kronecker product, 52
- linear error-correcting code, 6
  - dual of, 7
- monomial equivalence, 11
- multiplicities, 18
- mutually unbiased bases, 47
- orthogonal design, 13
- Parseval frame, 44

## Index

- periodic autocorrelation, 14
- quaternary orthogonal design, 13
- regular Hadamard matrix, 53
- Schur idempotents, 18
- simplex code, 8
- standard form, 19
- synthesis operator, 44
- tight frame, 44
- unbiased Hadamard matrices, 47
- unbiased orthogonal designs, 48
- weighing matrix, 9
- weighing schemes, 69