# Balancedly Splittable Orthogonal Designs

Thomas Pender
\*\*\*Joint work with Hadi Kharaghani, Sho Suda

Department of Mathematics and Computer Science University of Lethbridge

Alberta-Montana Combinatorics & Algorithms Days at BIRS June 2022



# Summary

- balancedly splittable Hadamard matrices
- balancedly splittable orthogonal designs
- construction
- related configurations

#### **Definition:** Hadmard Matrix

- A square matrix with entries from  $\{-1,1\}$  with pairwise orthogonal rows.
- Equiv.  $HH^t = nI_n$ , with H an  $n \times n$  (-1,1)-matrix of order n.



• a Hadamard matrix of order 16



#### **Definition:** Hadmard Matrix

- A square matrix with entries from  $\{-1,1\}$  with pairwise orthogonal rows.
- Equiv.  $HH^t = nI_n$ , with H an  $n \times n$  (-1,1)-matrix of order n.



#### **Definition:** Hadmard Matrix

- A square matrix with entries from  $\{-1,1\}$  with pairwise orthogonal rows.
- Equiv.  $HH^t = nI_n$ , with H an  $n \times n$  (-1, 1)-matrix of order n.

## **Definition:** Balancedly Splittable (Kharaghani and Suda, 2019)

- ullet A Hadamard matrix H of order n with  $n imes \ell$  submatrix  $H_1$  .
- H is  $(n, \ell, a, b)$  balancedly splittable, a < b, if

$$H_1H_1^t=\ell I_n+aA+b(J_n-I_n-A),$$

for symmetric (0,1)-matrix A with zero diagonal.



• a Hadamard matrix of order 16



6 / 47

• a balancedly splittable Hadamard matrix of order 16



```
\begin{pmatrix} 1-1-1&1&1-1&-1&1&-1&1&-\\ 1-1&-1&1&-1&-1&1&1&1&-&-1\\ 1&1&-1&-1&1&1&-1&-1&1&-1\\ 1&1&-&-1&1&1&-1&-1&1&1&1&-\\ 1-&-1&1&-1&-1&1&1&-&1&1\\ 1-&-1&-1&-1&1&1&1&-&1&1\\ 1-&-1&-1&-1&1&1&1&-&1&1 \end{pmatrix}
```

8 / 47

```
2 2 2 2 2 6 2 2 2 2 2 2 2 2 2 2 2
2 2 2 2 2 2 2 2 6 2 2 2 2 2 2 2
2 5 5 2 5 2 5 2 5 6 5 2 2 5 2 2 2
22222226222
22222222226
```

• The rows of the split are equiangular.



• There are four splits.



#### **Definition:** Orthogonal Designs

• A square  $\{\pm z_1, \dots, \pm z_u\}$ -matriz X of order n such that

$$XX^t = \sigma I_n, \qquad s_i \in \mathbb{N},$$

where 
$$\sigma = \sum_{i} s_{i} z_{i}^{2}$$
.

• Write  $OD(n; s_1, \ldots, s_u)$ .



ābāabbābāabbāba bābāabbabaabbabā a b ā b b ā a b b ā b a a b b ā a b ā b b a a b b a b ā a b b a bābabābāabbābaab ābababaabbabāab abaabbābaabbābā a b a a b b a b a a b b a b a babbaabbābā abbā b̄abbāabbabaabba a b ā b ā b̄ a a b b ā b ā a b a b ā b̄ a b ā a b b̄ a b̄ a a b

• An OD(16; 8, 8).

ababababababābābāb ababābābababāb ā b ā a b b ā b ā a b b ā b a ā a b b a b a a b b a b ā bābbāabbābaabbā bā bā a a bā a bā a bā a ābabābāabbābaab ā b a b a b a a b b a b a a b a b̄ a a b b ā b̄ a a b b ā b ā a b̄ a a b b̄ a b ā a b b̄ a b̄ a babbaabbabaabba  $\bar{b}$  a b  $\bar{b}$   $\bar{a}$  a b  $\bar{b}$  a  $\bar{b}$  a a b  $\bar{b}$  a a b ā b ā b a a b b ā b ā a b abābabāabbabaab



labababababā bābābāb ababābabābabābab abababābābābābabab ababābābababāb bābāabbābāabbāba bābāabbabaabbabā abābbāabbābaabbā abābbaabbabāabba bābabābāabbābaab bābababaabbabāab babaabbābaabbābā a b̄ a a b b̄ a b ā a b b̄ a b̄ a ābabbaabbābāabbā 

• Products in the set  $\{\pm 2a^2, \pm 2b^2, \pm 2ab\}$ .

## **Definition:** Orthogonal Designs

• A square  $\{\pm z_1, \dots, \pm z_u\}$ -matrix X of order n such that  $XX^t = \sigma I_n, s_i \in \mathbb{N}$ , where  $\sigma = \sum_i s_i z_i^2$ .



#### **Definition:** Orthogonal Designs

• A square  $\{\pm z_1, \dots, \pm z_u\}$ -matrix X of order n such that  $XX^t = \sigma I_n, s_i \in \mathbb{N}$ , where  $\sigma = \sum_i s_i z_i^2$ .

## Definition: Balancedly Splittable (Kharaghani et al., 2021)

• An OD X with  $n \times \ell$  submatrix  $X_1$  admits a stable  $(n, \ell, a, b)$  split if

$$X_1X_1^t = \sigma(cI_n + aA + b(J_n - I_n - A)).$$

The split is unstable if it is not stable and

$$X_1X_1^t = \ell I_n + aA + b(J_n - I_n - A)$$

after replacing each variable with 1.



## **Definition:** Auxiliary Matrices

- X a full  $OD(n; s_1, \ldots, s_u)$  and H the Hadamard matrix obtained by replacing each variable with 1.
- Label rows of X and H by  $x_0, \ldots, x_{n-1}$ , and  $h_0, \ldots, h_{n-1}$ .
- The auxiliary matrices of X are given by

$$c_i = h_i^t x_i, \qquad i = 0, \dots, n-1.$$



• An *OD*(2; 1, 1):

$$\begin{pmatrix} a & b \\ b & -a \end{pmatrix}$$

Auxiliary matrices:

$$c_0 = \begin{pmatrix} a & b \\ a & b \end{pmatrix}, \qquad c_1 = \begin{pmatrix} b & -a \\ -b & a \end{pmatrix}$$



Form the sequences

$$\alpha = (c_0, c_1, c_1), \quad \beta = (c_0, c_1, -c_1).$$

• The sum of the periodic autocorrelations of  $\alpha$  and  $\beta$  satisfy

$$\sum_{i} \alpha_{i} \alpha_{i+j}^{t} + \sum_{i} \beta_{i} \beta_{i+j}^{t} = 0, \qquad j = 1, 2.$$

Sums of cross-correlations

$$\sum_{i} \alpha_{i} \beta_{i+j}^{t} + \sum_{i} \beta_{i} \alpha_{i+j}^{t} = 0, \quad j = 1, 2.$$



Form the matrices

$$A = circ(\alpha), \qquad B = circ(\beta),$$

• Then the matrix

$$\Theta = \begin{pmatrix} A & B \\ B & A \end{pmatrix}.$$



20 / 47

$$\Theta = \begin{pmatrix} a & b & b & \bar{a} & b & \bar{a} & a & b & b & \bar{a} & \bar{b} & a \\ a & b & \bar{b} & a & \bar{b} & a & a & b & \bar{b} & a & b & \bar{a} \\ b & \bar{a} & a & b & \bar{b} & \bar{a} & \bar{b} & a & a & b & \bar{b} & \bar{a} \\ \bar{b} & a & a & b & \bar{b} & a & b & \bar{a} & a & b & \bar{b} & a \\ b & \bar{a} & b & \bar{a} & a & b & \bar{b} & a & b & \bar{a} & a & b \\ \bar{b} & a & \bar{b} & a & a & b & \bar{b} & a & \bar{b} & \bar{a} & \bar{b} \\ \bar{a} & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} & a & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} \\ \bar{b} & a & a & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} & a & b & \bar{b} & \bar{a} \\ \bar{b} & \bar{a} & a & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} & a & b & \bar{b} & \bar{a} \\ \bar{b} & \bar{a} & \bar{b} & \bar{a} & a & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} & \bar{b} \\ \bar{b} & \bar{a} & \bar{b} & \bar{a} & \bar{a} & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} & \bar{b} \\ \bar{b} & \bar{a} & \bar{b} & \bar{a} & \bar{a} & b & \bar{b} & \bar{a} & \bar{b} & \bar{a} & \bar{a} & \bar{b} \end{pmatrix}$$



$$\Theta\Theta^{t} = \begin{pmatrix} AA^{t} + BB^{t} & AB^{t} + BA^{t} \\ AB^{t} + BA^{t} & AA^{t} + BB^{t} \end{pmatrix}.$$



0



```
ā
        a b
        a b \bar{b}
  abbābaabbā
  a b b a b a a b b a
bābāabbābaab
        Ба b ā a b
    āb
        a b
 b b a b a a b b a b
  abbābāabbā
  abbabaabba
bābaabbābā
babāabbabaa
```

Define:

$$Y = \begin{pmatrix} 1 & 1 \\ 1 & - \end{pmatrix} \otimes X,$$
  
 $K = \begin{pmatrix} 1 & 1 \\ 1 & - \end{pmatrix} \otimes H.$ 

• Index rows as  $y_0, ..., y_{2n-1}$ , and  $k_0, ..., k_{2n-1}$ .



25 / 47

$$Y = \begin{pmatrix} 1 & 1 \\ 1 & - \end{pmatrix} \otimes X$$
  $K = \begin{pmatrix} 1 & 1 \\ 1 & - \end{pmatrix} \otimes H$   
Rows:  $y_0, \dots, y_{2n-1}$  Rows:  $k_0, \dots, k_{2n-1}$ 

#### Form

$$F = \begin{pmatrix} k_1^t x_0 & k_2^t x_0 & k_3^t x_0 \end{pmatrix},$$

$$E = \begin{pmatrix} h_0^t y_1 \\ h_0^t y_2 \\ h_0^t y_3 \end{pmatrix},$$

$$G = k_0^t y_0.$$



/ G	F							_F						
E	а	b	b	ā	b	ā	ć	7	b	b	ā	b	а	_
	a	b	Б	a	Б	а	ã		b	Б	a	b	ā	
	b	ā	a	b	b	ā	Ī	,	a	a	b	b	ā	
	$\bar{b}$	a	a	b	$\bar{b}$	а	Ł	)	ā	a	b	Б	a	
	Ь	ā	b	ā	a	b	L	)	ā	Б	a	a	b	
	$\bar{b}$	a	Б	a	a	b	Ē	5	a	b	ā	a	b	
E	а	b	b	ā	b	а	á	3	b	b	ā	b	ā	_
	a	b	$\bar{b}$	a	b	ā	ä	9	b	$\bar{b}$	a	$\bar{b}$	a	
	$\bar{b}$	a	a	b	b	ā	L		ā	a	b	b	ā	
	b	ā	a	b	$\bar{b}$	а	Ī	,	a	a	b	Б	a	
	b	ā	$\bar{b}$	a	a	b	Ł	)	ā	b	ā	a	b	
	$\bar{b}$	a	b	ā	а	b	Ē	,	a	$\bar{b}$	a	а	b	,



ababā bā bababā b bābāabbābāabbāba ā a b b a b a a b b a b ā bābbāabbābaabbā bābbaabbabāabba ābabābāabbābaab ābabaabbabāab a b a a b b ā b a a b b ā b ā baabbabāabbaba babbaabbabaabba Ба b b ā a b Ба b a a b Ба abābābaabbābāab abābabāabbabaab



• The following is a OD(2; 1, 1) of quaternions.

$$\begin{pmatrix} \bar{a} & bi \\ \bar{b}j & ak \end{pmatrix},$$

• where a and b are real variables.

bi ā ā bi bi bi ā а bi ā bi a bi ā bi a bi ā bi ā bi a bi a bi ā bi a bi a bi ā Бi ā bi Бі Бi a bi ā bi ā bi ā bi bj ak bj ak ā bi bj āk bj āk bj ak ā bi bj ak ā bi bj ak ā bi bj āk ā bi bj āk ā bi bj āk bj ak ā ak bj āk bj ak bj ak ā bi bj ak bj āk ak bi āk bi āk ā bi bj āk bj ak āk bj āk ā bi bj ak bj āk ā bi bj ak bj ak bj āk bj āk ā bi bj āk bj ak ā bi bj āk bj bi ā bi bj āk ā bi bj ak bj ak ā bi ā bi bj ak ā ā ā āk bj ak bj āk ā bi bj ak bj ak āk bj ak bj āk bj ak ā bi bj āk bj āk ā

< ロト < 個 ト < 重 ト < 重 ト ■ ■ の < @

#### **Definition:** Equiangular Lines

A collection  $\mathcal{L} \subset \mathbb{R}^{\ell}$  of lines (vectors) are equiangular if there is a constant a such that  $|\langle u, v \rangle| = a$  for every  $u, v \in L$ .

#### **Definition:** Unbiased Hadamard Matrices

Hadamard matrices H and K are unbiased if

$$HK^t = \sqrt{n}L$$

for some Hadamard matrix L.

<ロ > < 回 > < 回 > < 巨 > < 巨 > 三 の < (で)

• a Hadamard matrix of order 16

• a balanced split

◆ロト ◆個ト ◆差ト ◆差ト 差 めるぐ

33 / 47

- a collection of equiangular lines
- products are the same in absolute value

• H and K are unbiased

4回 → 4回 → 4 三 → 4 三 → 9 Q ○

### Proposition (Kharaghani and Suda, 2019)

 $H = (H_1 H_2)$  a Hadamard matrix of order n with

$$H_1H_1^t = \ell I_n + aS$$

where S is a (-1,0,1)-matrix. The following are equivalent:

- **1**  $K = (-H_1 H_2)$  is unbiased with H, and
- ②  $(\ell, a) = ((n \pm \sqrt{n})/2, \sqrt{n}/2).$

$$\begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$$

$$\begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$$

$$\begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$$

39 / 47

$$\begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}$$

- each split gives equiangular lines
- parametric conditions of the proposition are satisfied
- we can always construct unbiased Hadamard matrices

# **Definition:** Unbiased ODs (Kharaghani and Suda, 2018)

- $X_1$  and  $X_2$  be two  $OD(n, s_1, \ldots, s_u)$  in  $\{\pm z_1, \ldots, \pm z_u\}$ .
- $X_1$  and  $X_2$  are unbiased if there is a Hadamard matrix H such that

$$X_1 X_2^t = \left(\alpha^{-\frac{1}{2}} \sum_i s_i z_i^2\right) H.$$

for some  $\alpha \in \mathbb{R}^+$ .



#### Definition: Unbiased ODs (Kharaghani and Suda, 2018)

- $X_1$  and  $X_2$  be two  $OD(n, s_1, \ldots, s_u)$  in  $\{\pm z_1, \ldots, \pm z_u\}$ .
- $X_1$  and  $X_2$  are unbiased if there is a Hadamard matrix H such that

$$X_1 X_2^t = \left(\alpha^{-\frac{1}{2}} \sum_i s_i z_i^2\right) H.$$

for some  $\alpha \in \mathbb{R}^+$ .

stable split equiangular lines param. cond.  $\Rightarrow$  unbiased ODs



Recall:

$$X = \begin{pmatrix} G & F & -F \\ E & A & B \\ -E & B & A \end{pmatrix}.$$

Take

$$Y = \begin{pmatrix} G & F & F \\ E & A & -B \\ -E & B & -A \end{pmatrix}.$$

• Then  $XY^t = 2\left(\sum_i s_i z_i^2\right) H$ , where H is Hadamard.



(a a a a b b a a b b b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b a a b b a a b b a a b b a a b b b a a a b b b a a a b b b a a a b b b a a a b b b a a a b b b b a a a a b b b b b b b a a a a b b b b b b b b a a a a b b b a a a a b b b a b a a a b a b b a a a a b a b b a a a a b a b b a a a a b a b b b a b a a a b a b b a a a a b a b b a a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a a b a b a a a a b a b a a a b a b a a a a b a b a a a a b a b a a a b a b a a a a b a b a a a a b a b a a a a b a b a a a b a b a a a b a b a a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a b a b a a a a b a b a a a b a b a a a a b a b a a a a b a b a a a a b a b a a a a b a b a a a a b a b a a a a a b a b a a a a b a b a a a a a

4□ b 4□ b 4 = b 4 = b 3 = 90

# The End!!

Thank You!



## References

- Kharaghani, H., Pender, T., and Suda, S. (2021). Balancedly splittable orthogonal designs and equiangular tight frames. *Des. Codes Cryptogr.*, 89(9):2033–2050.
- Kharaghani, H. and Suda, S. (2018). Unbiased orthogonal designs. *Des. Codes Cryptogr.*, 86(7):1573–1588.
- Kharaghani, H. and Suda, S. (2019). Balancedly splittable Hadamard matrices. *Discrete Math.*, 342(2):546–561.