

A Six-Parameter Extension of the Direct Somos–4 Algebraic Generating Function

Thomas Scheuerle

Abstract

The preceding paper constructed a five-parameter quadratic algebraic generating function whose first four Hankel determinants are (λ, m, r, η) and whose Hankel sequence satisfies a normalized Somos–4 recurrence with coefficient τ . We ask whether the same method extends to the general two-coefficient recurrence

$$H_{k+4}H_k = \alpha H_{k+3}H_{k+1} + \tau H_{k+2}^2.$$

A weighted transport with $q^6 = \alpha$ gives an exact six-parameter construction and preserves the prescribed ordinary Hankel determinants. However, its selected generating-function branch contains fractional powers of α ; already its first nonconstant coefficient is $q^{-5}r$. This leads to the sharper question whether another quadratic-algebraic representative can be defined over the rational parameter field while retaining the companion normalization $D_N = H_{N+2}$.

We show that the difficulty belongs neither to the Somos–4 orbit nor to its associated genus-one curve. The Hankel sequence is defined over the rational six-parameter field, and the transported discriminant has the rational quartic model

$$Y^2 = (1 - (A + \alpha B)z + \tau z^2)^2 - 4\alpha^2 z^3.$$

We then compare the construction with the generic quadratic-orthogonal-pair realization of Chang and Liu. An explicit shift identifies their bilateral orbit with ours, and their subdiagonal data satisfy $d_n = B_{n+2}$. The comparison also separates three notions that need not descend together: the Hankel sequence, the algebraic curve, and a chosen moment branch. The first two descend, whereas the distinguished transported and Chang–Liu branches do not descend generically. We rule out several natural repairs, including parity rescaling, geometric scaling, and injective finite prefixes over the base field. A genuinely non-diagonal construction from the invariant even and odd parts remains possible, so no general nonexistence claim is made.

1 Introduction

The purpose of this paper is to extend the five-parameter generating function of [2] and to understand a difficulty that becomes visible in the extension. The previous construction prescribes

$$(H_1, H_2, H_3, H_4) = (\lambda, m, r, \eta)$$

and produces the recurrence

$$H_{k+4}H_k = H_{k+3}H_{k+1} + \tau H_{k+2}^2, \quad k \geq 1. \tag{1}$$

The natural generalization is to free the coefficient of the first bilinear term:

$$H_{k+4}H_k = \alpha H_{k+3}H_{k+1} + \tau H_{k+2}^2, \quad k \geq 1. \tag{2}$$

The six independent parameters are therefore $(\lambda, m, r, \eta, \alpha, \tau)$, and $\alpha = 1$ recovers the normalization of the preceding paper.

Two questions guide the discussion. First, can the earlier algebraic construction be extended to arbitrary values of these six parameters while preserving the four prescribed Hankel determinants? We answer this question affirmatively by a weighted change of parameters and variable. Second, can one obtain such a generating function directly over

$$K = \mathbb{Q}(\lambda, m, r, \eta, \alpha, \tau)$$

without fractional powers of α , while also retaining the ungraded companion identity $D_N = H_{N+2}$? The weighted construction does not answer the second question: it naturally lives over a field containing a sixth root of α , and its first nonconstant coefficient proves that this dependence is genuine for the selected branch.

The resulting investigation is easiest to follow by distinguishing three objects:

the Hankel sequence, the associated algebraic curve, a chosen generating-function branch.

The first two objects admit descriptions over K . A chosen branch need not. This distinction explains why removing radicals from the curve does not by itself remove them from the normalized generating function.

Sections 2 and 3 construct the six-parameter extension and locate the first fractional power. Section 4 shows that the Hankel orbit and the associated curve nevertheless descend to K . Section 5 compares the result with Chang and Liu's generic quadratic-algebraic Hankel construction and gives an explicit dictionary between the two descriptions. Section 6 studies the remaining branch and normalization problem and records several transformations that cannot solve it. Section 7 states precisely what remains open.

2 From the five-parameter construction to six parameters

2.1 The five-parameter input

Let

$$G_{1;\lambda,m,r,\eta,\tau}(x) = \sum_{k \geq 0} g_k x^k$$

be the selected algebraic series of [2]. Define

$$H_N(g) = \det(g_{i+j})_{0 \leq i,j \leq N-1}, \quad D_N(g) = \det(g_{i+j+1})_{0 \leq i,j \leq N-1}.$$

The normalization in the preceding paper is

$$(H_1, H_2, H_3, H_4) = (\lambda, m, r, \eta), \quad D_N = H_{N+2}. \quad (3)$$

Its quadratic discriminant factors as

$$\Delta_1(x) = (x\lambda^2\eta mr^2)^2 \left((1 - Jx + \tau x^2)^2 - 4x^3 \right), \quad (4)$$

where

$$J = \frac{\lambda\eta}{mr} + \frac{m^2}{\lambda r} + \frac{r^2}{m\eta} + \frac{\tau mr}{\lambda\eta}. \quad (5)$$

These statements are the only inputs from the five-parameter construction needed below.

2.2 Weighted transport

Work over

$$\mathbb{K} = \mathbb{Q}(\lambda, m, r, \eta, q, \tau), \quad q^6 = \alpha,$$

and introduce the primed parameters

$$\lambda' = \lambda, \quad m' = q^{-2}m, \quad r' = q^{-6}r, \quad \eta' = q^{-12}\eta, \quad \tau' = q^{-8}\tau. \quad (6)$$

Define

$$G_{\alpha; \lambda, m, r, \eta, \tau}(x) := G_{1; \lambda', m', r', \eta', \tau'}(qx). \quad (7)$$

If

$$G_{1; \lambda', m', r', \eta', \tau'}(x) = \sum_{k \geq 0} a_k x^k, \quad G_{\alpha; \lambda, m, r, \eta, \tau}(x) = \sum_{k \geq 0} b_k x^k,$$

then

$$b_k = q^k a_k. \quad (8)$$

Proposition 1 (Determinant weights). *For every $N \geq 1$,*

$$H_N(b) = q^{N(N-1)} H_N(a), \quad D_N(b) = q^{N^2} D_N(a). \quad (9)$$

Proof. In the ordinary Hankel matrix, its (i, j) entry is multiplied by q^{i+j} . Factoring q^i from row i and q^j from column j gives the exponent $2 \sum_{j=0}^{N-1} j = N(N-1)$. For the shifted matrix the entry is multiplied by q^{i+j+1} . The additional factor q in every row contributes N , so the total exponent is $N(N-1) + N = N^2$. $\square$

2.3 The six-parameter Somos-4 recurrence

Theorem 2. *Let $\hat{H}_N = H_N(b)$. Then*

$$(\hat{H}_1, \hat{H}_2, \hat{H}_3, \hat{H}_4) = (\lambda, m, r, \eta)$$

and

$$\hat{H}_{k+4} \hat{H}_k = \alpha \hat{H}_{k+3} \hat{H}_{k+1} + \tau \hat{H}_{k+2}^2, \quad k \geq 1. \quad (10)$$

Proof. Equation (9) and (6) give

$$\hat{H}_1 = \lambda, \quad \hat{H}_2 = q^2 m' = m, \quad \hat{H}_3 = q^6 r' = r, \quad \hat{H}_4 = q^{12} \eta' = \eta.$$

Write $H'_N = H_N(a)$. The primed sequence satisfies

$$H'_{k+4} H'_k = H'_{k+3} H'_{k+1} + \tau' H'_{k+2}^2.$$

The weight difference between the left-hand product and the first product on the right is 6, while that between the left-hand product and H'_{k+2}^2 is 8. Hence

$$\hat{H}_{k+4} \hat{H}_k = q^6 \hat{H}_{k+3} \hat{H}_{k+1} + q^8 \tau' \hat{H}_{k+2}^2.$$

Now use $q^6 = \alpha$ and $q^8 \tau' = \tau$. $\square$

Corollary 3 (Low-order recurrence check). *The first two values forced beyond the initial tuple are*

$$\hat{H}_5 = \frac{\alpha \eta m + \tau r^2}{\lambda}, \quad \hat{H}_6 = \frac{\alpha \hat{H}_5 r + \tau \eta^2}{m}.$$

These formulas provide a direct indexing check for any coefficient expansion of (7).

3 Why fractional powers appear

The weighted transport solves the ordinary Hankel-determinant problem, but it does not preserve the companion normalization in its original form. This is the point at which the stronger rationality question enters.

3.1 The transported companion identity

Proposition 4. *If the primed series obeys $D_N(a) = H'_{N+2}$, then*

$$\widehat{D}_N := D_N(b) = q^{-3N-2} \widehat{H}_{N+2}, \quad N \geq 1. \quad (11)$$

Proof. By (9),

$$D_N(b) = q^{N^2} H'_{N+2}, \quad \widehat{H}_{N+2} = q^{(N+2)(N+1)} H'_{N+2}.$$

Their exponent difference is $N^2 - (N+2)(N+1) = -3N - 2$. $\square$

Remark 5. A constant coefficient rescaling $b_k \mapsto cb_k$ changes the ratio $D_N(b)/\widehat{H}_{N+2}$ by c^{-2} . It cannot remove the factor q^{-3N} . Thus the literal identity $D_N = H_{N+2}$ is not invariant under this graded transport. Equation (11), rather than the ungraded identity, is the companion normalization naturally inherited from the five-parameter series.

3.2 The first coefficients and failure of descent

For the five-parameter selected series, the companion normalization gives $g_0 = \lambda$ and $g_1 = r$. Applied to the primed parameters in (6), this yields $a_0 = \lambda$ and $a_1 = r' = q^{-6}r$. Therefore (8) gives

$$b_0 = \lambda, \quad b_1 = q^{-5}r = \frac{qr}{\alpha}. \quad (12)$$

The next two coefficients obtained by direct expansion are

$$b_2 = \frac{m}{\lambda} + \frac{q^{-10}r^2}{\lambda} = \frac{m}{\lambda} + \frac{q^2r^2}{\alpha^2\lambda}, \quad (13)$$

$$b_3 = \frac{q^5m^2}{\lambda^2r} + \frac{q^3\eta}{\alpha r} + \frac{2qmr}{\alpha\lambda^2} + \frac{q^3r^3}{\alpha^3\lambda^2}. \quad (14)$$

Hence

$$\begin{aligned} G_\alpha(x) &= \lambda + \frac{qr}{\alpha}x + \left(\frac{m}{\lambda} + \frac{q^2r^2}{\alpha^2\lambda} \right) x^2 \\ &\quad + \left(\frac{q^5m^2}{\lambda^2r} + \frac{q^3\eta}{\alpha r} + \frac{2qmr}{\alpha\lambda^2} + \frac{q^3r^3}{\alpha^3\lambda^2} \right) x^3 + O(x^4). \end{aligned} \quad (15)$$

The first coefficient also follows from (11) at $N = 1$:

$$b_1 = D_1(b) = q^{-5}H_3(b) = q^{-5}r.$$

Thus the fractional power is forced by the transported companion data rather than caused by an insufficient simplification.

Put

$$K = \mathbb{Q}(\lambda, m, r, \eta, \alpha, \tau), \quad L = K(q), \quad q^6 = \alpha.$$

Proposition 6 (Non-descent of the transported branch). *Regard α as an indeterminate. In the generic nondegenerate case $r \neq 0$, the transported series (7) does not belong to $K[[x]]$.*

Proof. Its coefficient of x is $b_1 = q^{-5}r = qr/\alpha$, which is not in K for generic α . Equivalently, if $\sigma(q) = \zeta q$ with $\zeta^6 = 1$ and $\zeta \neq 1$, then $\sigma(b_1) = \zeta^{-5}b_1 = \zeta b_1 \neq b_1$. $\square$

Although the coefficients do not descend, the fractional powers cancel in the ordinary Hankel determinants. For example,

$$H_2(b) = b_0 b_2 - b_1^2 = \lambda \left(\frac{m}{\lambda} + \frac{q^{-10} r^2}{\lambda} \right) - q^{-10} r^2 = m.$$

Together with the theorem above, this establishes the precise distinction: ordinary Hankel descent holds, whereas descent of the selected transported series fails.

4 What is defined over the six-parameter field

Although the selected transported series does not descend, this does not imply that the underlying recurrence or its curve requires fractional parameters. We now separate these levels.

4.1 Transport of the discriminant

Separate the old invariant into

$$A = \frac{\lambda \eta}{mr} + \frac{\tau mr}{\lambda \eta}, \quad B = \frac{m^2}{\lambda r} + \frac{r^2}{m \eta}. \quad (16)$$

After substitution of (6),

$$J' = q^{-4} A + q^2 B. \quad (17)$$

Consequently the substitution $x \mapsto qx$ gives

$$qJ' = q^{-3} A + q^3 B = \alpha^{-1/2} A + \alpha^{1/2} B. \quad (18)$$

Proposition 7 (Transported discriminant). *The discriminant obtained by substituting (6) and $x \mapsto qx$ into (4) is*

$$\Delta_\alpha(x) = q^{-50} (x \lambda^2 \eta m r^2)^2 \left[\left(1 - (q^{-3} A + q^3 B) x + q^{-6} \tau x^2 \right)^2 - 4 q^3 x^3 \right]. \quad (19)$$

Equivalently,

$$\Delta_\alpha(x) = \alpha^{-25/3} (x \lambda^2 \eta m r^2)^2 \left[\left(1 - (\alpha^{-1/2} A + \alpha^{1/2} B) x + \frac{\tau}{\alpha} x^2 \right)^2 - 4 \alpha^{1/2} x^3 \right]. \quad (20)$$

Proof. The square prefactor has total q -weight $2(1 - 12 - 2 - 12) = -50$. The three terms in the residual factor transform as $J'x \mapsto qJ'x$, $\tau'x^2 \mapsto q^2\tau'x^2 = q^{-6}\tau x^2$, and $x^3 \mapsto q^3x^3$. Equations (17) and $q^6 = \alpha$ give the stated forms. $\square$

The overall monomial factor in (19) is a square in $\mathbb{K}(x)$ and does not affect the residual double cover. It should, however, be retained when comparing discriminants of particular polynomial representatives, because multiplying a quadratic equation by a scalar squares that scalar in its discriminant.

4.2 A rational model of the curve

Remove the square prefactor in (20) and put

$$x = \alpha^{1/2} z. \quad (21)$$

The residual equation becomes

$$Y^2 = \left(1 - (A + \alpha B) z + \tau z^2 \right)^2 - 4 \alpha^2 z^3. \quad (22)$$

All coefficients of (22) belong to $\mathbb{Q}(\lambda, m, r, \eta, \alpha, \tau)$.

For comparison with the quartic convention used in [2], set $X = z^{-1}$ and $Y = z^{-2}y$. Then (22) is birational to

$$y^2 = \left(X^2 - (A + \alpha B)X + \tau\right)^2 - 4\alpha^2 X. \quad (23)$$

Thus the curve descends to the rational six-parameter field. The coordinate change (21) nevertheless rescales coefficients of a formal power series and does not preserve the normalized Hankel data. Curve descent and descent of the distinguished generating series are therefore different problems.

4.3 Formal representatives with the same Hankel determinants

Preservation of the ordinary Hankel transform alone does not characterize the distinguished algebraic series. For a nondegenerate Hankel sequence define

$$\beta_n = \frac{H_{n-1}H_{n+1}}{H_n^2}, \quad H_0 = 1. \quad (24)$$

All H_n generated by (10), and hence all β_n , lie in K . For arbitrary $a_0, a_1, \dots \in K$, the Jacobi fraction

$$\tilde{G}(x) = \frac{H_1}{1 - a_0x - \frac{\beta_1 x^2}{1 - a_1x - \frac{\beta_2 x^2}{1 - a_2x - \ddots}}} \quad (25)$$

has ordinary Hankel determinants H_n . Thus radical-free formal-series representatives over K exist, but they need not be quadratic algebraic, preserve the graded companion identity, or arise from the distinguished Stieltjes splitting.

5 Comparison with the construction of Chang and Liu

The preceding results leave open whether a different quadratic-algebraic generating function can avoid fractional powers. The construction of Chang and Liu is directly relevant because it realizes a generic bilateral Somos–4 orbit by Hankel determinants. We therefore compare the two descriptions explicitly before considering further repairs.

5.1 The shifted Somos–4 orbit

Chang and Liu [1, Sections 3.3–4.2] treat the generic bilateral Somos–4 initial-value problem by Hankel determinants of quadratic orthogonal pairs. Their normalized recurrence is

$$\sigma_{n+2}\sigma_{n-2} = a\sigma_{n+1}\sigma_{n-1} + b\sigma_n^2, \quad n \in \mathbb{Z}, \quad (26)$$

with $\sigma_{-1} = x$, $\sigma_0 = 1$, $\sigma_1 = y$, and $\sigma_2 = z$. Our freely prescribed determinants fit this normalization after the shift

$$\sigma_n = \frac{H_{n+2}}{m}, \quad (x, y, z; a, b) = \left(\frac{\lambda}{m}, \frac{r}{m}, \frac{\eta}{m}; \alpha, \tau\right). \quad (27)$$

This uses all six parameters and avoids the constraint $\eta = \alpha\lambda r + \tau m^2$ that would follow from imposing our recurrence at $N = 0$ with $H_0 = 1$.

Write $u^2 = \alpha$ and define

$$\mathcal{N} = \tau r^2 m + \lambda^2 \eta^2 + \alpha \eta m^3 + \alpha \lambda r^3, \quad (28)$$

$$\mathcal{N}_0 = \tau r^2 m - \lambda^2 \eta^2 + \alpha \eta m^3 + \alpha \lambda r^3. \quad (29)$$

Under (27), the Chang–Liu parameters include

$$d_0 = \frac{\lambda r}{m^2}, \quad u_0 = \frac{2r}{m}, \quad (30)$$

$$\frac{v}{u} = \frac{\mathcal{N}}{2\alpha\lambda r\eta m}, \quad uv = \frac{\mathcal{N}}{2\lambda r\eta m}, \quad (31)$$

$$\frac{v_0}{u} = \frac{\mathcal{N}_0}{2\alpha\lambda r\eta m}, \quad uv_0 = \frac{\mathcal{N}_0}{2\lambda r\eta m}, \quad (32)$$

and

$$f = \frac{\tau}{\alpha} - \frac{\mathcal{N}^2}{4\alpha\lambda^2 r^2 \eta^2 m^2}. \quad (33)$$

The sign difference between $\mathcal{N}$ and $\mathcal{N}_0$ is essential because v_0 enters the distinguished moment branch. The scaling $X = uZ$ turns their curve into

$$Y^2 = (\alpha Z^2 + f)^2 + 4\alpha Z - 4uv, \quad (34)$$

which is defined over K . This is curve-level descent, not coefficientwise descent of a selected moment branch.

5.2 Translation to Jacobi data

Use the local Jacobi convention

$$F(t) = \frac{\lambda}{1 - A_0 t - \frac{B_1 t^2}{1 - A_1 t - \frac{B_2 t^2}{1 - A_2 t - \frac{B_3 t^2}{1 - A_3 t - B_4 t^2 M_4(t)}}}}, \quad M_4(0) = 1. \quad (35)$$

The product formula

$$H_N = \lambda^N \prod_{j=1}^{N-1} B_j^{N-j} \quad (36)$$

gives

$$B_1 = \frac{m}{\lambda^2}, \quad B_2 = \frac{\lambda r}{m^2}, \quad B_3 = \frac{m\eta}{r^2}, \quad B_4 = \frac{r(\alpha\eta m + \tau r^2)}{\lambda\eta^2}. \quad (37)$$

The companion identities through $D_3 = H_5$ force

$$A_0 = \frac{r}{\lambda}, \quad A_1 = \frac{\eta\lambda^2 + m^2}{\lambda m r}, \quad A_2 = \frac{\alpha\eta m^3 + \lambda r^3 + \tau m^2 r^2}{\eta\lambda m r}. \quad (38)$$

Proposition 8 (Fourth shifted determinant). *For the convention (35),*

$$D_4 = \lambda^4 B_1^3 B_2^2 B_3 (A_0 A_1 A_2 A_3 - A_0 A_1 B_3 - A_0 A_3 B_2 - A_2 A_3 B_1 + B_1 B_3). \quad (39)$$

After substitution of (37) and (38), this reduces to

$$D_4 = \frac{\eta}{\lambda r^2} [A_3 r(\alpha\eta m + \tau r^2) - \eta^2 \lambda]. \quad (40)$$

Consequently $D_4 = H_6$, where

$$H_6 = \frac{\alpha^2 \eta m r + \alpha \tau r^3 + \tau \eta^2 \lambda}{\lambda m},$$

uniquely forces

$$A_3 = \frac{\alpha^2 \eta m r^3 + \alpha \tau r^5 + \eta^3 \lambda m + \tau \eta^2 \lambda r^2}{\eta m r(\alpha\eta m + \tau r^2)}. \quad (41)$$

Proof. Equation (39) follows by direct expansion of the shifted 4×4 Hankel determinant in Jacobi parameters. Substitution and collection of the terms linear in A_3 gives (40). Solving (40) for A_3 after inserting the displayed value of H_6 gives (41). $\square$

An equivalent useful form is

$$A_3 = \frac{\alpha r^2}{\eta m} + \frac{\eta \lambda (\eta m + \tau r^2)}{m r (\alpha \eta m + \tau r^2)}.$$

No B_4 occurs in D_4 : the fourth ordinary Hankel step determines B_4 , while the fourth shifted step determines A_3 , without imposing any extra relation among the six parameters.

Chang and Liu use

$$d_n = \frac{\sigma_{n-1} \sigma_{n+1}}{\sigma_n^2}.$$

Substitution of (27) gives the exact all-order interface

$$d_n = \frac{H_{n+1} H_{n+3}}{H_{n+2}^2} = B_{n+2}. \quad (42)$$

Thus $d_0 = B_2$, $d_1 = B_3$, and $d_2 = B_4$. At the subdiagonal level only B_1 precedes the Chang–Liu tail; the diagonal and moment normalizations remain separate issues.

6 What the comparison resolves

The comparison confirms the same basic distinction found for the weighted transport: the orbit and the curve may descend even when one distinguished moment branch does not.

6.1 Even and odd parts of the Chang–Liu branch

Let

$$S(t; u) = \sum_{n \geq 0} s_n(u) t^n, \quad u^2 = \alpha,$$

be the distinguished positive Chang–Liu moment branch. The two choices of square root satisfy

$$S(t; -u) = S(-t; u), \quad s_n(-u) = (-1)^n s_n(u). \quad (43)$$

Hence the two branches have equal ordinary Hankel determinants, but a single branch is generically not fixed by $u \mapsto -u$ and therefore does not descend coefficientwise to $K[[t]]$. Writing

$$S(t; u) = E(t^2; u) + tO(t^2; u)$$

gives $E(z; u) \in K[[z]]$ and $O(z; u)/u \in K[[z]]$, whereas O is generically anti-invariant.

Proposition 9 (Parity-rescaling obstruction). *Let $r_{2k} = a s_{2k}$ and $r_{2k+1} = b s_{2k+1}$. If the generic identity $H_3(r) = \kappa H_3(s)$ holds coefficientwise, then $a^3 = \kappa$, $ab^2 = \kappa$, and therefore $b^2 = a^2$.*

Proof. Compare the coefficients of the independent monomials in the two generic 3×3 Hankel determinants. The pure even contribution has weight a^3 , while the terms containing two odd moments have weight ab^2 . $\square$

In particular, the descended recombination $R(t) = E(t^2; u) + tO(t^2; u)/u$ fails already at order two:

$$H_2(R) - H_2(S) = \left(1 - \frac{1}{\alpha}\right) s_1^2.$$

Proposition 10 (Geometric-scaling obstruction). *No generic scaling $r_n = cd^n s_n$ converts the normalized Chang–Liu determinants $H_N(s) = H_{N+2}/m$ into $H_N(r) = H_{N+2}$ for all N .*

Proof. Such a scaling gives $H_N(r) = c^N d^{N(N-1)} H_N(s)$, so one would need $c^N d^{N(N-1)} = m$. The cases $N = 1, 2$ force $c = m$ and $d^2 = m^{-1}$, while $N = 3$ then gives $c^3 d^6 = 1$, not m , generically. $\square$

Finally, the Galois-equivariant obstruction proved above applies to a finite Jacobi prefix over K : as a generically invertible linear-fractional transformation in its tail, it cannot turn one non-descending Chang–Liu branch into a descending one. This excludes the direct prefix strategy, but not every possible quadratic-algebraic representative.

6.2 Why base-field transforms cannot repair a branch

The following observation excludes a broad class of attempted rationalizations.

Proposition 11 (Galois-equivariant obstruction). *Let $T : L[[x]] \rightarrow L[[x]]$ be injective and commute with every K -automorphism of L . If $T(G) \in K[[x]]$, then $G \in K[[x]]$. If T is invertible, the two conditions are equivalent.*

Proof. For every $\sigma \in \text{Gal}(L/K)$, $T(\sigma G) = \sigma T(G) = T(G)$. Injectivity gives $\sigma G = G$, so G is fixed by the Galois action and belongs to $K[[x]]$. $\square$

For $c \in K$, the binomial transform

$$B_c(G)(x) = \frac{1}{1 - cx} G\left(\frac{x}{1 - cx}\right)$$

is defined over K , commutes with the Galois action, and is invertible with inverse B_{-c} . Likewise, the invert transform

$$I_c(G)(x) = \frac{G(x)}{1 - cxG(x)}$$

is defined over K , Galois-equivariant, and invertible with inverse I_{-c} . Consequently neither transform can turn the non-descending transported branch into a series in $K[[x]]$. This conclusion concerns these transforms over the base field; it does not exclude a new algebraic construction with different companion data.

7 Remaining possibilities

The established facts now separate three levels cleanly: the Chang–Liu curve descends; its ordinary Hankel sequence descends; an individual distinguished moment branch does not descend generically. The rational Jacobi calculation extends through A_3 and B_4 , and the subdiagonal interface is exactly $d_n = B_{n+2}$.

The remaining constructive possibility should therefore be sought in the invariant pair

$$E(z), \quad R(z) := O(z)/u,$$

rather than by attaching an injective K -rational prefix to one raw branch. Starting from the quadratic equation for $S(t; u)$, one should substitute

$$S(t; u) = E(t^2) + utR(t^2)$$

and separate invariant and anti-invariant parts. This yields coupled algebraic equations over K for E and R . A non-diagonal rational construction from this pair can then be tested against

$$f_0 = \lambda, \quad f_1 = r, \quad f_2 = \frac{r^2 + m}{\lambda}$$

and the determinant conditions through $D_4 = H_6$.

Thus no general nonexistence claim is made. What is excluded is the narrower class consisting of parity rescalings, constant mass and argument scalings, alternating signs, and injective finite prefixes over K applied to a single distinguished Chang–Liu branch. A constructive all-order theorem, or a precise nonexistence theorem for a broader class, remains open.

8 Conclusion

We began with the five-parameter generating function of [2] and asked whether it extends to the general two-coefficient Somos–4 recurrence. The weighted substitution (7) gives an exact extension for arbitrary $(\lambda, m, r, \eta, \alpha, \tau)$. It preserves the prescribed ordinary Hankel determinants and yields (10). Thus the six-parameter Hankel problem is solved at the level of the recurrence.

The same construction reveals a second and more delicate problem. Its companion determinants satisfy the graded identity (11), and the coefficient $b_1 = q^{-5}r$ shows that the selected branch does not generically descend to $K = \mathbb{Q}(\lambda, m, r, \eta, \alpha, \tau)$. This is not merely an unsimplified formula. On the other hand, the rational quartic (22) shows that the associated curve does descend. The Hankel sequence, the curve, and a distinguished generating-function branch must therefore be treated as separate objects.

The comparison with Chang and Liu makes this separation explicit in a second construction. The shifted dictionary (27) identifies their generic bilateral orbit with ours, and the all-order relation $d_n = B_{n+2}$ identifies the subdiagonal Jacobi data. The rational restoration through A_3 and B_4 shows that the prescribed companion data are consistent at the computed levels. Nevertheless, the parity relation (43) shows that an individual distinguished moment branch is generically non-descending.

The remaining question is consequently narrower than the original one. It is unknown whether another quadratic-algebraic representative can simultaneously have coefficients in K , realize the prescribed six-parameter Hankel orbit, and satisfy the ungraded identity $D_N = H_{N+2}$. Parity rescaling, geometric scaling, and injective finite prefixes over K cannot achieve all three goals. These obstructions do not constitute a general nonexistence theorem. A non-diagonal construction based on the invariant pair $(E, O/u)$ remains a concrete possibility for further investigation.

Disclosure on authoring and verification tools

The manuscript was revised with assistance from AI-based language and symbolic calculation tools. Their use was directed toward notation control, LaTeX editing, and exact algebraic checks. The author remains responsible for the mathematical statements and the final text.

References

- [1] X.-K. Chang and J. Liu, Hankel determinants from quadratic orthogonal pairs for hyperelliptic functions and their applications, arXiv:2603.11670v4 [nlin.SI], 2026; to appear in the *Journal of the London Mathematical Society*.
- [2] T. Scheuerle, Direct generation of a Somos–4 sequence from an algebraic generating function, arXiv:2609.15754 [math.CO], 2026.
- [3] H. S. Wall, *Analytic Theory of Continued Fractions*, Van Nostrand, New York, 1948.
- [4] P. Flajolet, Combinatorial aspects of continued fractions, *Discrete Mathematics* **32** (1980), 125–161.

- [5] J. W. Layman, The Hankel transform and some of its properties, *Journal of Integer Sequences* **4** (2001), Article 01.1.5.
- [6] C. French, Transformations preserving the Hankel transform, *Journal of Integer Sequences* **10** (2007), Article 07.7.3.
- [7] A. N. W. Hone, Elliptic curves and quadratic recurrence sequences, *Bulletin of the London Mathematical Society* **37** (2005), 161–171.
- [8] S. Fomin and A. Zelevinsky, The Laurent phenomenon, *Advances in Applied Mathematics* **28** (2002), 119–144.
- [9] A. J. van der Poorten and C. S. Swart, Recurrence relations for elliptic sequences: every Somos–4 is a Somos– k , *Bulletin of the London Mathematical Society* **38** (2006), 546–554.