

Australian, Aboriginal & Torres Strait Islander Nutrition & Physical Activity data

A/Prof Kim Bell-Anderson



THE UNIVERSITY OF
SYDNEY



We recognise and pay respect to the Elders and communities – past, present, and emerging – of the lands that the University of Sydney's campuses stand on. For thousands of years they have shared and exchanged knowledges across innumerable generations for the benefit of all.

Ngyni ngalawangun mari budjari Gadinurada



THE UNIVERSITY OF
SYDNEY




Australian Health Survey 2011-13

- The largest and most comprehensive health surveys conducted in Australia
- It combines:
 - National Health Survey (NHS)
 - National Aboriginal and Torres Strait Islander Health Survey
 - National Nutrition and Physical Activity Survey (NNPAS)
 - National Health Measures Survey (NHMS)

[Australian Health Survey: Users' Guide, 2011-13](#)

STRUCTURE OF THE 2011-13 AUSTRALIAN HEALTH SURVEY



APPROACHED SAMPLE, FINAL SAMPLE AND RESPONSE RATES

	New South Wales	Victoria	Queensland	South Australia	Western Australia	Tasmania	Northern Territory	Australian Capital Territory	Australia
Households approached (after sample loss)	3 269	3 051	2 907	2 285	2 385	1 635	1 307	1 516	18 355
Households in sample	2 736	2 516	2 457	1 962	2 144	1 469	975	1 306	15 565
Response rate (%)	83.7	82.5	84.5	85.9	89.9	89.8	74.6	86.1	84.8
Persons in sample	3 602	3 287	3 244	2 508	2 847	1 909	1 304	1 725	20 426

ABS record identifiers

Variable	Description
ABSHID/ABSLID	Household identifier
ABSPID	Person identifier within each household
ABSH/LID + ABSPID	Unique 14 digit random identifier
ABSFID	Food records within each person record
ABSPID + ABSFID	Uniquely identifies each food record
ABSSID	Supplement records within each person record
ABSPID + ABSSID	Uniquely identifies each supplement record
ABSLFID	ADG summary records within each person record
ABSPID + ABSLFID	Uniquely identifies each ADG summary record

Data source and definitions

- Usual daily intake of fruit and vegetables in the AHS is based on self-reported data for the number of serves of fruit and vegetables that people usually ate each day.
- A **serve of vegetables** was defined as:
 - half a cup of cooked vegetables
 - one medium potato or
 - one cup of salad vegetables
 - (approximately 75 grams).
- A **serve of fruit** was defined as:
 - one medium piece or two small pieces of fresh fruit,
 - one cup of diced fruit,
 - a quarter of a cup of sultanas, or four dried apricot halves
 - (approximately 150 grams of fresh fruit or 50 grams of dried fruit).
- Tomatoes were included as a vegetable rather than a fruit, and legumes were excluded.
- Fruit juices were not considered to be fruit.

WAIST MEASUREMENT GUIDELINES, Adults

	Men	Women
Not at risk	Waist circumference less than 94 cm	Waist circumference less than 80 cm
Increased risk	Waist circumference more than or equal to 94 cm	Waist circumference more than or equal to 80 cm
Greatly increased risk	Waist circumference more than or equal to 102 cm	Waist circumference more than or equal to 88 cm

Exercise

- Types of exercise covered in the AHS were walking, moderate and vigorous exercise for sport, recreation or fitness.
- **Moderate exercise** consists of activity undertaken for fitness, recreation or sport that causes a moderate increase in heart rate or breathing
- **Vigorous exercise** causes a large increase in a person's heart rate or breathing.
- Level of exercise is determined based on the frequency, intensity and duration of exercise.

Index of Relative Socio-Economic Disadvantage

This is one of four Socio-Economic Indexes for Areas (SEIFA) compiled by ABS following each Census of Population and Housing. The indexes are compiled from various characteristics of persons resident in particular areas: the Index of Relative Socio-Economic Disadvantage summarises attributes such as low income, low educational attainment, high unemployment and jobs in relatively unskilled occupations. A lower Index of Relative Socio-Economic Disadvantage quintile (e.g. the first quintile) indicates relatively greater disadvantage and a lack of advantage in general. A higher Index of Relative Socio-Economic Disadvantage (e.g. the fifth quintile) indicates a relative lack of disadvantage and greater advantage in general. For further information about SEIFA see the Australian Health Survey: Users' Guide, 2011-13 (cat. no. 4363.0.55.001) scheduled for release in November 2012.

CPC Publications

- Sui, Z., Raubenheimer, D., & Rangan, A. (2017). **Exploratory analysis of meal composition in Australia: meat and accompanying foods.** *Public Health Nutrition*, 20(12), 2157–2165.
- Sui, Z., Raubenheimer, D., & Rangan, A. (2017). **Consumption patterns of meat, poultry, and fish after disaggregation of mixed dishes: secondary analysis of the Australian National Nutrition and Physical Activity Survey 2011–12,** 1–12.
- Brand-Miller, J. C., & Barclay, A. W. (2017). **Declining consumption of added sugars and sugar-sweetened beverages in Australia: a challenge for obesity prevention.** *The American Journal of Clinical Nutrition*
- Engelen, L., Gale, J., Chau, J. Y., Hardy, L. L., Mackey, M., Johnson, N., et al. (2017). **Who is at risk of chronic disease? Associations between risk profiles of physical activity, sitting and cardio-metabolic disease in Australian adults.** *Australian and New Zealand Journal of Public Health*, 41(2), 178–183.

USER NOTE:

Delete the social media icons
below that you do not require and
align the icons you need neatly

Insert team/discipline area

Name Surname Title

+61 2 1234 5678

[sydney.edu.au/xxxx](#)

