

# Annals of Surgery

## Laparoscopic Hartmann's procedure: is it a good option for acute colonic diverticulitis with generalized peritonitis? A national multicenter comparative matched cohort study --Manuscript Draft--

|                                                                                                                                            |                                                                                                                                                                                    |
|--------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Manuscript Number:</b>                                                                                                                  |                                                                                                                                                                                    |
| <b>Full Title:</b>                                                                                                                         | Laparoscopic Hartmann's procedure: is it a good option for acute colonic diverticulitis with generalized peritonitis? A national multicenter comparative matched cohort study      |
| <b>Article Type:</b>                                                                                                                       | Original Study                                                                                                                                                                     |
| <b>Keywords:</b>                                                                                                                           | diverticulitis; Laparoscopy; Perforated diverticulitis; Sigmoid resection, Hartmann Procedure, Propensity score.                                                                   |
| <b>Corresponding Author:</b>                                                                                                               | Thomas Husson, M.D.<br>Hopital Europeen Georges Pompidou<br>Paris, FR FRANCE                                                                                                       |
| <b>Corresponding Author Secondary Information:</b>                                                                                         |                                                                                                                                                                                    |
| <b>Corresponding Author's Institution:</b>                                                                                                 | Hopital Europeen Georges Pompidou                                                                                                                                                  |
| <b>Corresponding Author's Secondary Institution:</b>                                                                                       |                                                                                                                                                                                    |
| <b>First Author:</b>                                                                                                                       | Thomas Husson, M.D.                                                                                                                                                                |
| <b>First Author Secondary Information:</b>                                                                                                 |                                                                                                                                                                                    |
| <b>Order of Authors:</b>                                                                                                                   | Thomas Husson, M.D.                                                                                                                                                                |
|                                                                                                                                            | Charles Sabbagh, M.D., PhD                                                                                                                                                         |
|                                                                                                                                            | Laura Beyer-Berjot, M.D., PhD                                                                                                                                                      |
|                                                                                                                                            | Mehdi Ouaiissi, M.D., PhD                                                                                                                                                          |
|                                                                                                                                            | Valérie Bridoux, M.D., PhD                                                                                                                                                         |
|                                                                                                                                            | Mehdi Karoui, M.D., PhD                                                                                                                                                            |
|                                                                                                                                            | Arnaud Alves, M.D., PhD                                                                                                                                                            |
|                                                                                                                                            | Antoine Mariani, M.D., PhD                                                                                                                                                         |
|                                                                                                                                            | French Surgical Association                                                                                                                                                        |
| <b>Order of Authors Secondary Information:</b>                                                                                             |                                                                                                                                                                                    |
| <b>Manuscript Region of Origin:</b>                                                                                                        | FRANCE                                                                                                                                                                             |
| <b>Suggested Reviewers:</b>                                                                                                                | Guillaume Meurette, M.D., PhD<br>Agrégé, HUG: Hopitaux Universitaires Geneve<br>guillaume.meurette@hug.ch<br>Affiliation with the topic, member of European Coloproctology Society |
| <b>Opposed Reviewers:</b>                                                                                                                  |                                                                                                                                                                                    |
| <b>Additional Information:</b>                                                                                                             |                                                                                                                                                                                    |
| <b>Question</b>                                                                                                                            | <b>Response</b>                                                                                                                                                                    |
| Does your manuscript describe a Randomized Controlled Trial?                                                                               | No                                                                                                                                                                                 |
| Briefly indicate the type of study your paper describes.<br>as follow-up to "Does your manuscript describe a Randomized Controlled Trial?" | Retrospective cohort study with propensity score matching                                                                                                                          |

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                         |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------|
| Have all coauthors viewed and approved of this version manuscript?                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Yes - All coauthors have seen & approved                |
| Do you or any of your coauthors have any potential conflicts of interest you need to declare?                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | The authors declare there are no conflicts of interest. |
| Was this submission funded by an outside agency (e.g., business, government agency, etc.)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | No funding was received in support of this work.        |
| If your submission contains figures, are any of them in color?                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Yes, my submission has colored figures.                 |
| Are you and your coauthors willing to pay for color printing?<br>as follow-up to "If your submission contains figures, are any of them in color?"                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | Yes, we will pay for color printing.                    |
| <p>RETAINED RIGHTS: Except for copyright, other proprietary rights related to the Work (e.g., patent or other rights to any process or procedure) shall be retained by the author. To reproduce any text, figures, tables, or illustrations from this Work in future works of their own, the author must obtain written permission from Wolters Kluwer Health, Inc. ("WKH").</p> <p>ORIGINALITY: Each author warrants that his or her submission to the Work is original, does not infringe upon, violate, or misappropriate any copyright or other intellectual property rights, or any other proprietary right, contract or other right or interest of any third party, and that he or she has full power to enter into this agreement. Neither this Work nor a similar work has been published nor shall be submitted for publication elsewhere while under consideration by this Publication.</p> <p>AUTHORSHIP RESPONSIBILITY: Each author warrants that he or she has participated sufficiently in the intellectual content, the analysis of data, if applicable, and the writing of the Work to</p> | I agree                                                 |

take public responsibility for it. Each has reviewed the final version of the Work, believes it represents valid work, and approves it for publication. Moreover, should the editors of the Publication request the data upon which the work is based, they shall produce it.

**PREPRINTS:** Upon acceptance of the article for publication, each author warrants that he/she will promptly remove any prior versions of this Work (normally a preprint) that may have been posted to an electronic server.

**DISCLAIMER:** Each author warrants that this Work contains no libelous or unlawful statements and does not infringe or violate the publicity or privacy rights of any third party, libel or slander any third party, contain any scandalous, obscene, or negligently prepared information, or infringe or violate any other personal or proprietary right of others. Each author warrants that the Work does not contain any fraudulent, plagiarized or incorrectly attributed material. Each author warrants that all statements contained in the Work purporting to be facts are true, and any formula or instruction contained in the Work will not, if followed accurately, cause any injury, illness, or damage to the user. If excerpts (e.g., text, figures, tables, illustrations, or audio/video files) from copyrighted works are included, a written release will be secured by the author prior to submission, and credit to the original publication will be properly acknowledged. Each author further warrants that he or she has obtained, prior to submission, written releases from patients whose names or likenesses are submitted as part of the Work. Should the Editor or WKH request copies of such written releases, the author shall provide them in a timely manner.

TRADE COMPLIANCE: Each author warrants that if the author, any of the author's coauthors or any other individual whose content is included in the Work resides in Iran, Cuba, Syria, North Korea, Crimea, the Donetsk People's Republic (DNR) or the Luhansk People's Republic (LNR) regions of the Ukraine, the Work has been prepared in a personal, academic, or research capacity and not as an official representative or otherwise on behalf of the relevant government.

#### DISCLOSURES/CONFLICT OF INTEREST

Each author must identify any financial interests or affiliations with institutions, organizations, or companies relevant to the manuscript by completing the form below. Additionally, any financial associations involving a spouse, partner or children must be disclosed as well.

Note: Some sections below come from the ICMJE Uniform Disclosure Form for Potential Conflicts of Interest at [http://www.icmje.org/downloads/coi\\_disclosure.pdf](http://www.icmje.org/downloads/coi_disclosure.pdf) (dated July 2010).

Did you or your institution at any time receive payment or support in kind for any aspect of the submitted work (including but not limited to grants, consulting fee or honorarium, support for travel to meetings for the study or other purposes, fees for participation in review activities such as data monitoring boards, statistical analysis, end point committees, and the like, payment for writing or reviewing the manuscript, provision of writing assistance, medicines, equipment, or administrative support, etc...)?

No

Please indicate whether you have financial relationships (regardless of amount of compensation) with entities.

No

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                                                    |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|
| <p>You should report relationships that were present during the 36 months prior to submission including board membership, consultancy, employment, expert testimony, grants/grants pending, payment for lectures including service on speakers bureaus, payment for manuscript preparation, patents (planned, pending or issued), royalties, payment for development of educational presentations, stock/stock options, travel/accommodations/meeting expenses unrelated to activities listed (for example, if you report a consultancy above there is no need to report travel related to that consultancy), etc.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                    |
| <p>Other Relationships</p> <p>Are there other relationships or activities that readers could perceive to have influenced, or that give the appearance of potentially influencing, what you wrote in the submitted work?</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | <p>No other relationships/conditions/circumstances that present potential conflict of interest</p> |
| <p>AUTHOR'S OWN WORK: In consideration of WKH's publication of the Work, the author hereby transfers, assigns, and otherwise conveys all his/her copyright ownership worldwide, in all languages, and in all forms of media now or hereafter known, including electronic media such as CD-ROM, Internet, and Intranet, to WKH. If WKH should decide for any reason not to publish the Work, WKH shall give prompt notice of its decision to the corresponding author, this agreement shall terminate, and neither the author nor WKH shall be under any further liability or obligation. Each author grants WKH the rights to use his or her name and biographical data (including professional affiliation) in the Work and in its or the journal's promotion. Notwithstanding the foregoing, this paragraph shall not apply, and any transfer made pursuant to this paragraph shall be null and void if (i) the Work has been accepted by WKH for publication, and (ii) the author chooses to have the Work published by WKH as an open access publication.</p> | <p>I agree</p>                                                                                     |

WORK MADE FOR HIRE: If this Work or any element thereof has been commissioned by another person or organization, or if it has been written as part of the duties of an employee, an authorized representative of the commissioning organization or employer must also sign this form stating his or her title in the organization.

GOVERNMENT EMPLOYEES: If the Work or a portion of it has been created in the course of any author's employment by the United States Government, check the "Government" box at the end of this form. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" and is not subject to copyright. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

INSTITUTIONAL REVIEW BOARD/ANIMAL CARE COMMITTEE APPROVAL: Each author warrants that his or her institution has approved the protocol for any investigation involving humans or animals and that all experimentation was conducted in conformity with ethical and humane principles of research.

WARRANTIES: Each author warranty made in this form is for the benefit of WKH and the Editor; each author agrees to defend, indemnify, and hold harmless those parties for any breach of such warranties.

Please select:

Author's Own Work

Any additional comments?

I am the person in question for this submission or otherwise have approval to complete this agreement.

I agree

CME/CE Disclosure

I agree

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                              |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|
| <p>Each author must identify and disclose any financial associations involving a spouse, partner or children by completing the Family Disclosure question below, and whether any off-label uses or unapproved drugs or devices are discussed in his/her manuscript by completing the Off-Label Use/Unapproved Drugs or Products question below. In the event that the Work is published as a continuing education or continuing medical education article, this information will be provided to the accrediting body and may be included in the published article. When applicable, articles accepted for publication may need to comply with additional standards related to CME or CE accreditation. Please refer to guidelines for authors for details.</p> <p>WKH and its affiliates reserve the right to publish the manuscript as a continuing education article.</p> |                                                                                              |
| <p>Off-Label Use/Unapproved Drugs or Products</p> <p>If your manuscript discusses an unlabeled use of a commercial product or device or an investigational use of a product or device not yet approved by the FDA for any purpose, you must specifically disclose in the manuscript that the product is not labeled for the use under discussion or that the product is still investigational. Please check the item below that applies to you</p>                                                                                                                                                                                                                                                                                                                                                                                                                          | <p>I will not discuss unlabeled/investigational uses of any commercial product or device</p> |

## Cover Letter

Corresponding Author: Dr Antoine Mariani

Department of oncologic and digestive surgery, Hôpital Européen Georges Pompidou

Address: [20 Rue Leblanc, 75015 Paris](#), France

E-mail: [antoine.mariani@aphp.fr](mailto:antoine.mariani@aphp.fr)

Phone : +33 1 56 09 35 62

May 5<sup>th</sup>, 2026

Professor Justin B. Dimick,

Dear Editor-in-Chief of Annals of Surgery

Please find enclosed our manuscript entitled: "Laparoscopic Hartmann's procedure: is it a good option for acute colonic diverticulitis with generalized peritonitis? A national multicenter comparative matched cohort study" for consideration as an original investigation.

This manuscript is an original contribution; all authors have read and complied with author guidelines. Authors, status and affiliations are at the end of this letter.

Acute perforated sigmoid diverticulitis with generalized peritonitis remains a challenging surgical emergency. Although recent trials and current guidelines favor resection with primary anastomosis in stable patients, Hartmann's procedure is still used in more complex or high-risk situations. At the same time, minimally invasive colorectal surgery has progressively expanded to emergency settings, and laparoscopic Hartmann's procedure has been proposed as an attractive strategy. In theory, laparoscopy could reduce postoperative morbidity, and limit adhesion formation, thereby facilitating subsequent restoration of bowel continuity. This minimally invasive two-stage strategy could therefore appear to combine the immediate safety of a non-restorative procedure with improved long-term functional outcomes. However, whether these expected benefits are achieved in generalized diverticular peritonitis remains uncertain.

The aim of this multicenter national matched cohort study was to compare laparoscopic Hartmann's procedure with open sigmoidectomy and primary anastomosis with diverting ileostomy, with particular attention to postoperative outcomes and 12-month stoma-free survival. Our results indicate that laparoscopic Hartmann's procedure is feasible in this setting, but the anticipated advantages of the minimally invasive approach were not demonstrated.



Short-term outcomes were comparable between strategies, with no significant reduction in postoperative morbidity, major complications, mortality, or length of stay. In addition, conversion was frequent. Most importantly, laparoscopic Hartmann's procedure was associated with fewer stoma reversals and poorer 1-year stoma-free survival than open primary anastomosis with diversion.

This manuscript has not been previously published or submitted elsewhere for publication and will not be sent to another journal until a decision is made concerning publication by Annals of Surgery. Possible conflicts of interest are described in the manuscript file in the section "Conflict of Interest/Disclosure Statement". We have no financial ties to disclose. All authors have seen and approved the final version of the manuscript being submitted and that all authors fulfill the COPE (Committee on Publication Ethics) requirements for authorship.

Thank you for receiving our manuscript and considering it for publication. We hope that you will find it interesting and suitable for publication in Annals of Surgery  
Sincerely yours,

Antoine Mariani, M.D., PhD

**For all the authors:**

Thomas Husson, M.D.<sup>1</sup>; Charles Sabbagh, M.D., PhD<sup>2</sup>; Laura Beyer-Berjot, M.D., PhD<sup>3</sup>;  
Mehdi Ouaiissi M.D., PhD<sup>4</sup>; Valérie Bridoux M.D., PhD<sup>5</sup>; Karoui, M.D., PhD<sup>1</sup>; Arnaud  
Alves, M.D, PhD<sup>6</sup>; Antoine Mariani, M.D., PhD<sup>1</sup> **and the French Surgical Association**

**Cooperative Group**

1. Department of Surgery, European Georges Pompidou Hospital, Assistance Publique Hôpitaux de Paris, Paris, France
2. Department of Digestive Surgery, University of Caen Normandy, Caen, France
3. Department of Digestive Surgery, North University Hospital, Marseille, Assistance Publique Hôpitaux de Marseille, France
4. Department of Digestive, Oncological, Endocrine, Hepatobiliary and Liver Transplant, Trousseau Hospital, University Hospital of Tours, Tours, France

5. Department of Digestive Surgery, University Hospital of Rouen, Rouen, France
6. Department of Digestive Surgery, Amiens University Hospital, Amiens, France

## **Mini-Abstract**

In this French national propensity-matched cohort study of generalized diverticular peritonitis, laparoscopic Hartmann's procedure and open primary anastomosis with diverting ileostomy yielded similar short-term outcomes. However, laparoscopic Hartmann's procedure was associated with fewer stoma reversals and poorer 1-year stoma-free survival.

## **Abstract**

### **Objective**

Compare laparoscopic Hartmann's procedure (LHP) with open sigmoidectomy with primary anastomosis and diverting ileostomy (OPA).

### **Summary Background Data**

Laparoscopic surgery is increasingly used in emergency settings, but its role in generalized diverticular peritonitis remains debated.

### **Methods**

Retrospective study using a French multicenter cohort, including 7,053 patients who underwent surgery for colonic diverticulitis. Patients were included if they had a generalized peritonitis (Hinchey III or IV) treated in emergency by laparoscopic Hartmann or open anastomosis with ileostomy. Perioperative outcomes and stoma-free survival were compared after propensity score matching.

### **Results**

284 patients met inclusion criteria (138 LHP, 146 OPA). After matching, 101 patients were included in each group. Baseline characteristics were comparable. 30% of the overall population had Hinchey IV diverticulitis. Conversion was required in 49% of LHP cases. LHP did not confer significant benefits concerning postoperative outcomes including 90-day morbidity (53% vs 50%,  $p = 0.8$ ), major complications (27% vs 26%,  $p > 0.9$ ), 90-day mortality (5% vs 3%,  $p = 0.7$ ), or length of hospital stay (12 vs 10 days,  $p = 0.095$ ), and was associated with fewer stoma reversal (70% vs 88%,  $p = 0.002$ ). Stoma-free survival at 1 year was significantly better in the OPA group (85% vs 65%, respectively; log-rank  $p < 0.0001$ ; Cox model HR = 1.84 [1.39–2.44],  $p < 0.001$ ). No treatment-by-Hinchey interaction was observed ( $p = 0.269$ ).

## **Conclusion**

Although early outcomes were comparable, the minimally invasive Hartmann's procedure was associated with fewer stoma reversals and poorer stoma-free survival.

When feasible, open primary anastomosis should remain the preferred approach.

## **MeSH Keywords :**

Diverticulitis; Laparoscopy; Perforated diverticulitis; Sigmoid resection, Hartmann Procedure, Propensity score.

***Laparoscopic Hartmann's procedure: is it a good option for acute  
colonic diverticulitis with generalized peritonitis? A national multicenter  
comparative matched cohort study***

**Table of Contents**

|                                                         |                                            |
|---------------------------------------------------------|--------------------------------------------|
| <b><i>Abstract.....</i></b>                             | <b><i>Error! Bookmark not defined.</i></b> |
| <b><i>MeSH Keywords:.....</i></b>                       | <b><i>Error! Bookmark not defined.</i></b> |
| <b><i>Text of the Paper .....</i></b>                   | <b><i>6</i></b>                            |
| <b><i>Acknowledgements.....</i></b>                     | <b><i>19</i></b>                           |
| <b><i>References .....</i></b>                          | <b><i>24</i></b>                           |
| <b><i>Legend for Illustrations .....</i></b>            | <b><i>29</i></b>                           |
| <b><i>List of Supplemental Digital Content.....</i></b> | <b><i>30</i></b>                           |

## Title Page

# Laparoscopic Hartmann's procedure: is it a good option for acute colonic diverticulitis with generalized peritonitis? A national multicenter comparative matched cohort study

## Authors

Thomas Husson, M.D<sup>1</sup>, Arnaud Alves, M.D, PhD <sup>2</sup>, Laura Beyer-Berjot<sup>3</sup>, M.D., PhD, M  
Ouaisi<sup>4</sup>, M.D., PhD, V Bridoux<sup>5</sup>, M.D., PhD, Mehdi Karoui, M.D., PhD<sup>1</sup>; Charles  
Sabbagh M.D., PhD<sup>6</sup>, Antoine Mariani, M.D., PhD<sup>1</sup> and the French Surgical

## Association Cooperative Group

1. Department of Surgery, European Georges Pompidou Hospital, Assistance Publique  
Hôpitaux de Paris, Paris, France
2. Department of Digestive Surgery, University of Caen Normandy, Caen, France
3. Department of Digestive Surgery, North University Hospital, Marseille, Assistance  
Publique Hôpitaux de Marseille, France
4. Department of Digestive, Oncological, Endocrine, Hepatobiliary and Liver  
Transplant, Trousseau Hospital, University Hospital of Tours, Tours, France
5. Department of Digestive Surgery, University Hospital of Rouen, Rouen, France
6. Department of Digestive Surgery, Amiens University Hospital, Amiens, France

Corresponding author: Antoine Mariani, M.D, Ph.D.

Department of digestive surgery and surgical oncology

Hôpital Européen Georges Pompidou, Assistance Publique des Hôpitaux de Paris

Phone : +33 1 56 09 35 62

E-mail : antoine.mariani@aphp.fr

### ***Conference presentation***

The results reported in this paper were presented as a poster presentation at the 20<sup>th</sup> congress of ESCP (September 6<sup>th</sup>, 2025) and oral communication at the 1<sup>st</sup> mutual French annual congress of surgery (3CVD). The manuscript, including related data, figures and tables has not been previously published and the manuscript is not under consideration elsewhere.



### ***Contribution of each author:***

All authors confirm that they meet the current criteria of the International Committee of Medical Journal Editors (ICMJE) for authorship.

All authors did:

- Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work;
- And drafted the work or reviewed it critically for important intellectual content
- Approved this version to be published
- And agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

In details:

Design of the study: TH, MK, AA, CS and AM

Collection of data: TH, AA, CS, AM

Statistical analysis: TH, AM

Initial manuscript: All authors

Reviewing before publication: All authors

***Category: Original investigation***

***Counts: words, references, figures:***

- Words:
  - Mini-Abstract: 41 words
  - Abstract: 245 words
  - Text (without references): 2967 words
- References: 37
- Table and Figures:
  - Total: 5
  - Tables: 3
  - Figures: 2
- Supplementary Data: 2 (2 figures)

## Text of the Paper

### *Introduction*

Acute perforated sigmoid diverticulitis with generalized peritonitis (Hinchey III–IV) is a surgical emergency requiring rapid source control while balancing postoperative morbidity and the risk of a permanent stoma (1,2). Historically, Hartmann’s procedure was the standard of care but was associated with substantial morbidity, non-negligible mortality, and low reversal rates (3,4). Following recent randomized trials and meta-analyses, national and international guidelines have shifted the paradigm, now favoring sigmoid resection with primary anastomosis when possible, as it offers similar short-term safety but significantly higher reversal rates and better long-term stoma-free survival than Hartmann’s Procedure, reserving the latter for unstable or very high-risk patients (5–12). Regarding the omission of a diverting stoma, guidelines remain cautious, suggesting unprotected anastomosis only for highly selected cases (13).

Over the same period, minimally invasive colorectal surgery has expanded into the emergency setting (14,15). While laparoscopic peritoneal lavage has fallen out of favor due to high rates of persistent sepsis, laparoscopic sigmoid resection has emerged as a feasible option(16–18). Current data suggest that in experienced hands, emergency laparoscopic resection could reduce parietal morbidity and hospital stay compared to open surgery (19–22). However, the specific role of the Laparoscopic Hartmann’s Procedure (LHP) remains debated. Retrospective or registry-based studies suggest that this procedure can be performed safely and may reduce overall complications and length of stay compared with open Hartmann’s procedure.

Crucially, the potential benefits of LHP may extend beyond the acute phase. The low reversal rate of the Hartmann’s strategy is often attributed to dense intra-abdominal

adhesions formed after open surgery. Emerging evidence suggests that performing the initial Hartmann's procedure laparoscopically may reduce adhesion formation, thereby facilitating a subsequent laparoscopic reversal (22,23). This "minimally invasive two-stage strategy" could potentially combine the immediate safety of a non-restorative procedure with a higher likelihood of successful future restoration of bowel continuity. It remains unclear, however, whether this minimally invasive approach can compete with the current standard of care. The aim of the present study was to evaluate, in a large national cohort, whether Laparoscopic Hartmann's Procedure offers comparable short-term outcomes and stoma-free survival to Open Sigmoid Resection with Primary Anastomosis and diverting ileostomy in patients admitted for generalized peritonitis (Hinchey III–IV) related to acute perforated sigmoid diverticulitis.

## ***Methods***

### ***Patients***

Data were obtained from a retrospective multicenter database including all adult patients operated on for sigmoid diverticulitis (SD) between January 2010 and September 2021 across 43 French centers. This analysis is part of a study registered at ClinicalTrials.gov (NCT06200857) (24). The need for informed consent was waived owing to the retrospective study design.

Patients were included if they underwent emergency surgery for SD peritonitis classified as Hinchey III or IV, treated either by open resection with primary anastomosis and diverting stoma or by laparoscopic Hartmann's procedure. Hinchey stage was determined from operative reports according to the original Hinchey classification and current guideline definitions, with stage III defined as generalized purulent peritonitis and stage IV as generalized fecal peritonitis (1,25).

Exclusion criteria were non-urgent surgery, Hinchey stages < III, surgical approaches other than the two evaluated strategies, American Society of Anesthesiologists (ASA) score = 4 and/or preoperative admission to an intensive care unit. These latter criteria were used as a surrogate for hemodynamic instability, a parameter not recorded in the database.

### ***Endpoints***

#### **Primary Endpoint**

The primary outcome was the 12-month stoma-free survival, defined as the time from the index emergency surgery to the successful restoration of intestinal continuity, with

1 patients censored at death, loss to follow-up, or at 12 months if no stoma reversal had  
2 occurred.  
3  
4

## 5 **Secondary Endpoints**

6  
7  
8 Secondary outcomes included intraoperative morbidity (any adverse event occurring  
9 during the index procedure), the conversion rate in the LHP group, 90-day postoperative  
10 morbidity according to Clavien–Dindo classification(26), length of hospital stay,  
11  
12 recurrent diverticulitis during follow-up, and the occurrence of incisional hernias.  
13  
14  
15  
16  
17

## 18 ***Surgical procedure***

19  
20  
21 Two types of procedure were assessed: the Laparoscopic Hartmann’s Procedure (LHP  
22 group)(27), consisting of laparoscopic sigmoidectomy with rectal closure and an end  
23 colostomy, with or without conversion to open surgery; and the Open sigmoidectomy  
24 with a primary anastomosis and a defunctioning stoma (OPA group). In both groups,  
25  
26 decision concerning the lowering of the splenic flexure and abdominal drainage were  
27 left at the surgeon’s discretion.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

## 40 ***Statistical analysis***

41  
42  
43 In this observational study, propensity score matching was used to reduce selection bias  
44 related to the surgical approach and other confounding factors, ensuring comparability  
45 between groups by adjusting for baseline differences between patients undergoing either  
46  
47 LHP or OPA. Propensity scores were estimated using multivariable logistic regression  
48 including all pre-specified covariates. These covariates comprised age, sex, smoking  
49  
50 history, BMI, ASA score, immunodepression, anticoagulant therapy, previous  
51  
52 laparotomy, and the Hinchey classification. These factors were selected as known  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 confounders of both the choice of surgical approach and perioperative outcomes. An  
2 "optimal" case-control match without replacement was used, minimizing absolute  
3  
4 differences in propensity scores (28). Covariate balance after matching was assessed  
5  
6 using standardized mean differences (SMD), with values < 0.1 indicating adequate  
7  
8 balance (Supplementary Figure S1) (28). Missing data were addressed using pairwise  
9  
10 deletion. Continuous variables were reported as median  $\pm$  interquartile range (IQR) and  
11  
12 compared using paired t-tests for approximately normally distributed variables  
13  
14 (normality assessed visually and with the Shapiro–Wilk test), or Wilcoxon signed-rank  
15  
16 tests otherwise. Categorical variables were expressed as frequency (%) and compared  
17  
18 within matched pairs using McNemar’s test.  
19  
20  
21  
22  
23

24 A Cox proportional hazards model was used to compare 12-month stoma-free survival  
25  
26 between groups in the matched cohort. To account for the matched design and within-  
27  
28 pair correlation, a robust sandwich standard errors clustered by matched pairs was  
29  
30 applied (29). A treatment-by-Hinchee interaction was tested in the matched Cox model.  
31  
32  
33 Additionally, a supplementary analysis separated patients who underwent a  
34  
35 laparoscopic Hartmann procedure with or without conversion, to explore whether  
36  
37 conversion modified stoma-free survival.  
38  
39  
40

41 Results are presented as hazard ratios (HR) with 95% confidence intervals (CI).  
42  
43 Statistical analyses were performed using RStudio software version 4.5.1 (R Foundation  
44  
45 for Statistical Computing, Vienna, Austria). A p-value < 0.05 was considered  
46  
47 statistically significant.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## **Results**

### ***Preoperative characteristics***

A total of 1,521 patients underwent surgery for diverticular peritonitis (Hinchey III or IV) between January 2010 and September 2021 across 43 centers, including 34 university hospitals. The final study population comprised 284 patients: 138 in the LHP group and 146 in the OPA group (Fig. 1). After propensity score matching, two comparable groups of 101 patients each were obtained, with adequate balance across all baseline covariates (SMD < 0.1 for covariates included; Supplementary Fig. S1).

Before matching, patients in the LHP group were older (median age 63 vs 50 years,  $p = 0.03$ ), more frequently classified as ASA 3 (26% vs 17%,  $p = 0.04$ ), and had a higher proportion of Hinchey IV peritonitis (37% vs 23%,  $p = 0.01$ ). Other baseline characteristics were similar between groups.

After matching, severe diverticulitis remained frequent, with Hinchey IV peritonitis observed in 31% of patients in the LHP group and 29% in the OPA group, and no other differences in preoperative characteristics were observed (Table 1).

### ***Intraoperative findings and adverse events***

In the propensity score–matched (PSM) cohort, intraoperative outcomes are summarized in Table 2. Median operative time tended to be shorter in the LHP group compared with the OPA group (138 [100–169] vs 180 [120–240] minutes), although this difference did not reach statistical significance ( $p = 0.065$ ). Conversion to open surgery occurred in 49% of patients undergoing LHP.

Splenic flexure mobilization was performed far less frequently in the LHP group than in the OPA group (15% vs 72%,  $p < 0.001$ ), as was high ligation of the inferior mesenteric



artery (4.9% vs 14%,  $p = 0.003$ ), with high ligation defined as division of the inferior mesenteric artery proximal to the origin of the left colic artery (30). As expected, anastomosis was performed exclusively in the OPA group, most commonly as a latero-lateral configuration (59%), Intraoperative transfusion rates were low and comparable between groups (3.3% vs 4.3%,  $p > 0.9$ ).

### ***Postoperative outcomes***

Postoperative outcomes are detailed in Table 3. Overall 90-day morbidity was high but comparable between groups (53% in the LHP group vs 50% in the OPA group,  $p = 0.6$ ). Ninety-day mortality did not significantly differ (5.0% vs 3.0%,  $p = 0.5$ ), nor did the rate of intensive care unit admission (16% vs 11%,  $p = 0.3$ ).

The severity of postoperative complications was similar between groups, with no difference in the distribution of Clavien–Dindo grades ( $p = 0.5$ ). Severe complications (Clavien–Dindo  $>2$ ) occurred in 27% of patients in the LHP group and 26% in the OPA group ( $p > 0.9$ ). Anastomotic leakage was observed in 8 patients (8%) in the OPA group. Reintervention rates for complication were comparable between two groups (10% vs 16%,  $p = 0.2$ ), as well as median length of stay and thirty-day readmission rates were comparable.

Longer-term postoperative events, including incisional hernia (21% vs 23%), and recurrence of diverticulitis, were not significantly different between groups.

Anastomotic stenosis concerned 2 patients (2,2%) in OPA group.

### ***Stoma-Free Survival***

At 12 months, the probability of being stoma-free was higher in the OPA group than in the LHP group (85% vs 65%, respectively; stratified log-rank  $p < 0.0001$ ) (Figure 1).

1 The median follow-up duration, estimated using the reverse Kaplan-Meier method, was  
2 29 months in the LHP group and 70 months in the OPA group. In the Cox proportional  
3 hazards model accounting for the matched design with robust standard errors, OPA was  
4 associated with a significantly higher rate of stoma closure compared with LHP (HR =  
5 1.84, 95% CI 1.39–2.44;  $p < 0.001$ ), corresponding to a shorter time to stoma-free status  
6 (Figure 2). As an effect-modification analysis by peritonitis severity, the hazard ratio for  
7 stoma closure (OPA vs LHP) was 2.07 (95% CI, 1.44–2.99) in Hinchey III and 1.44  
8 (95% CI, 0.88–2.38) in Hinchey IV, with no significant treatment-by-Hinchey  
9 interaction ( $p = 0.269$ ).  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

21 Additionally, a supplementary analysis according to conversion status within the LHP  
22 group was performed (Supplementary Fig. S2). OPA was associated with a higher rate  
23 of stoma closure compared with both converted (HR = 1.59, 95% CI 1.12–2.27;  $p =$   
24 0.01) and non-converted LHP (HR = 2.09, 95% CI 1.44–3.01;  $p < 0.001$ ). No significant  
25 difference in stoma-free survival was observed between converted and non-converted  
26 LHP (HR = 1.34, 95% CI 0.85–2.13;  $p = 0.21$ ).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Discussion

Randomized trials have established primary anastomosis as the preferred reconstructive strategy over Hartmann's procedure for perforated diverticulitis with generalized peritonitis. In both the LADIES (DIVA arm) and DIVERTI trials, short-term outcomes were broadly comparable between strategies, whereas long-term stoma outcomes favored primary anastomosis (8,10). However, these trials were not designed to assess surgical approach and do not address whether laparoscopy can mitigate the long-term limitations of a non-restorative strategy. Our study aimed to specifically study the impact of laparoscopy when performing a Hartmann's procedure (LHP) in comparison with the reference intervention including open sigmoid resection with primary anastomosis (OPA) and diverting ileostomy.

The main finding of this propensity score-matched comparison is that long-term stoma-free survival was lower after LHP than after OPA, despite similar short-term outcomes. At 12 months, the probability of being stoma-free was 85% in the OPA group versus 65% in the LHP group (log-rank  $p < 0.0001$ ), with an almost twofold higher hazard of stoma closure after OPA than after LHP (HR 1.84, 95% CI 1.39-2.44). Importantly, we found no statistically significant interaction between treatment strategy and Hinchey stage, suggesting that the long-term stoma-free survival advantage associated with OPA was not restricted to one severity stratum. This interaction analysis should nonetheless be interpreted cautiously given the lower precision in Hinchey IV.

Accordingly, we selected open primary anastomosis with diverting ileostomy as the reference strategy, as it reflects contemporary practice for stable patients with generalized peritonitis. Although omission of diversion has been proposed in selected cases, current guidelines remain cautious and generally restrict unprotected anastomosis

1 to highly selected patients.(5–7) The anastomotic leak rate observed in the OPA group  
2 (8%) in our study warrants specific discussion, as it directly intersects with the issue of  
3 diversion. A protective stoma can effectively mitigate the clinical consequences of an  
4 anastomotic leak; however, it is also associated with its own morbidity, and its reversal  
5 requires an additional operation, which has been reported to carry complication rates at  
6 14 – 21% (31,32). Generalized peritonitis has historically raised concerns regarding  
7 anastomotic safety. Nevertheless, contemporary randomized trials consistently  
8 demonstrate acceptable leak rates even in the absence of diversion, with short-term  
9 outcomes comparable to those of primary anastomosis with diverting ileostomy  
10 (LADIES and DIVERTI trials). The present results do not provide sufficient evidence to  
11 advocate routine omission of diversion in this setting. Rather, they support a pragmatic  
12 interpretation whereby a protected anastomosis represents a reasonable compromise  
13 between short-term safety and the long-term benefit of higher stoma-free survival  
14 (13,33,34). The randomized controlled trial DIVERTI2 specifically investigated the role  
15 of a protective stoma following primary anastomosis for Hinchey III-IV peritonitis  
16 patients, with overall morbidity at one year as the primary endpoint (13). To avoid  
17 stoma formation, the latest alternative remains laparoscopic lavage. Based principally  
18 on three randomized controlled trials (DILALA, LOLA, and SCANDIV), this  
19 intervention has been shown to be safe and technically straightforward; however, it is  
20 associated with notable drawbacks, including a higher risk of recurrent diverticulitis and  
21 an increased need for reinterventions.

22 The present findings can be positioned within the limited and heterogeneous  
23 observational literature on minimally invasive emergency surgery for perforated  
24 diverticulitis. Vennix et al. reported that emergency laparoscopic sigmoid resection is  
25 feasible in selected settings and may be associated with lower morbidity and shorter  
26

hospital stay compared with open surgery, but their analysis primarily addressed the impact of surgical approach within mixed reconstructive strategies and therefore cannot establish strategy-level equivalence (20). Turley et al., using a large propensity-matched NSQIP analysis restricted to Hartmann's procedures, found no significant short-term advantage of laparoscopy over open surgery, although interpretation is limited by the absence of granular clinical data and by the lack of information on Hinchey status or on long-term stoma outcomes (19). At the other end of the spectrum, Cassini et al. described a totally laparoscopic two-step strategy with high feasibility of laparoscopic reversal in selected patients treated in expert centers, providing proof of technical feasibility rather than comparative evidence against reconstructive strategies (22). Taken together, these studies support that LHP is feasible and may improve selected perioperative metrics in some contexts, but they do not demonstrate equivalence with the contemporary benchmark strategy of primary anastomosis with diversion for long-term stoma-free survival, an issue directly addressed by the present matched comparative design

A key complementary result comes from the analysis according to conversion status within the LHP group, which occurred in approximately half of cases. This is similar to Vennix et al. ( $\approx 54\%$ ) and contrasts with the lower rate reported in a highly selected expert series by Cassini et al. (10%) (20,22). Conversion rates cannot be reliably extracted from registry-based analyses such as Turley et al. (19,20,22). Despite this high conversion rate, OPA remained associated with a higher rate of stoma closure compared with both converted and non-converted LHP, while no significant difference in stoma-free survival was observed between converted and non-converted LHP. These findings indicate that the unfavorable long-term trajectory after LHP is not explained by secondary conversion to an open Hartmann's procedure but rather reflects the intrinsic

1 limitations of the Hartmann strategy itself, regardless of surgical approach. The present  
2 dataset did not allow analysis of the specific causes of conversion, which prevents  
3  
4 exploring whether conversion was driven by disease severity, technical difficulty, or  
5  
6 intraoperative findings.  
7

8  
9 Laparoscopic Hartmann's procedure has been advocated on theoretical grounds in this  
10 emergency setting, including reduced parietal trauma and wound morbidity, potentially  
11  
12 faster postoperative recovery, and the hypothesis that a minimally invasive first stage  
13  
14 may facilitate subsequent reversal by limiting adhesions (35). However, in our study,  
15  
16 short-term outcomes were similar between strategies, with no significant differences in  
17  
18 90-day morbidity or mortality. This aligns with findings from Turley et al., in which a  
19  
20 laparoscopic Hartmann's procedure did not significantly reduce postoperative morbidity  
21  
22 or mortality compared with open surgery in emergency diverticulitis, with comparable  
23  
24 30-day mortality (4.5% open vs 3.0% laparoscopic) and overall morbidity (30% vs  
25  
26 25%) between matched groups (19). Although laparoscopy is associated with improved  
27  
28 outcomes in elective colorectal surgery, recent reviews suggests these benefits are not  
29  
30 consistently seen in emergency settings, and reviews of emergency laparoscopic  
31  
32 colorectal surgery indicate that the technical challenges of peritonitis can limit the  
33  
34 expected advantages of a minimally invasive approach (33,34,36,37).  
35  
36

37  
38 This study has several strengths that support the robustness and relevance of the present  
39  
40 findings. It relies on a national multicenter surgical database (AFC), provides one of the  
41  
42 largest reported series specifically focused on laparoscopic Hartmann's procedure in  
43  
44 this emergency indication, and uses propensity score matching to reduce baseline  
45  
46 imbalance when benchmarking LHP against the contemporary reference strategy. The  
47  
48 present findings can be positioned within the limited and heterogeneous observational  
49  
50 literature. As a retrospective analysis, the comparison remains exposed to residual  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 confounding despite propensity-score matching. Key clinical variables were not  
2 available, including the initial intention regarding restoration of continuity, which may  
3 have influenced both treatment allocation and subsequent stoma management.  
4  
5 Approximately half of LHP required conversion, and the dataset did not capture the  
6 reasons for conversion. Additionally, detailed information on the modalities of  
7 restoration of continuity was not available (surgical approach at reversal and whether  
8 reconstruction was performed with or without diversion), and quality-of-life outcomes  
9 were not collected.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 ***Conclusions***

22  
23 Our findings support a pragmatic interpretation for current practice in generalized  
24 diverticular peritonitis. In this matched national cohort, performing Hartmann's  
25 procedure laparoscopically did not translate into clearly improved short-term outcomes  
26 and did not offset the long-term functional disadvantage inherent to a non-restorative  
27 strategy. When patients are stable and eligible for reconstruction, open sigmoid  
28 resection with protected primary anastomosis therefore remains the reference approach  
29 to maximize the likelihood of long-term stoma-free survival whereas Hartmann's  
30 procedure should be reserved for unstable or very high-risk patients.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Acknowledgements

The authors acknowledge the contribution of all investigators and collaborators of the AFC group involved in the creation and maintenance of the national database :

Jean-Marc Regimbeau, Amiens University Hospital, Amiens, France; Julien Gillet, Angers University Hospital, Angers, France; Antoine Hamy, Angers University Hospital, Angers, France; Amine Lahmidi, Argentan Hospital, Argentan, France; Mohamed Sebbagh, Argentan Hospital, Argentan, France; Antoine Bres, Argentan Hospital, Argentan, France; Nicolas Girard, Bayeux Hospital, Bayeux, France; Aurélie Desjouis, Bayeux Hospital, Bayeux, France; Nadjib Ainseba, Beauvais Hospital, Beauvais, France; Julien Epailly, Beauvais Hospital, Beauvais, France; Fabienne Huysman, Beauvais Hospital, Beauvais, France; Maud Dezeustre, Besançon University Hospital, Besançon, France; Alexandre Doussot, Besançon University Hospital, Besançon, France; Célia Turco, Besançon University Hospital, Besançon, France; Diane Bourrhis, Brest University Hospital, Brest, France; Camille Godet, Caen University Hospital, Caen, France; Morgane Finocchi, Caen University Hospital, Caen, France; Yoan Marion, Caen University Hospital, Caen, France; Gil Lebreton, Caen University Hospital, Caen, France; Flavie Benard, INSERM 1086 ANTICIPE Unity, Caen, France; Elodie Pineau, INSERM 1086 ANTICIPE Unity, Caen, France; Véronique Bouvier, INSERM 1086 ANTICIPE Unity, Caen, France; Pablo Ortega-Deballon, Dijon University Hospital, Dijon, France; Nathan Moreno-Lopez, Dijon University Hospital, Dijon, France; Patrick Rat, Dijon University Hospital, Dijon, France; Bassem Makar, Flers Hospital, Flers, France; Paola Fara, Flers Hospital, Flers, France; Marc Pompilio, Flers Hospital, Flers, France; Hugo Hertault, Martinique Hospital, Fort-de-France, France; Corentin Guezennec, Martinique Hospital, Fort-de-France, France; Habchi



Saida, Avranches-Granville Hospital, Avranches, France; Alexandre Brinza,  
Avranches-Granville Hospital, Avranches, France; Laurent Plard, Avranches-Granville  
Hospital, Avranches, France; Antoine Ailhaud, Grenoble University Hospital, Grenoble,  
France; Léa Gutierrez, Grenoble University Hospital, Grenoble, France; Vincent  
Venchiarutti, Grenoble University Hospital, Grenoble, France; Jean-Luc Faucheron,  
Grenoble University Hospital, Grenoble, France; Bertrand Morin, Le Havre Hospital,  
Le Havre, France; Bertrand Pech de la clause, Le Havre Hospital, Le Havre, France;  
Eglantine Voiteiller, Le Havre Hospital, Le Havre, France; Guillaume Gagnat, Le Havre  
Hospital, Le Havre, France; Barbara Noiret, University Hospital of Lille, Lille, France;  
Constance Laroye, University Hospital of Lille, Lille, France; Guillaume Piessen,  
University Hospital of Lille, Lille, France; Katia Lécolle, Huriez University Hospital,  
Lille, France; Jean Robert, Huriez University Hospital, Lille, France;  
Nzamushelepanmabla, Huriez University Hospital, Lille, France; Perrine Engel,  
Limoges University Hospital, Limoges, France; Muriel Mathonnet, Limoges University  
Hospital, Limoges, France; Cyprien Rodriguez, Lisieux Hospital, Lisieux, France;  
Ammar Makki, Lisieux Hospital, Lisieux, France; Anne-Cécile Couchard, Longjumeau  
Hospital, Longjumeau, France; François Duramé, Longjumeau Hospital, Longjumeau,  
France; Cécile Marceau, Longjumeau Hospital, Longjumeau, France; Jean-Christophe  
Paquet, Longjumeau Hospital, Longjumeau, France; Vahan Kepenekian, Edouard-  
Herriot University Hospital, Lyon, France; Guillaume Passot, Edouard-Herriot  
University Hospital, Lyon, France; Olivier Glehen, Edouard-Herriot University  
Hospital, Lyon, France; Ninon Wallach, La Croix-Rousse University Hospital, Lyon,  
France; Kayvan Mohkam, La Croix-Rousse University Hospital, Lyon, France; Marie-  
Caroline Blanc, North University Hospital, Marseille, France; Stéphane Berdah, North  
University Hospital, Marseille, France; Mathilde Aubert, Timone University Hospital,

Marseille, France; Ahmet Ayav, Nancy University Hospital, Nancy, France; Hervé  
Chanty, Nancy University Hospital, Nancy, France; Alexandre Didier, Nancy  
University Hospital, Nancy, France; Estelle Vigneron, Nancy University Hospital,  
Nancy, France; Noémie Périard, Nantes University Hospital, Nantes, France; Marie  
Thaumoux, Nantes University Hospital, Nantes, France; Océane Grenouilloux, Nice  
University Hospital, Nice, France; Seila Fall, Nice University Hospital, Nice, France;  
Sophoclis Thomareis, Nîmes University Hospital, Nîmes, France; Chloé Marsollier,  
Nîmes University Hospital, Nîmes, France; Michel Prud'homme, Nîmes University  
Hospital, Nîmes, France; Christos Antonopulos, Hospital Bichat-Claude-Bernard  
(APHP), Paris, France; Simon Msika, Hospital Bichat-Claude-Bernard (APHP), Paris,  
France; Marie Monsinjon, Hospital Beaujon (APHP), Clichy, France; Massimo Giacca,  
Hospital Beaujon (APHP), Clichy, France; Antoine Cazelles, European Georges  
Pompidou and Pitié-Salpêtrière Hospitals (APHP), Paris, France; Mehdi Karoui,  
European Georges Pompidou and Pitié-Salpêtrière Hospitals (APHP), Paris, France;  
Valentina Malerba, Antoine Beclere Hospital (APHP), Clamart, France; Ibrahim  
Dagher, Antoine Beclere Hospital (APHP), Clamart, France; Nicola de Angelis, Henri-  
Mondor Hospital (APHP), Créteil, France; Francesco Brunetti, Henri-Mondor Hospital  
(APHP), Créteil, France; Omar Ahmed, Avicenne University Hospital (APHP), Paris,  
France; Jean-Marc Ferraz, Montsouris Institut, Paris, France; Yassir Ramjaun,  
Montsouris Institut, Paris, France; Christine Denet, Montsouris Institut, Paris, France;  
Basile Fuchs, Cochin Hospital (APHP), Paris, France; Daniel Pietrasz, Hôpital Saint-  
Louis (APHP), Paris, France; Pénélope Raimbert, Hôpital Saint-Louis (APHP), Paris,  
France; Juliette Benedetti, Hôpital Saint-Louis (APHP), Paris, France; Chloé Coton,  
Hôpital Saint-Louis (APHP), Paris, France; Hélène Corte, Hôpital Saint-Louis (APHP),  
Paris, France; Aude Medrignac, Rennes University Hospital, Rennes, France; Daphné

Chavrier, Rennes University Hospital, Rennes, France; Robin Girard, Rouen University  
 Hospital, Rouen, France; Julia Berthelot, Rouen University Hospital, Rouen, France;  
 Matthieu Monge, Rouen University Hospital, Rouen, France; Eloïse Papet, Rouen  
 University Hospital, Rouen, France; Jean Pinson, Rouen University Hospital, Rouen,  
 France; Elise Dolbeau, Rouen University Hospital, Rouen, France; Jean-Jacques Tuech,  
 Rouen University Hospital, Rouen, France; Julien Chautard, Saint-Lô Hospital, Saint-  
 Lô, France; Mathilde Boullier, Saint-Lô Hospital, Saint-Lô, France; François Goujard,  
 Saint-Lô Hospital, Saint-Lô, France; Laurent Perrot, Saint-Lô Hospital, Saint-Lô,  
 France; Ludovic Loge, Saint-Lô Hospital, Saint-Lô, France; Jean-Michel Petiot, Saint-  
 Lô Hospital, Saint-Lô, France; Simone Mandefredelli, Strasbourg University Hospital,  
 Strasbourg, France; Diane Charleux-Muller, Strasbourg University Hospital,  
 Strasbourg, France; Guillaume Sauvinet, Strasbourg University Hospital, Strasbourg,  
 France; Jean-Baptiste Bertin, Strasbourg University Hospital, Strasbourg, France; Louis  
 Vincent, Strasbourg University Hospital, Strasbourg, France; Blandine Simeu,  
 Strasbourg University Hospital, Strasbourg, France; Jean-Baptiste Delhorme,  
 Strasbourg University Hospital, Strasbourg, France; Cécile Brigand, Strasbourg  
 University Hospital, Strasbourg, France; Serge Rohr, Strasbourg University Hospital,  
 Strasbourg, France; Manon Bouchet, Toulouse University Hospital; Jean-Pierre Duffas,  
 Toulouse University Hospital; Laurent Ghouti, Toulouse University Hospital; Antoine  
 Philis, Toulouse University Hospital; Guillaume Portier, Toulouse University Hospital;  
 Elias Karam, Tours University Hospital; Nicolas Michot, Tours University Hospital;  
 Alice Artus, Tours University Hospital; Emilien Mauny, Tours University Hospital;  
 Céline Bourbao-Tournois, Tours University Hospital; Hugo Barbin, Tours University  
 Hospital; Julien Thiery, Tours University Hospital.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Hinchey EJ, Schaal PG, Richards GK. Treatment of perforated diverticular disease of the colon. *Adv Surg.* 1978;12:85–109.
2. Hanna MH, Kaiser AM. Update on the management of sigmoid diverticulitis. *World J Gastroenterol.* 2021;27:760–781.
3. Regenbogen SE, Hardiman KM, Hendren S, et al. Surgery for Diverticulitis in the 21st Century: A Systematic Review. *JAMA Surg.* 2014;149:292–303.
4. Hallam S, Mothe BS, Tirumulaju R. Hartmann’s procedure, reversal and rate of stoma-free survival. *Ann R Coll Surg Engl.* 2018;100:301–307.
5. Sartelli M, Weber DG, Kluger Y, et al. 2020 update of the WSES guidelines for the management of acute colonic diverticulitis in the emergency setting. *World J Emerg Surg.* 2020;15:32.
6. Hall J, Hardiman K, Lee S, et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Treatment of Left-Sided Colonic Diverticulitis. *Dis Colon Rectum.* 2020;63:728.
7. Haute Autorité de Santé HAS. Prise en charge médicale et chirurgicale de la diverticulite colique. Saint-Denis La Plaine; 2017.
8. Bridoux V, Regimbeau JM, Ouaisi M, et al. Hartmann’s Procedure or Primary Anastomosis for Generalized Peritonitis due to Perforated Diverticulitis: A Prospective Multicenter Randomized Trial (DIVERTI). *J Am Coll Surg.* 2017;225:798–805.

9. Oberkofler CE, Rickenbacher A, Raptis DA, et al. A Multicenter Randomized Clinical Trial of Primary Anastomosis or Hartmann's Procedure for Perforated Left Colonic Diverticulitis With Purulent or Fecal Peritonitis. *Ann Surg*. 2012;256:819.
10. Lambrichts DPV, Vennix S, Musters GD, et al. Hartmann's procedure versus sigmoidectomy with primary anastomosis for perforated diverticulitis with purulent or faecal peritonitis (LADIES): a multicentre, parallel-group, randomised, open-label, superiority trial. *Lancet Gastroenterol Hepatol*. 2019;4:599–610.
11. Cirocchi R, Afshar S, Shaban F, et al. Perforated sigmoid diverticulitis: Hartmann's procedure or resection with primary anastomosis-a systematic review and meta-analysis of randomised control trials. *Tech Coloproctology*. 2018;22:743–753.
12. Acuna SA, Wood T, Chesney TR, et al. Operative Strategies for Perforated Diverticulitis: A Systematic Review and Meta-analysis. *Dis Colon Rectum*. 2018;61:1442–1453.
13. Pinson J, Tuech JJ, Ouaisi M, et al. Role of protective stoma after primary anastomosis for generalized peritonitis due to perforated diverticulitis-DIVERTI 2 (a prospective multicenter randomized trial): rationale and design (nct04604730). *BMC Surg*. 2022;22:191.
14. Esparza Monzavi CA, Naffouje SA, Chaudhry V, et al. Open vs Minimally Invasive Approach for Emergent Colectomy in Perforated Diverticulitis. *Dis Colon Rectum*. 2021;64:319–327.

15. Lee YF, Brown RF, Battaglia M, et al. Laparoscopic Versus Open Emergent Sigmoid Resection for Perforated Diverticulitis. *J Gastrointest Surg Off J Soc Surg Aliment Tract.* 2020;24:1173–1182.
16. Schultz JK, Yaqub S, Wallon C, et al. Laparoscopic Lavage vs Primary Resection for Acute Perforated Diverticulitis: The SCANDIV Randomized Clinical Trial. *JAMA.* 2015;314:1364–1375.
17. Hoek VT, Edomskis PP, Stark PW, et al. Laparoscopic peritoneal lavage versus sigmoidectomy for perforated diverticulitis with purulent peritonitis: three-year follow-up of the randomised LOLA trial. *Surg Endosc.* 2022;36:7764–7774.
18. Marshall JR, Buchwald PL, Gandhi J, et al. Laparoscopic Lavage in the Management of Hinchey Grade III Diverticulitis: A Systematic Review. *Ann Surg.* 2017;265:670.
19. Turley RS, Barbas AS, Lidsky ME, et al. Laparoscopic versus open Hartmann procedure for the emergency treatment of diverticulitis: a propensity-matched analysis. *Dis Colon Rectum.* 2013;56:72–82.
20. Vennix S, Lips DJ, Di Saverio S, et al. Acute laparoscopic and open sigmoidectomy for perforated diverticulitis: a propensity score-matched cohort. *Surg Endosc.* 2016;30:3889–3896.
21. Vennix S, Boersema GS, Buskens CJ, et al. Emergency Laparoscopic Sigmoidectomy for Perforated Diverticulitis with Generalised Peritonitis: A Systematic Review. *Dig Surg.* 2015;33:1–7.

22. Cassini D, Miccini M, Manoochehri F, et al. Emergency Hartmann's Procedure and Its Reversal: A Totally Laparoscopic 2-Step Surgery for the Treatment of Hinchey III and IV Diverticulitis. *Surg Innov*. 2017;24:557–565.
23. Tan L, Liu XY, Zhang B, et al. Laparoscopic versus open Hartmann reversal: a propensity score matching analysis. *Int J Colorectal Dis*. 2023;38:22.
24. Alves A, Sabbagh C, Ouaisi M, et al. Influence of Socioeconomic Deprivation on Surgical Outcomes for Patients With Sigmoid Diverticulitis in France: A Multicenter Retrospective Study. *Ann Surg*. 2024;280:858–869.
25. Klarenbeek BR, de Korte N, van der Peet DL, et al. Review of current classifications for diverticular disease and a translation into clinical practice. *Int J Colorectal Dis*. 2012;27:207–214.
26. Clavien PA, Barkun J, de Oliveira ML, et al. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg*. 2009;250:187–196.
27. Hartmann H. Nouveau procédé d'ablation des cancers de la partie terminale du colon pelvien. In Strasbourg, France; 1921.
28. Austin PC. An Introduction to Propensity Score Methods for Reducing the Effects of Confounding in Observational Studies. *Multivar Behav Res*. 2011;46:399–424.
29. Austin PC, Cafri G. Variance estimation when using propensity-score matching with replacement with survival or time-to-event outcomes. *Stat Med*. 2020;39:1623–1640.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
30. Pezim ME, Nicholls RJ. Survival after high or low ligation of the inferior mesenteric artery during curative surgery for rectal cancer. *Ann Surg.* 1984;200:729–733.
31. Luglio G, Pendlimari R, Holubar SD, et al. Loop Ileostomy Reversal After Colon and Rectal Surgery: A Single Institutional 5-Year Experience in 944 Patients. *Arch Surg.* 2011;146:1191–1196.
32. Sharma A, Deeb AP, Rickles AS, et al. Closure of defunctioning loop ileostomy is associated with considerable morbidity. *Colorectal Dis.* 2013;15:458–462.
33. Wu KL, Lee KC, Liu CC, et al. Laparoscopic versus Open Surgery for Diverticulitis: A Systematic Review and Meta-Analysis. *Dig Surg.* 2016;34:203–215.
34. Lin H, Zhuang Z, Huang X, et al. The role of emergency laparoscopic surgery for complicated diverticular disease. *Medicine (Baltimore).* 2020;99:e22421.
35. Capitano S. Laparoscopic Hartmann: Believe in It. *Dis Colon Rectum.* 2021;64:e41.
36. Abraha I, Binda GA, Montedori A, et al. Laparoscopic versus open resection for sigmoid diverticulitis. *Cochrane Database Syst Rev.* 2017;2017:CD009277.
37. Klarenbeek BR, Veenhof AA, Bergamaschi R, et al. Laparoscopic Sigmoid Resection for Diverticulitis Decreases Major Morbidity Rates: A Randomized Control Trial: Short-term Results of the Sigma Trial. *Ann Surg.* 2009;249:39.

## Legend for Illustrations

### Figure 1: Flow chart for patient inclusion

Flow diagram of patient selection and propensity score matching. Among 1521 patients in the database, 284 met inclusion criteria (LHP n = 138; OPA n = 146). After 1:1 propensity score matching, 101 patients were included in each group.

### Table 1: Preoperative characteristics

### Table 2: Intraoperative outcomes of propensity-score matched patients

### Table 3: Postoperative outcomes of propensity-score matched patients

### Figure 2: Stoma free survival according to surgical approach

Kaplan–Meier estimates of 12-month stoma-free survival in propensity score–matched patients undergoing LHP (red) versus OPA (blue). Comparisons used a stratified log-rank test; HRs were derived from a Cox model with robust variance accounting for matched pairs. Numbers at risk are shown.

## List of Supplemental Digital Content

### **Supplementary Figure S1: Covariate balance before and after propensity score matching.**

Absolute standardized mean differences (SMD) for all covariates in the unmatched and matched samples. The vertical dashed line indicates the predefined threshold for acceptable balance (SMD = 0.1). All covariates achieved SMD < 0.1 after matching, indicating adequate covariate balance between groups.

### **Supplementary Figure S2: Stoma-free survival according to surgical approach and conversion status**

Kaplan–Meier estimates of stoma-free survival up to 12 months after index surgery in propensity score–matched patients, comparing laparoscopic Hartmann’s procedure not converted to open surgery (LHP non-converted, orange), LHP converted to open surgery (LHP converted, pink), and open procedure (OPA, blue). Follow-up was censored at 12 months. Hazard ratios and corresponding p-values were estimated using Cox proportional hazards models with robust standard errors clustered by matched pairs. Numbers at risk at selected time points are displayed below the plot.

**Table 1: Preoperative characteristics**

| Characteristic                                                                                                                                      | All patients (n=284) |                   |       | Matched patients (n=202) |                   |      |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|----------------------|-------------------|-------|--------------------------|-------------------|------|
|                                                                                                                                                     | LHP<br>(n=138)       | OPA<br>(n=146)    | P     | LHP<br>(n=101)           | OPA<br>(n=101)    | P    |
| Age (year)                                                                                                                                          | 63 [51, 76]          | 50 [50, 69]       | 0.03* | 58 [49, 72]              | 60 [51, 73]       | 0.2  |
| Gender, male                                                                                                                                        | 73 (52,9%)           | 88 (59,9%)        | 0.2   | 58 (57%)                 | 55 (55%)          | 0.6  |
| ASA score > 2                                                                                                                                       | 36 (26%)             | 25 (17%)          | 0.04* | 21 (21%)                 | 18 (18%)          | 0.6  |
| BMI kg/m <sup>2</sup>                                                                                                                               | 25.5 [22.3, 28.3]    | 25.4 [22.9, 29.4] | 0.7   | 25.5 [22.3, 28.3]        | 25.1 [21.7, 28.6] | 0.9  |
| Active smoker                                                                                                                                       | 30 (21,7%)           | 45 (30,6%)        | 0.09  | 30 (30%)                 | 27 (27%)          | 0.6  |
| Immunodepression                                                                                                                                    | 24 (17,4%)           | 14 (9,5%)         | 0.051 | 15 (15%)                 | 14 (14%)          | 0.8  |
| Anticoagulation                                                                                                                                     | 13 (9,4%)            | 9 (6,1%)          | 0,3   | 8 (8.5%)                 | 5 (5.1%)          | 0.2  |
| <b>Medical antecedents</b>                                                                                                                          |                      |                   |       |                          |                   |      |
| Cardiologic                                                                                                                                         | 50 (36,2%)           | 59 (40,1%)        | 0.5   | 29 (30%)                 | 41 (42%)          | 0.13 |
| Neurologic                                                                                                                                          | 23 (16,7%)           | 19 (12,9%)        | 0.4   | 14 (14%)                 | 12 (12%)          | 0.7  |
| Pulmonary                                                                                                                                           | 24 (17,4%)           | 18 (12,2%)        | 0.2   | 15 (16%)                 | 11 (11%)          | 0.6  |
| Diabetes                                                                                                                                            | 10 (7,2%)            | 10 (6,8%)         | 0.9   | 6 (6.2%)                 | 7 (7.1%)          | 0.6  |
| Previous laparotomy                                                                                                                                 | 15 (10,9%)           | 13 (8,8%)         | 0.6   | 10 (9.9%)                | 12 (12%)          | 0.6  |
| Previous diverticulitis                                                                                                                             | 29 (21%)             | 41 (27,9%)        | 0.2   | 23 (24%)                 | 28 (29%)          | 0.5  |
| <b>Actual diverticulitis</b>                                                                                                                        |                      |                   |       |                          |                   |      |
| Hinchey Score                                                                                                                                       |                      |                   |       |                          |                   |      |
| III                                                                                                                                                 | 87 (63%)             | 113 (76,9%)       | 0.01* | 70 (69%)                 | 72 (71%)          | 0.7  |
| IV                                                                                                                                                  | 51 (37%)             | 34 (23,1%)        | 0.01* | 31 (31%)                 | 29 (29%)          | 0.7  |
| Associated hemorrhage                                                                                                                               | 1 (0,7%)             | 0 (0%)            | 0.5   | 1 (1.0%)                 | 0 (0%)            |      |
| Associated occlusion                                                                                                                                | 7 (5,1%)             | 10 (6,8%)         | 0.5   | 6 (5.9%)                 | 6 (5.9%)          | >0.9 |
| Associated fistula                                                                                                                                  | 2 (1,4%)             | 5 (3,4%)          | 0.4   | 2 (2.0%)                 | 4 (4.0%)          | 0.4  |
| Surgery within 24 hours of diagnosis                                                                                                                | 134 (97,1%)          | 139 (94,6%)       | 0.3   | 98 (97%)                 | 98 (97%)          | >0.9 |
| Surgery after 24 hours                                                                                                                              | 1 (0,7%)             | 5 (3,4%)          | 0.6   | 1 (1.0%)                 | 1 (1.0%)          | >0.9 |
| Quantitative variables are expressed as median (interquartile range [IQR]), and qualitative variables are expressed as frequencies and percentages. |                      |                   |       |                          |                   |      |
| ASA, American Society of Anesthesiologists; BMI, body mass index; CRP, C-reactive protein;SD, standard deviation.                                   |                      |                   |       |                          |                   |      |
| *Statistically significant differences.                                                                                                             |                      |                   |       |                          |                   |      |

**Table 2: Intraoperative outcomes of propensity-score matched patients**

|                               | LHP (N = 101)  | OPA (N = 101)  | p-value |
|-------------------------------|----------------|----------------|---------|
| Intraoperative outcome        |                |                |         |
| Operative time, min*          | 138 [100, 169] | 180 [120, 240] | 0.065   |
| Conversion to open surgery†   | 49 (49%)       | Not concerned  |         |
| Splenic flexure mobilization* | 15 (15%)       | 73 (72%)       | <0.001* |
| High IMA ligation*            | 5 (4.9%)       | 14 (14%)       | 0.003*  |
| Anastomose type               |                |                |         |
| Latero-terminal               | Not concerned  | 60 (59%)       |         |
| Termino-terminal              | Not concerned  | 32 (32%)       |         |
| Transfusion                   | 3 (3.3%)       | 4 (4.3%)       | >0.9    |

Quantitative variables are expressed as median (interquartile range [IQR]), and qualitative variables are expressed as frequencies and percentages.

IMA, Inferior Mesentery Artery

\* Statistically significant differences.

† Reasons for conversion are unknown as they were not collected in the database.

**Table 3: Postoperative outcomes of propensity-score matched patients**


|                            | LHP (N = 101) | OPA (N = 101) | p-value |
|----------------------------|---------------|---------------|---------|
| Postoperative outcome      |               |               |         |
| Overall 90 days morbidity  | 54 (53%)      | 50 (50%)      | 0.6     |
| Overall 90 days mortality  | 5 (5.0%)      | 3 (3.0%)      | 0.5     |
| ICU hospitalization, %     | 16 (16%)      | 11 (11%)      | 0.3     |
| Dindo-Clavien              |               |               | 0.5     |
| 1                          | 29 (29%)      | 34 (34%)      |         |
| 2                          | 24 (24%)      | 19 (19%)      |         |
| 3                          | 8 (7.9%)      | 13 (13%)      |         |
| 4                          | 14 (14%)      | 10 (9.9%)     |         |
| 5                          | 5 (5.0%)      | 3 (3.0%)      |         |
| Dindo-Clavien >2           | 27 (27%)      | 26 (26%)      | >0.9    |
| Intraperitoneal abscess    | 7 (7.1%)      | 9 (9.0%)      | 0.6     |
| Anastomotic leakage        |               | 8 (8.0%)      |         |
| Length of hospital stay, d | 12 [8, 19]    | 10 [8, 16]    | 0.095   |
| 30-day readmission         | 15 (16%)      | 19 (21%)      | 0.5     |

Quantitative variables are expressed as median (interquartile range [IQR]), and qualitative variables are expressed as frequencies and percentages.

ICU, intensive care unit; NS, not significant


**Figure 1: Flow chart for patient inclusion**

Flow diagram of patient selection and propensity score matching. Among 1521 patients in the database, 284 met inclusion criteria (LHP n = 138; OPA n = 146). After 1:1 propensity score matching, 101 patients were included in each group.



**Figure 2: Stoma free survival according to surgical approach**


Kaplan–Meier estimates of 12-month stoma-free survival in propensity score–matched patients undergoing LHP (red) versus OPA (blue). Comparisons used a stratified log-rank test; HRs were derived from a Cox model with robust variance accounting for matched pairs. Numbers at risk are shown.






# Supplementary Figure S1: Covariate balance before and after propensity score matching.

Absolute standardized mean differences (SMD) for all covariates in the unmatched and matched samples. The vertical dashed line indicates the predefined threshold for acceptable balance (SMD = 0.1). All covariates achieved SMD < 0.1 after matching, indicating adequate covariate balance between groups.



## Supplementary Figure S2: Stoma-free survival according to surgical approach and conversion status

Kaplan–Meier estimates of stoma-free survival up to 12 months after index surgery in propensity score–matched patients, comparing laparoscopic Hartmann’s procedure not converted to open surgery (LHP non-converted, orange), LHP converted to open surgery (LHP converted, pink), and open procedure (OPA, blue). Follow-up was censored at 12 months. Hazard ratios and corresponding p-values were estimated using Cox proportional hazards models with robust standard errors clustered by matched pairs. Numbers at risk at selected time points are displayed below the plot.



**Table 1: Preoperative characteristics**

|                                                                                                                                                     | All patients (n=284) |                   |       | Matched patients (n=202) |                   |      |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|----------------------|-------------------|-------|--------------------------|-------------------|------|
| Characteristic                                                                                                                                      | LHP<br>(n=138)       | OPA<br>(n=146)    | P     | LHP<br>(n=101)           | OPA<br>(n=101)    | P    |
| Age (year)                                                                                                                                          | 63 [51, 76]          | 50 [50, 69]       | 0.03* | 58 [49, 72]              | 60 [51, 73]       | 0.2  |
| Gender, male                                                                                                                                        | 73 (52,9%)           | 88 (59,9%)        | 0.2   | 58 (57%)                 | 55 (55%)          | 0.6  |
| ASA score > 2                                                                                                                                       | 36 (26%)             | 25 (17%)          | 0.04* | 21 (21%)                 | 18 (18%)          | 0.6  |
| BMI kg/m2                                                                                                                                           | 25.5 [22.3, 28.3]    | 25.4 [22.9, 29.4] | 0.7   | 25.5 [22.3, 28.3]        | 25.1 [21.7, 28.6] | 0.9  |
| Active smoker                                                                                                                                       | 30 (21,7%)           | 45 (30,6%)        | 0.09  | 30 (30%)                 | 27 (27%)          | 0.6  |
| Immunodepression                                                                                                                                    | 24 (17,4%)           | 14 (9,5%)         | 0.051 | 15 (15%)                 | 14 (14%)          | 0.8  |
| Anticoagulation                                                                                                                                     | 13 (9,4%)            | 9 (6,1%)          | 0,3   | 8 (8.5%)                 | 5 (5.1%)          | 0.2  |
| <b>Medical antecedents</b>                                                                                                                          |                      |                   |       |                          |                   |      |
| Cardiologic                                                                                                                                         | 50 (36,2%)           | 59 (40,1%)        | 0.5   | 29 (30%)                 | 41 (42%)          | 0.13 |
| Neurologic                                                                                                                                          | 23 (16,7%)           | 19 (12,9%)        | 0.4   | 14 (14%)                 | 12 (12%)          | 0.7  |
| Pulmonary                                                                                                                                           | 24 (17,4%)           | 18 (12,2%)        | 0.2   | 15 (16%)                 | 11 (11%)          | 0.6  |
| Diabetes                                                                                                                                            | 10 (7,2%)            | 10 (6,8%)         | 0.9   | 6 (6.2%)                 | 7 (7.1%)          | 0.6  |
| Previous laparotomy                                                                                                                                 | 15 (10,9%)           | 13 (8,8%)         | 0.6   | 10 (9.9%)                | 12 (12%)          | 0.6  |
| Previous diverticulitis                                                                                                                             | 29 (21%)             | 41 (27,9%)        | 0.2   | 23 (24%)                 | 28 (29%)          | 0.5  |
| <b>Actual diverticulitis</b>                                                                                                                        |                      |                   |       |                          |                   |      |
| Hinchey Score                                                                                                                                       |                      |                   |       |                          |                   |      |
| III                                                                                                                                                 | 87 (63%)             | 113 (76,9%)       | 0.01* | 70 (69%)                 | 72 (71%)          | 0.7  |
| IV                                                                                                                                                  | 51 (37%)             | 34 (23,1%)        | 0.01* | 31 (31%)                 | 29 (29%)          | 0.7  |
| Associated hemorrhage                                                                                                                               | 1 (0,7%)             | 0 (0%)            | 0.5   | 1 (1.0%)                 | 0 (0%)            |      |
| Associated occlusion                                                                                                                                | 7 (5,1%)             | 10 (6,8%)         | 0.5   | 6 (5.9%)                 | 6 (5.9%)          | >0.9 |
| Associated fistula                                                                                                                                  | 2 (1,4%)             | 5 (3,4%)          | 0.4   | 2 (2.0%)                 | 4 (4.0%)          | 0.4  |
| Surgery within 24 hours of diagnosis                                                                                                                | 134 (97,1%)          | 139 (94,6%)       | 0.3   | 98 (97%)                 | 98 (97%)          | >0.9 |
| Surgery after 24 hours                                                                                                                              | 1 (0,7%)             | 5 (3,4%)          | 0.6   | 1 (1.0%)                 | 1 (1.0%)          | >0.9 |
| Quantitative variables are expressed as median (interquartile range [IQR]), and qualitative variables are expressed as frequencies and percentages. |                      |                   |       |                          |                   |      |
| ASA, American Society of Anesthesiologists; BMI, body mass index; CRP, C-reactive protein; SD, standard deviation.                                  |                      |                   |       |                          |                   |      |
| *Statistically significant differences.                                                                                                             |                      |                   |       |                          |                   |      |

**Table 2: Intraoperative outcomes of propensity-score matched patients**

| Intraoperative outcome        | LHP (N = 101)  | OPA (N = 101)  | p-value |
|-------------------------------|----------------|----------------|---------|
| Operative time, min*          | 138 [100, 169] | 180 [120, 240] | 0.065   |
| Conversion to open surgery†   | 49 (49%)       | Not concerned  |         |
| Splenic flexure mobilization* | 15 (15%)       | 73 (72%)       | <0.001* |
| High IMA ligation*            | 5 (4.9%)       | 14 (14%)       | 0.003*  |
| Anastomose type               |                |                |         |
| Latero-terminal               | Not concerned  | 60 (59%)       |         |
| Termino-terminal              | Not concerned  | 32 (32%)       |         |
| Transfusion                   | 3 (3.3%)       | 4 (4.3%)       | >0.9    |

Quantitative variables are expressed as median (interquartile range [IQR]), and qualitative variables are expressed as frequencies and percentages.

IMA, Inferior Mesentery Artery

\* Statistically significant differences.

† Reasons for conversion are unknown as they were not collected in the database.

**Table 3: Postoperative outcomes of propensity-score matched patients**

|                                                                                                                                                     | LHP (N = 101) | OPA (N = 101) | p-value |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|---------------|---------------|---------|
| <b>Postoperative outcome</b>                                                                                                                        |               |               |         |
| Overall 90 days morbidity                                                                                                                           | 54 (53%)      | 50 (50%)      | 0.6     |
| Overall 90 days mortality                                                                                                                           | 5 (5.0%)      | 3 (3.0%)      | 0.5     |
| ICU hospitalization, %                                                                                                                              | 16 (16%)      | 11 (11%)      | 0.3     |
| Dindo-Clavien                                                                                                                                       |               |               | 0.5     |
| 1                                                                                                                                                   | 29 (29%)      | 34 (34%)      |         |
| 2                                                                                                                                                   | 24 (24%)      | 19 (19%)      |         |
| 3                                                                                                                                                   | 8 (7.9%)      | 13 (13%)      |         |
| 4                                                                                                                                                   | 14 (14%)      | 10 (9.9%)     |         |
| 5                                                                                                                                                   | 5 (5.0%)      | 3 (3.0%)      |         |
| Dindo-Clavien >2                                                                                                                                    | 27 (27%)      | 26 (26%)      | >0.9    |
| Intraperitoneal abscess                                                                                                                             | 7 (7.1%)      | 9 (9.0%)      | 0.6     |
| Anastomotic leakage                                                                                                                                 |               | 8 (8.0%)      |         |
| Length of hospital stay, d                                                                                                                          | 12 [8, 19]    | 10 [8, 16]    | 0.095   |
| 30-day readmission                                                                                                                                  | 15 (16%)      | 19 (21%)      | 0.5     |
| Quantitative variables are expressed as median (interquartile range [IQR]), and qualitative variables are expressed as frequencies and percentages. |               |               |         |
| ICU, intensive care unit; NS, not significant                                                                                                       |               |               |         |

Figure 1 : Flow-chart





Figure 2 : Stoma-Free Survival



### **Conflict of interest/Disclosure**

None of the authors have any conflict of interest.



**Funding/Support**



This research was conducted without external or internal sources of funding.



[Click here to access/download](#)

**Supplemental Data File**

**7S1\_Love\_Plot\_with\_legend\_pdf.pdf**






[Click here to access/download](#)

**Supplemental Data File**

9Sup2\_curves\_with\_legend\_pdf.pdf





[Click here to access/download](#)

**Supplemental Data File**  
**strobe\_hartmann\_annals.docx**

