

A simple two parameter distribution for modelling neuronal activity and capturing neuronal association

Thomas Delaney¹ and Cian O'Donnell¹

¹*School of Computer Science, Electrical and Electronic Engineering, and Engineering
Mathematics, University of Bristol, Bristol, United Kingdom.*

Abstract

Recent developments in electrophysiological technology have lead to an increase in the size of electrophysiological datasets. Consequently, there is a requirement for new analysis techniques that can make use of these new datasets, while remaining easy to use in practice. In this work, we fit the Conway-Maxwell-binomial distribution to spiking data read from a mouse exposed to visual stimuli.

1 Introduction

Motivate by pointing out how much computational power it can require to calculate n th order correlations.

Point out that we don't necessarily need to measure correlations anyway.

17

2 Results

18

3 Discussion

19

4 Data

20

Details from data cortex lab here.

21

5 Methods

22

Details about all kinds of things here.

23

5.1 Binning data

24

We binned the spiking data into very small bins.

25

6 Discussion

26

Point out that the Conway-Maxwell-binomial distribution could be used to measure

27

activity and association without having to sort the voltage traces into spikes. That

28

does defeat the purpose slightly, however.

References

- [1] Patricia M. E. Altham, *Two Generalizations of the Binomial Distribution*. Journal of the Royal Statistical Society 27, 162-167, (1978)
- [2] Mark M Churchland, Byron M Yu, John P Cunningham, Leo P Sugrue, Marlene R Cohen, Greg S Corrado, William T Newsome, Andrew M Clark, Paymon Hosseini, Benjamin B Scott, David C Bradley, Matthew A Smith, Adam Kohn, J Anthony Movshon, Katherine M Armstrong, Tirin Moore, Steve W Chang, Lawrence H Snyder, Stephen G Lisberger, Nicholas J Priebe, Ian M Finn, David Ferster, Stephen I Ryu, Gopal Santhanam, Maneesh Sahani, Krishna V Shenoy, *Stimulus onset quenches neural variability: A widespread cortical phenomenon*. Nature Neuroscience 13, 369-378, (2010)
- [3] Fraser Daly, Robert E. Gaunt, *The Conway-Maxwell-Poisson distribution: Distributional theory and approximation*. ALEA 13, 635-658, (2016)
- [4] Joseph B. Kadane, *Sums of Possibly Associated Bernoulli Variables: The Conway-Maxwell-Binomial Distribution*. Bayesian Analysis 11, 403-420, (2016)
- [5] Joseph B. Kadane, Galit Shmueli, Thomas P. Minka, Sharad Borle, Peter Boatwright, *Conjugate Analysis of the Conway-Maxwell-Poisson Distribution*. Bayesian Analysis 1, (2006)
- [6] R. L. Prentice, *Binary Regression Using an Extended Beta-Binomial Distribution, With Discussion of Correlation Induced by Covariate Measurement Errors*. Journal of American Statistical Association 81, 321-327, (1986)
- [7] Galit Shmueli, Thomas P. Minka, Joseph B. Kadane, Sharad Borle, Peter Boatwright, *A useful distribution for fitting discrete data: revival of the Conway-Maxwell-Poisson distribution*. Applied Statistics 54, 127-142, (2005)
- [8] Joseph S. Verducci, Michael E. Mack, Morris H. DeGroot, *Estimating multiple rater agreement for a rare diagnosis*. Journal of Multivariate Analysis 27, 512-535, (1988)