
# **Evolution of Biodiversity**

Chapter 5



# II. The Living World (10-15%)

- A. Ecosystem Structure
- B. Energy Flow
- C. Ecosystem Diversity
- D. Natural Ecosystems Change
- E. Natural Biogeochemical Cycles



https://ecologynature.com/?p=56

# Module 14: The Biodiversity of Earth


After this module you will be able to.....

- 1) Understand how we estimate the number of species living on Earth
- 2) Quantify biodiversity
- 3) Describe patterns of relatedness among species using phylogeny

# Estimating the number of species can be difficult!

The number of species in any given place is a common measure of biodiversity, but estimating the total number of species on Earth is a challenge.

Scientists have named approximately 2 million species, which means the total must be larger.



https://tenor.com/view/the-hangover-zach-galifianakis-math-card-counting-gambling-gif-3938292

# We can measure biodiversity in terms of species richness and evenness

Species richness The number of species in a given area.

Species evenness The relative proportion of individuals within the different species in a given area.

Knowing the species richness or evenness of an ecosystem gives environmental scientists a baseline they can use to determine how much an ecosystem has changed.


**Shannon's Index:** a measure of biodiversity that takes into account the number of species present and the relative abundance of species present


\_\_


# **Measuring Biodiversity**

Which population has greater richness?

Evenness?







Community 2 A: 70% B: 10% C: 10% D: 10%

Figure 14.2


Environmental Science for AP®, Second Edition
© 2015 W.H. Freeman and Company


**Measures of species diversity.** Species richness and species evenness are two different measures of species diversity. Although both communities contain the same number of species, community 1 has a more even distribution of species and is therefore more diverse than community 2.

The evolutionary relationship among species can be illustrated using a phylogeny

Phylogeny The branching pattern of evolutionary relationships.

The more similar the traits of two species, the more closely related the two species are assumed to be.






A phylogenetic tree. Phylogenies are based on the similarity of traits among species. Scientists can assemble phylogenetic trees that indicate how different groups of organisms are related and show where speciation events have occurred. The brown boxes indicate when major morphological changes evolved over evolutionary time.

# **Module 15: How Evolution Creates Biodiversity**

After this module you will be able to.....

- 1) Identify the processes that cause genetic diversity
- 2) Explain how evolution can occur through artificial selection
- 3) Explain how evolution can occur through natural selection
- 4) Explain how evolution can occur through random processes

# Genetic diversity is created through mutation and recombination



**Evolution** A change in the genetic composition of a population over time

Microevolution Evolution below the species level

Macroevolution Evolution that gives rise to new species, genera, families, classes, or phyla

# Genotype vs Phenotype FLASHBACK!

Gene A physical location on the chromosomes within each cell of an organism

**Genotype** The complete set of genes in an individual

Phenotype A set of traits expressed by an individual

#### Genotype vs. Phenotype

Genotype	Phenotype
BB Homozygous dominant	<b>©</b>
Bb Heterozygous	<b>@</b>
bb Homozygous recessive	<b>Q</b>

http://www.sliderbase.com/spitem-647-1.html

#### **Mutation**



https://www.syfy.com/syfywire/marvels-return-of-wolverine-miniseries-gives-logan-new-abilities

Mutation A random change in the genetic code produced by a mistake in the copying process

When mutations occur in cells responsible for reproduction (gametes) those mutations can be passed on to the next generation.


Sometimes a mutation improves an organism's chance of survival or reproduction. If such a mutation is passed along to the next generation, it adds new genetic diversity to the population.

### Recombination

Recombination The genetic process by which one chromosome breaks off and attaches to another chromosome during reproductive cell division. (CROSSING-OVER)

This process does not create new genes, but brings together new combinations of alleles on a chromosome, producing new traits.

For example, the human immune system must battle a large variety of viruses and bacteria that regularly attempt to invade the body. Recombination allows new allele combinations to come together, which provides new immune defenses.




Each gamete gets one copy of the chromosome, each with a unique combination of alleles.

https://learn.genetics.utah.edu/content/pigeons/geneticlinkage/

## Evolution can occur through artificial selection

Evolution by artificial selection The process in which humans determine which individuals breed, typically with a preconceived set of traits in mind. (SELECTIVE BREEDING)



Artificial selection has produced numerous breeds of livestock and pets.

Most modern agricultural crops are the result of many years of careful breeding.

Artificial selection can also produce unintended results such as herbicide resistance.





Figure 15.3
Environmental Science for AP®, Second Edition
Data from H. G. Parker et al., Science 304 (2004): 1160–1164

**Artificial selection on animals.** The diversity of domesticated dog breeds is the result of artificial selection on wolves. The wolf is the ancestor of the various breeds of dogs. It is illustrated at the same level as the dogs in this phylogeny because it is a species that is still alive today.


### Evolution can occur through natural selection

Evolution by natural selection The process in which the environment determines which individuals survive and reproduce.



Key Ideas of Darwin's theory of evolution

- Individuals produce an excess of offspring
- Not all offspring can survive
- Individuals differ in their traits
- Differences in traits can be passed on from parents to offspring
- Differences in traits are associated with differences in the ability to survive and reproduce



Natural Selection All species produce an excess number of offspring. Only those offspring with the fittest genotypes will pass on their genes to the next generation.


\_\_

#### **Natural Selection**

Natural selection favors any combination of traits that improves an individual's fitness.

Fitness An individual's ability to survive and reproduce.

Adaptation A trait that improves an individual's fitness.



https://platofootnote.wordpress.com/2017/06/26/how-to-properly-study-adaptation/

# Evolution can also occur through random processes

There are five random processes through which evolution occurs:

Mutation A random change in the genetic code produced by a mistake in the copying process

Gene flow The process by which individuals move from one population to another and thereby alter the genetic composition of both populations

Genetic drift A change in the genetic composition of a population over time as a result of random mating


Bottleneck effects A reproduction in the genetic diversity of a population caused by a reduction in its size

Founder effects A change in the genetic composition of a population as a result of descending from a small number colonizing individuals

#### **Mutations**

As the number of mutations accumulates in a population over time, evolution occurs.

If a mutation is not lost, it may increase in frequency over time.



**Figure 15.7** *Environmental Science for AP®*, Second Edition © 2015 W.H. Freeman and Company

### **Gene Flow**


Gene flow The process by which individuals move from one population to another and thereby alter the genetic composition of both populations.

The arrival of individuals from adjacent populations alters the frequency of alleles in the population.

In a population that is experiencing natural or artificial selection, high gene flow from outside can prevent the population from responding to selection.

Gene flow can be helpful in bringing in genetic variation to a population that lacks it.

#### **Gene Flow**



#### Florida

As the population of Florida panthers declined to very low numbers, the percentage of kinked tails increased to approximately 90 percent.

#### Texas

Several Texas panthers were brought to Florida.

#### Florida

After breeding with the Texas panthers, the Florida panther population was less inbred and the percentage of kinked tails declined.

#### Evolution by gene flow.

As the Florida panther declined in population size, the animals experienced low genetic variation and showed signs of inbreeding, which lead to kinky tails, heart defects, and low sperm counts. With the introduction of eight panthers from Texas, the Florida population experienced a decline in the prevalence of defects and a growth in population from 30 to 160 individuals.

Figure 15.8
Environmental Science for AP®, Second Edition
© 2015 W.H. Freeman and Company

### **Genetic Drift**

Genetic drift A change in the genetic composition of a population over time as a result of random mating.

Like mutation and gene flow, genetic drift is a nonadaptive, random process.

Can be particularly important role in altering the genetic composition of small populations.

### **Genetic Drift**

Evolution by genetic drift. (a) In a small population, some lesscommon genotypes can be lost by chance as random mating among a small number of individuals can result in the less-common genotype not mating. As a result, the genetic composition can change over time. (b) In a large population, it is more difficult for the less-common genotypes to be lost by chance because the absolute number of these individuals is large. As a result, the genetic composition tends to remain the same over time in larger populations.

#### (a) Small population



#### (b) Large population




Figure 15.9
Environmental Science for AP®, Second Edition
© 2015 W.H. Freeman and Company

#### **Bottleneck Effect**

Bottleneck effect A reduction in the genetic diversity of a population caused by a reduction in its size.

Reduced population numbers means reduced genetic variation.

Low genetic variation in a population can cause increased risk of disease and low fertility.

The bottleneck effect means species are less able to adapt to future environmental changes.

Resulting low diversity can lead to decline and extinction.

**Extinction** The death of the last member of a species.

# Bottleneck Effect

**Evolution by the bottleneck** effect. If a population experiences a drastic decrease in size (goes through a "bottleneck"), some genotypes will be lost, and the genetic composition of the survivors will differ from the composition of the original group.




Figure 15.10
Environmental Science for AP®, Second Edition
© 2015 W.H. Freeman and Company

### **Founder Effect**

Founder effect A change in the genetic composition of a population as a result of descending from a small number of colonizing individuals.

Evolution by the founder effect. If a few individuals from a mainland population colonize an island, the genotypes on the island will represent only a subset of the genotypes present in the mainland population. As with the bottleneck effect, some genotypes will not be present in the new population.




Figure 15.11
Environmental Science for AP®, Second Edition
© 2015 W.H. Freeman and Company

# Module 16: Speciation and the Pace of Evolution

After this module you will be able to.....

- 1) Explain the processes of allopatric and sympatric speciation
- 2) Understand the factors that affect the pace of evolution

# **Allopatric vs Sympatric**



New species commonly evolve through two processes

Allopatric speciation The process of speciation that occurs with geographic isolation

Tip: Imagine that two groups of people are separated by a bunch of ALLOgators


Sympatric speciation The evolution of one species into two, without geographic isolation

# **Allopatric Speciation**


Geographic Isolation

#### **Geographic isolation**

Physical separation of a group of individuals from others of the same species




## **Allopatric Speciation**



#### Reproductive isolation

The result of two populations within a species evolving separately to the point that they can no longer interbreed and produce viable offspring




# **Allopatric Speciation**

In allopatric speciation Geographic barriers can split populations. Natural selection may favor different traits in the environment of each isolated population, resulting in different adaptations. Over time, the two populations may become so genetically distinct that they are no longer capable of interbreeding.

**Figure 16.1** *Environmental Science for AP®*, Second Edition © 2015 W.H. Freeman and Company

# **Sympatric Speciation**



Sympatric speciation Flowering plants such as wheat commonly form new species through the process of polyploidy, an increase in the number of sets of chromosomes beyond the normal two sets. (a) The ancestral einkorn wheat (Triticum boeoticum) has two sets of chromosomes and produces small seeds. (b) Durum wheat (Triticum durum), which is used to make pasta, was bred to have four sets of chromosomes and produces mediumsized seeds. (c) Common wheat (*Triticum* aestivum), which is used mostly for bread, was bred to have six sets of chromosomes and produces the largest seeds.

Figure 16.3
Environmental Science for AP®, Second Edition
After http://zr.molbiol.ru/poaceae znachenije.html

#### The Pace of Evolution

#### The pace of evolution is dependent on many factors

- A species can survive an environmental change if it can quickly evolve adaptations to new conditions.
- Slow rates of evolution occur when a population has long generation times or contains low genetic variation.
- Evolution by artificial selection can be very rapid.
- Genetically modified organism (GMO) An organism produced by copying genes from a species with a desirable trait and inserting them into another species.

# Module 17: Evolution of Niches and Species Distributions

After this module you will be able to.....

- 1) Explain the difference between a fundamental and a realized niche
- 2) Describe how environmental change can alter species distributions
- 3) Discuss how environmental change can cause species extinctions

# **Every species has a niche**


Range of tolerance The limits to the abiotic conditions that a species can tolerate

Fundamental niche The suite of abiotic conditions under which a species can survive, grow, and reproduce

Realized niche The range of abiotic and biotic conditions under which a species actually lives

Distribution Areas of the world in which a species lives

### **Species niches**



Range of tolerance All species have an ideal range of abiotic conditions, such as temperature, under which their members can survive, grow, and reproduce. Under more extreme conditions, their ability to perform these essential functions declines.


**Temperature** 

Figure 17.1

Environmental Science for AP®, Second Edition
© 2015 W.H. Freeman and Company

# **Species niches**

#### Realized vs. Fundamental Niche



What happened with Balanus was removed?


- Fundamental niche the niche potentially occupied by a species
- Realized niche the niche actually occupied by a species

**Species niches mnemonic** 

# **RAFT**

**Realized Actual** 

**Fundamental Theoretical** 



## **Species niches**


Niche generalist A species that can live under a wide range of abiotic or biotic conditions

Niche specialist A species that is specialized to live in a specific habitat or to feed on a small group of species

Niche generalists fare better under changing conditions because they have a number of alternative habitats and food sources available.

Niche specialists do well when environmental conditions remain relatively constant; however loss of a favored habitat or food source leaves them with few alternatives for survival.

### Environmental changes affect distribution of species



Changes in tree species distributions over time. Pollen recovered from lake sediments indicates that plant species moved north as temperatures warmed following the retreat of the glaciers, beginning about 12,000 years ago. Areas shown in color or white were sampled for pollen, whereas areas shown in gray were not sampled.



**Figure 17.3** *Environmental Science for AP®*, Second Edition
Data from http://veimages.gsfc.nasa.gov//3453 /boreal\_model.gif

## **Species Extinction**




Environmental changes can cause species extinction...

If environmental conditions change, species that cannot adapt to the changes or move to more favorable environments will eventually go extinct.

The average lifespan of a species appears to be only about 1 million to 10 million years; 99 percent of the species that have ever lived on Earth are now extinct.

Mass extinction A large extinction of species in a relatively short period of time.

### **Five Global Mass Extinctions**



**Figure 17.6**Environmental Science for AP®, Second Edition
Data from GreenSpirit, http://www.greenspirit.org.uk/resources/TimeLines.jpg

Mass extinctions Five global mass extinction events have occurred since the evolution of complex life roughly 500 million years ago.