## Mesh Model Equations Cheat Sheet

### Overview

This cheat sheet summarizes key equations from the Mesh Model framework, covering discrete mesh mechanics, quantum field emergence, and geometric structure. It reflects the core pipeline developed in the Mesh-Field Transformer and its alignment with both classical and quantum field theory.

### Legend

- $q_i$  Generalized coordinate
- $\dot{q}_i$  Time derivative of  $q_i$
- T Kinetic energy
- V Potential energy
- m Mass
- R Radial or rotational coordinate
- $\theta$  Angular position
- G Gravitational constant
- M Mass of a black hole or object
- r Radial coordinate
- c Speed of light
- $\rho(x)$  Energy density at position x
  - ħ Reduced Planck's constant
- $\omega$  Angular frequency
- $\hat{H}$  Hamiltonian operator
- $\hat{p}$  Momentum operator
- $\hat{x}$  Position operator
- $E_n$  Energy of the *n*-th quantum state
- $\phi_i$  Field value at node i
- $\phi(x)$  Continuous field approximation
- $\pi(x)$  Canonical momentum density
- a Lattice spacing
- $\mathcal{L}$  Lagrangian density
- ds Line element in spacetime
- f(x) Emergent metric factor
- S Action
- $\Delta S$  Change in action across a curvature gradient

### 1 Lagrangian Mechanics

### 1.1 Generalized Coordinates

$$L(q_i, \dot{q}_i, t) = T - V$$

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0$$

#### 1.2 Particle on a Circle

$$L = \frac{1}{2}mR^2\dot{\theta}^2$$

### 2 General Relativity Integration

### 2.1 Schwarzschild Metric

$$ds^{2} = -\left(1 - \frac{2GM}{r}\right)c^{2}dt^{2} + \left(1 - \frac{2GM}{r}\right)^{-1}dr^{2} + r^{2}d\Omega^{2}$$

#### 2.2 Mesh-Emergent Metric

$$ds^{2} = -f(x)^{2}dt^{2} + dx^{2}, \quad f(x) = \sqrt{1 - \frac{2G\rho(x)}{c^{2}}}$$

### 3 Quantum Mechanics

### 3.1 Quantum Harmonic Oscillator

$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}k\hat{x}^2$$

$$E_n = \hbar\omega \left( n + \frac{1}{2} \right)$$

### 4 Field Theory on a Mesh

### 4.1 Discrete Scalar Field (Lattice)

$$L = \sum_{i} \left[ \frac{1}{2} \dot{\phi}_{i}^{2} - \frac{1}{2a^{2}} (\phi_{i+1} - \phi_{i})^{2} \right]$$

4.2 Continuum Limit (Wave Equation)

$$m\frac{\partial^2 \phi}{\partial t^2} = k\frac{\partial^2 \phi}{\partial x^2}$$

4.3 Lagrangian Density (Continuum)

$$\mathcal{L} = \frac{1}{2}m\left(\frac{\partial\phi}{\partial t}\right)^2 - \frac{1}{2}k\left(\frac{\partial\phi}{\partial x}\right)^2$$

4.4 Canonical Quantization

$$[\hat{\phi}(x), \hat{\pi}(y)] = i\hbar\delta(x-y)$$

- 5 Mesh Drive and Coherence Gradient
- 5.1 Curvature Propulsion

$$\Delta S = S_{\text{front}} - S_{\text{rear}} < 0$$

Motion arises from asymmetry in the mesh curvature gradient ion: Directional causal availability

- 6 Core Equations of the Mesh Model
- 6.1 1. Coupled Lagrangian (Structure + Curvature + Coherence)

$$\begin{split} \mathcal{L} &= -T_0 \sqrt{1 - \frac{1}{T_0} g^{\mu\nu} \partial_{\mu} \phi \, \partial_{\nu} \phi} - V(\phi) \\ &\quad + \frac{1}{2} K^{\mu\nu\alpha\beta} \nabla_{\mu} h_{\alpha\beta} \nabla_{\nu} h^{\alpha\beta} \\ &\quad - \frac{1}{2 \kappa} \phi^2 g^{\mu\nu} h_{\mu\nu} \end{split}$$

**Tagline:** "Spacetime emerges when coherence meets resistance."

6.2 2. Mesh–Field Transformer (Curved-Compatible)

$$\phi(x) = \sum_{i} \phi_{i} \, \psi_{i}(x), \quad \text{with}$$

$$\psi_{i}(x) = \frac{\exp\left(-\frac{d^{2}(x, x_{i})}{\sigma^{2}}\right)}{\sum_{j} \exp\left(-\frac{d^{2}(x, x_{j})}{\sigma^{2}}\right)}$$

$$V_{\rm field} = \int \frac{1}{2} k \, g^{\mu\nu}(x) \, \partial_{\mu} \phi(x) \, \partial_{\nu} \phi(x) \, \sqrt{-g(x)} \, d^4x$$

**Tagline:** "Structure becomes field — the mesh becomes physics."

6.3 3. Inversion Equation (Structure Defines Geometry)

$$g^{\mu\nu}(x) \propto \frac{1}{\mathcal{E}(x)} \sum_{i,j} \phi_i \phi_j \, \partial^{\mu} \psi_i(x) \, \partial^{\nu} \psi_j(x)$$

**Tagline:** "Curvature is structure. Geometry is earned."

- 7 Causal Structure and Collapse Dynamics
- 7.1 1. Coherence Vector Field

$$\vec{C}(x,t) = \nabla \phi(x,t) \cdot \chi(x,t)$$

from local phase gradient and coherence support.

7.2 2. Tension-Dependent Propagation Velocity

$$v^{2}(x) = \frac{T(x)}{\mu(x)} \quad \Rightarrow \quad \vec{v}(x) = \sqrt{\frac{T_{ij}(x)}{\mu}} \cdot \hat{n}$$

**Interpretation:** Anisotropic signal velocity determined by mesh stiffness and density.

7.3 3. Curvature Resistance (Accumulated Coherence Loss)

$$\mathcal{R}(x) = \int_{\gamma} (1 - \chi(x(s))) \, ds$$

**Interpretation:** Pathwise delay and bending from coherence failure — analog of gravitational curvature.

7.4 4. Effective Causal Cone

$$\operatorname{Cone}_{\operatorname{eff}}(x) = f\left(\vec{C}(x), \vec{v}(x), \mathcal{R}(x)\right)$$

**Interpretation:** Emergent light cone from composite geometry of field structure.

### 7.5 5. Causal Transport Equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \vec{v}) + \rho \Gamma(x) = 0$$
$$\Gamma(x) = \nabla \cdot \vec{C}(x)$$

**Interpretation:** General transport law governing ripple propagation, coherence collapse, and decoherence.

# 7.6 6. Effective Mass from Collapse and Resistance

$$m_{\rm eff}^2(x) \propto \Gamma(x) + \mathcal{R}(x)$$

**Interpretation:** Mass emerges structurally from coherence divergence and accumulated resistance.

# 7.7 7. Entropy Bound from Coherence Divergence

$$S_{\max} \leq \frac{1}{4} \int_{\Sigma} \left| \nabla \cdot \vec{C}(x) \right| \, dA$$

**Interpretation:** A structural analog of the Bousso bound based on ripple support loss.

# 7.8 8. Decay Probability from Divergence

$$P(t) = 1 - \exp\left(-\int \Gamma(x(t)) dt\right)$$

**Interpretation:** Structural analog of exponential decay from coherence divergence.

#### 7.9 9. Horizonless Tunneling Rate

$$\Gamma \sim \exp\left(-\frac{\Delta \mathcal{R}}{\hbar}\right)$$

Interpretation: Black hole radiation as ripple tunneling through steep coherence gradient.

# 7.10 10. Interference Region (Double Slit Geometry)

$$\mathcal{I}(x) = \left\{ x \mid \vec{C}_L(x) \cdot \vec{C}_R(x) > 0, \quad \mathcal{R}(x) < \infty \right\}$$

**Interpretation:** Interference only forms in regions with overlapping coherence and finite resistance.

# 7.11 11. Commutator from Coherence Misalignment

$$[\vec{C}^a, \vec{C}^b] := f^{abc}\vec{C}^c, \quad \text{where} \quad f^{abc}(x) \propto \epsilon^{\mu\nu} \partial_{\mu} \chi^a \, \partial_{\nu} \chi^b$$

**Interpretation:** Effective SU(N)-like structure from interference of coherence gradients.

# 8 Particle Structure and Coherence Algebra

### 8.1 Spin- $\frac{1}{2}$ Phase Winding

$$\Psi(x) = e^{i\theta(x)/2} \quad \Rightarrow \quad \Psi \to -\Psi \text{ under } \oint \nabla \theta \cdot d\ell = 2\pi$$

Interpretation: Spin- $\frac{1}{2}$  behavior from double-valued coherence phase geometry.

### 8.2 Chiral Collapse Asymmetry

$$\Delta\Gamma^a(x) = \Gamma_L^a(x) - \Gamma_R^a(x)$$

Interpretation: Structural origin of chirality from directional coherence collapse.

### 8.3 Quark Fractional Charge

$$Q^{a} = \frac{1}{2\pi} \oint \nabla \theta^{a}(x) \cdot d\ell = \frac{n_{a}}{k_{a}}$$

Interpretation: Fractional charge from coherence winding density.

### 8.4 Gluon Field Strength Tensor

$$\mathcal{F}^{ab}_{\mu\nu} = \partial_{\mu}C^{a}_{\nu} - \partial_{\nu}C^{a}_{\mu} + f^{abc}C^{b}_{\mu}C^{c}_{\nu}$$

Interpretation: SU(3)-like curvature from coherence cone interactions.

#### 8.5 Gluon Field Evolution

$$\nabla^{\mu}\mathcal{F}^{ab}_{\mu\nu} = J^{b}_{\nu}(x) \quad \text{with} \quad J^{a}_{\nu} = \phi^{b}\partial_{\nu}\phi^{c}f^{abc}$$

Interpretation: Gluon transport sourced by quark coherence mode transitions.