

Global Medical Affairs Cover Sheet

Asset Name	Budesonide/Formoterol-Novel START Medical Reactive Slide Deck		
Intended Use	Reactive		
Approved for Distribution	Yes, pending local market approval	Therapy Area	Respiratory

Asset Owner	Norbert Feigler/Leslee Falls		New or Updated	Updated
Document Number	ML-3010-ALL-0123		Based on Asset <i>(include Vault MedComms or PromoMats document number or N/A for not applicable)</i>	N/A
Approved Date	05/19	Expiration Date	05/20	
Technical Review/Fact Check Completed by Global Medical Information <i>(Yes/No)</i>		Yes	Applicable Copyright Permissions Obtained for Graphics <i>(Yes/No)</i>	Yes

This material is globally approved for use by AstraZeneca Medical Affairs Scientific Personnel only. The local market is responsible for interpreting, reviewing, and approving the content according to their local label, rules, and regulations.

AstraZeneca does not, under any circumstances, promote its products for off-label or unapproved uses.

This slide deck is for verbal, reactive use only.

This slide deck is for verbal, reactive use only. Use of any of the content must be approved by your local nominated signatory/local medical process to ensure compliance with local regulations.

Global Medical Affairs

Novel Symbicort Turbuhaler Asthma Reliever Therapy (START) – Study Overview

Veeva Vault MedComms Document Number: ML-3010-ALL-0123

Approval Date: 05/19

Expiration Date: 05/20

UNCONTROLLED COPY

Clinical Programs Investigating Budesonide/Formoterol as an Anti-inflammatory Reliever in Mild Asthma

START Study Analysis: Mild patients benefit from early introduction and long-term ICS (budesonide)¹



SYGMA²

Novel START:

As-needed budesonide/formoterol in mild asthma³

To investigate if an open-label, clinical trial of as-needed budesonide/formoterol in adults previously treated with as-needed SABA only could extend the results from SYGMA to a real world setting

PRACTICAL:^a

An independent study
As-needed budesonide/formoterol^{4,5}


^aPRACTICAL is not an AstraZeneca study.

ICS = inhaled corticosteroid; Novel START = Novel Symbicort Turbuhaler Asthma Reliever Therapy; PRACTICAL = PeRsonalised Asthma Combination Therapy: with Inhaled Corticosteroid And fast-onset Long-acting beta agonist; SABA = short-acting beta₂-agonist; START = Steroid Treatment As Regular Therapy; SYGMA = SYmbicort Given as needed in Mild Asthma.

1. Reddel HK et al. *Lancet*. 2017;389:157-166; 2. O'Byrne PM et al. *Trials*. 2017;18:12. <https://doi.org/10.1186/s13063-016-1731-4>. Accessed March 4, 2019; 3. 2. Beasley R et al. *Eur Respir J*. 2016;47:981-984.; 4. Study ACTRN12616000377437. Australian New Zealand Clinical Trials Registry website. <https://www.anzctr.org.au/Trial/Registration/TrialReview.aspx?id=370122>. Accessed March 4, 2019; 5. Fingleton J et al. *BMJ Open Resp Res*. 2017;4:e000217. <https://bmjopenrespres.bmj.com/content/4/1/e000217>. Accessed March 7, 2019.

Study Design^a

12-month, **pragmatic**, randomized, open-label, parallel-group, multicenter study (N=675) to assess the long-term efficacy and safety of budesonide/formoterol anti-inflammatory reliever compared to SABA as-needed or ICS maintenance + SABA as-needed in patients with mild asthma **previously treated with SABA alone**^{1,2}



^aAnalyses were intention-to-treat; ^bResulting in medical care consultation and/or systemic glucocorticoids and/or high beta₂-agonist use; ^cPrescription of systemic glucocorticoids for ≥3 days for asthma and/or hospitalization or ED visit due to asthma leading to systemic glucocorticoids; ^dA total of three exacerbations, one severe exacerbation, or unstable asthma resulting in change in randomized treatment.

ACQ-5 = Asthma Control Questionnaire-5; DPI = dry powder inhaler; ED = emergency department; FE_{NO} = fraction of exhaled nitric oxide; FEV₁ = forced expiratory volume in 1 second; ICS = inhaled corticosteroid; OCS = oral corticosteroid; pMDI = pressurized metered-dose inhaler; SABA = short-acting β₂-agonist; SCE = screening, consent, enrollment.

1. Beasley R et al. *Eur Respir J.* 2016;47;981-984; 2. Beasley R et al. *N Engl J Med.* 2019;380:2020-2030.

Baseline Characteristics

	Albuterol 100 µg x 2 Inhalations As-Needed (n=223)	Budesonide/Formoterol 200/6 µg x 1 Inhalation As-Needed (n=220)	Budesonide Maintenance 200 µg x 1 Inhalation BID + Albuterol 100 µg x 2 Inhalations As-Needed (n=225)
Age, years, mean (SD)	35.8 (14.0)	36 (14.1)	34.9 (14.3)
Female sex, n (%)	113 (50.7)	122 (55.5)	129 (57.3)
Current smoker, ^a n (%)	24 (10.8)	18 (8.2)	22 (9.8)
Patient-reported SABA use in the 4 weeks prior to enrollment:			
Occasions per week, n, mean (SD)	3.4 (3.3)	3.8 (3.5)	3.2 (3.0)
Patients with ≤2 occasions per week, n (%)	127 (57.0)	105 (47.7)	132 (58.7)
≥1 severe exacerbations in the prior 12 months, ^b n (%)	20 (9.0)	12 (5.5)	17 (7.6)
ACQ-5 score, mean (SD)	1.1 (0.7)	1.1 (0.7)	1.1 (0.7)
On-treatment ^c FEV ₁ % predicted, mean (SD)	89.2 (13.7)	89.8 (14.1)	90.3 (13.6)
Median FE _{NO} (range), ppb	40 (5-235)	37 (3-300)	38 (5-200)
Blood eosinophils x 10 ⁻⁹ /L, mean (SD)	0.3 (0.2)	0.3 (0.2)	0.3 (0.2)

^aPatient-reported smoking history >20 pack-years or onset of respiratory symptoms after the age of 40 years in current or previous smokers with a smoking history of ≥10 pack-years was a key exclusion criterion; ^bHospitalization for asthma in the previous 12 months was a key exclusion criterion; ^cThere was no specific instruction for patients to withhold their bronchodilator use prior to the baseline measurement.

ACQ-5 = Asthma Control Questionnaire-5; FE_{NO} = fraction of exhaled nitric oxide; FEV₁ = forced expiratory volume in 1 second; SABA = short-acting β₂-agonist; SD = standard deviation.

Beasley R et al. *N Engl J Med*. 2019;380:2020-2030.

UNCONTROLLED COPY

Exacerbation Definitions

Primary Endpoint

Exacerbations:

Worsening asthma resulting in ≥ 1 of the following:

- **An urgent medical care consultation:**
 - Primary care visit
 - Emergency department visit
 - Hospital admission
- **A prescription for glucocorticoids for any duration**
- **An episode of high β_2 -agonist use:**
 - >16 actuations of albuterol within 24 hours
 - >8 actuations of budesonide/formoterol within 24 hours

Secondary Endpoint

Severe Exacerbations:


Based on ATS/ERS Task Force criteria

Worsening asthma leading to:


- **A prescription** for systemic glucocorticoid treatment for ≥ 3 days or
- **Hospitalization or emergency department visit** leading to a prescription for systemic glucocorticoid treatment

Exacerbation Outcomes

Annualized Exacerbation Rate¹
(Primary Outcome)



Number of Severe^a Exacerbations
(Secondary Outcome)^{1b}




^aFor severe exacerbations, the *relative risk* was estimated as opposed to the *relative rate*, as participants could only have 1 severe exacerbation after which they were withdrawn from the study;² ^bSecondary endpoints were not adjusted for multiplicity. RR = relative rate (left graph); RR = relative risk (right graph).
1. Beasley R et al. *N Engl J Med.* 2019;380:2020-2030; 2. Beasley R et al. Supplementary appendix. *N Engl J Med.* 2019. https://www.nejm.org/doi/suppl/10.1056/NEJMoa1901963/suppl_file/nejmoa1901963_appendix.pdf. Accessed 19 May, 2019.

UNCONTROLLED COPY

© AstraZeneca 2019


Number of Exacerbations By Exacerbation Treatment Criteria



^aSecondary endpoints were not adjusted for multiplicity.
Beasley R et al. *N Engl J Med.* 2019;380:2020-2030.

UNCONTROLLED COPY


First Exacerbation



^aSecondary endpoints were not adjusted for multiplicity.
BUD = budesonide; FORM = formoterol; HR = hazard ratio; Novel START = Symbicort Turbuhaler Asthma Reliever Therapy; SABA = short-acting β_2 -agonist.
Beasley R et al. *N Engl J Med.* 2019;380:2020-2030.

UNCONTROLLED COPY

Fewer Severe Exacerbations With >70% Lower Corticosteroid Load




A 56% reduction ($p=0.038$)^a in the number of severe exacerbations with budesonide/formoterol as-needed compared to budesonide maintenance plus SABA as-needed was achieved with a >70% lower corticosteroid load

^aSecondary endpoints were not adjusted for multiplicity.

IQR = interquartile range; Novel START = Novel Symbicort Turbuhaler Asthma Reliever Therapy.

Beasley R et al. *N Engl J Med*. 2019;380:2020-2030.

Fraction of Exhaled Nitric Oxide



Ratio of Geometric Mean FE_{NO} at 52 weeks:^a BUD/FORM as-needed vs. albuterol as-needed, 0.83 (95% CI, 0.75-0.91); BUD/FORM as-needed vs. BUD maintenance, 1.13 (95% CI, 1.02-1.25)

^aSecondary endpoints were not adjusted for multiplicity. BUD = budesonide; FE_{NO} = fraction of exhaled nitric oxide; FORM = formoterol; IQR = interquartile range; ppb = parts per billion.

Beasley R et al. *N Engl J Med.* 2019;380:2020-2030.

© AstraZeneca 2019

Safety and Tolerability

Patients, n (%)	Albuterol 100 µg x 2 Inhalations As-Needed (n=226)	Budesonide/Formoterol 200/6 µg x 1 Inhalation As-Needed (n=222)	Budesonide Maintenance 200 µg x 1 Inhalation BID + Albuterol 100 µg x 2 Inhalations As-Needed (n=227)
Patients with ≥1 AE	185 (81.9)	174 (78.4)	190 (83.7)
Most common AEs (occurring in ≥2% of patients)			
Upper respiratory tract infection	75 (33.2)	71 (32.0)	75 (33.0)
Viral upper respiratory tract infection	5 (2.2)	6 (2.7)	8 (3.5)
Asthma	46 (20.4)	17 (7.7)	26 (11.5)
Patients with ≥1 SAE (including death)	6 (2.7)	13 (5.9) ^a	7 (3.1) ^b

Adverse events were consistent with prior trials and with reports in clinical use

^aDeath was due to motor vehicle accident (n=1); ^bDeath was due to suicide (n=1).
AE = adverse event; BID = twice daily; SAE = serious adverse event.
Beasley R et al. *N Engl J Med.* 2019;380:2020-2030.

Conclusions

In an open-label study of patients with mild asthma who were previously treated with SABA as-needed, treatment with budesonide/formoterol as-needed resulted in the following:

- Compared to albuterol as-needed:
 - Reduced rate of any exacerbation by 51%
 - Reduced rate of severe exacerbations by 60%
- Compared to maintenance budesonide:
 - Comparable rate of any exacerbation
 - Reduced rate of severe exacerbations by 56% at a >70% lower steroid load