

Number Field Counting, L -Functions, and Automorphic Forms

Frank Thorne

University of South Carolina

Modular Forms and Related Topics, Beirut, May 2018

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\mathrm{Disc}(K)| < X\}.$$

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup $G \subseteq S_d$,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

Definition

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup $G \subseteq S_d$,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

so that

$$N_d(X) = \sum_{\substack{G \subseteq S_d \\ \text{transitive}}} N_d(X, G).$$

Theorem (Finiteness – Hermite)

For each d and X , $N_d(X)$ is finite.

Theorem (Finiteness – Hermite)

For each d and X , $N_d(X)$ is finite.

Theorem (Minkowski's Lower Bound)

If $[K : \mathbb{Q}] = d$, then

$$|\mathrm{Disc}(K)| \geq \left(\frac{d^d}{d!}\right)^2 \left(\frac{\pi}{4}\right)^d.$$

Theorem (Finiteness – Hermite)

For each d and X , $N_d(X)$ is finite.

Theorem (Minkowski's Lower Bound)

If $[K : \mathbb{Q}] = d$, then

$$|\mathrm{Disc}(K)| \geq \left(\frac{d^d}{d!}\right)^2 \left(\frac{\pi}{4}\right)^d.$$

In other words,

$$N_d(X) = 0 \quad \text{for } X < (5.803 \cdots + o(1))^d.$$

The Inverse Galois Problem

Conjecture

For every d and transitive subgroup $G \subseteq S_d$,

$$X \text{ big enough} \implies N_d(X, G) \neq 0.$$

The Inverse Galois Problem

Conjecture

For every d and transitive subgroup $G \subseteq S_d$,

$$X \text{ big enough} \implies N_d(X, G) \neq 0.$$

Proof.

????????



Three Methods to Count Number Fields

- ▶ Polynomial methods (upper and lower bounds);

Three Methods to Count Number Fields

- ▶ Polynomial methods (upper and lower bounds);
- ▶ Abelian methods (upper bounds and asymptotics);

Three Methods to Count Number Fields

- ▶ Polynomial methods (upper and lower bounds);
- ▶ Abelian methods (upper bounds and asymptotics);
- ▶ Parametrization methods (asymptotics in limited cases).

If $\alpha \in \mathcal{O}_K$ is a generator of K/\mathbb{Q} , then $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$ and

$$\begin{aligned} |\mathrm{Disc}(\mathcal{O}_K)| &= \mathrm{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \mathrm{Disc}(\min_{\alpha}(X)) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

If $\alpha \in \mathcal{O}_K$ is a generator of K/\mathbb{Q} , then $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$ and

$$\begin{aligned} |\mathrm{Disc}(\mathcal{O}_K)| &= \mathrm{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \mathrm{Disc}(\min_{\alpha}(X)) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

Theorem (Schmidt)

For each d we have

$$N_d(X) \ll X^{\frac{d+2}{4}}.$$

If $\alpha \in \mathcal{O}_K$ is a generator of K/\mathbb{Q} , then $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$ and

$$\begin{aligned} |\mathrm{Disc}(\mathcal{O}_K)| &= \mathrm{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \mathrm{Disc}(\min_{\alpha}(X)) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

Theorem (Schmidt)

For each d we have

$$N_d(X) \ll X^{\frac{d+2}{4}}.$$

See Ellenberg-Venkatesh (2006), Dummit (2017) for refinements.

Given a base number field K .

Given a base number field K .

Theorem (Class Field Theory)

Abelian extensions L/K of degree d and conductor dividing \mathfrak{m} are in bijection with index d quotients of $\text{Cl}_{\mathfrak{m}}(L)$.

Given a base number field K .

Theorem (Class Field Theory)

Abelian extensions L/K of degree d and conductor dividing \mathfrak{m} are in bijection with index d quotients of $\text{Cl}_{\mathfrak{m}}(L)$.

Theorem (Kummer Theory)

If in addition $\mu_d \subseteq K$, then abelian extensions L/K of exponent d are in bijection with subgroups of $K^{\times}/(K^{\times})^d$.

Sample theorem

Definition


If K is an S_3 -cubic field, its *quadratic resolvent* is $\mathbb{Q}(\sqrt{\text{Disc}(K)})$, the unique quadratic subfield of K^c .

Theorem (Cohen, Morra, T.)

Let $D \neq 0, 1$ be a fundamental discriminant. Then,

$$\sum_{\substack{[K:\mathbb{Q}]=3 \\ \mathbb{Q}(\sqrt{D}) \text{ is the} \\ \text{quadratic resolvent of } K}} |\text{Disc}(K)| = \{\text{explicit finite sum of Euler products}\}.$$

The parametrization method



A sample theorem

Theorem (Bhargava, *Annals*, 2004)

There exists an explicit, discriminant preserving bijection between the following two sets:

Theorem (Bhargava, *Annals*, 2004)

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $\mathrm{GL}_3(\mathbb{Z}) \times \mathrm{GL}_2(\mathbb{Z})$ -orbits on the lattice $(\mathrm{Sym}^2 \mathbb{Z}^3 \otimes \mathbb{Z}^2)$ of pairs of integral ternary quartic forms.

Theorem (Bhargava, *Annals*, 2004)

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $\mathrm{GL}_3(\mathbb{Z}) \times \mathrm{GL}_2(\mathbb{Z})$ -orbits on the lattice $(\mathrm{Sym}^2 \mathbb{Z}^3 \otimes \mathbb{Z}^2)$ of pairs of integral ternary quartic forms.
- ▶ Pairs (Q, R) , where Q is a quartic ring and R is a cubic resolvent of Q .

A Bhargava-style metatheorem

Theorem

There exists an explicit, discriminant preserving bijection between the following two sets:

A Bhargava-style metatheorem

Theorem

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ *$G(\mathbb{Z})$ -orbits on a lattice $V(\mathbb{Z})$; where G is an **algebraic group** acting (often **prehomogeneously**) on a vector space V ;*

A Bhargava-style metatheorem

Theorem

There exists an explicit, discriminant preserving bijection between the following two sets:

- ▶ $G(\mathbb{Z})$ -orbits on a lattice $V(\mathbb{Z})$; where G is an **algebraic group** acting (often **prehomogeneously**) on a vector space V ;
- ▶ Some nice class of arithmetic objects we want to count.

How to prove Bhargava-style theorems

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration.

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)
- ▶ Pick your favorite complex representation (G, V) (which should be defined over \mathbb{Z} , and for which the invariant theory should be nice).

How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)
- ▶ Pick your favorite complex representation (G, V) (which should be defined over \mathbb{Z} , and for which the invariant theory should be nice).
- ▶ Try to prove that the $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ parametrize something. Hope to get lucky.

An Arithmetic Statistics Theorem

Theorem (Davenport-Heilbronn 1971 + BBPSTTT*)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3+\epsilon}).$$

An Arithmetic Statistics Theorem

Theorem (Davenport-Heilbronn 1971 + BBPSTTT*)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3+\epsilon}).$$

*: Belabas (1999), Belabas, Bhargava, Pomerance (~2006), Bhargava, Shankar, Tsimerman (2010), Taniguchi, T. (2011), Bhargava, Taniguchi, T. (2018? hopefully??)

Three Connections to L -Functions and Automorphic Forms

To be (briefly) discussed today:

- ▶ The Colmez conjecture;
- ▶ Automorphy of Sato-Shintani zeta functions;
- ▶ Equidistribution of shapes of number fields.

Connection 1: The Colmez Conjecture

(work with [Adrian Barquero-Sanchez](#) and [Riad Masri](#))

The Main Theorem

Theorem

Assume a weak form of [Malle's Conjecture](#).

Then, [the Colmez conjecture](#) is true for 100% of [CM fields](#) of any fixed degree, when ordered by discriminant.

The Colmez Conjecture

Conjecture (Colmez '93)

Let E be a CM field, and let X_Φ be a CM abelian variety of type (\mathcal{O}_E, Φ) . Then,

$$h_{\text{Fal}}(X_\Phi) = -Z(A_{E,\Phi}^0, 0) - \frac{1}{2}\mu_{\text{Art}}(A_{E,\Phi}^0),$$

where in the above

The Colmez Conjecture

Conjecture (Colmez '93)

Let E be a CM field, and let X_Φ be a CM abelian variety of type (\mathcal{O}_E, Φ) . Then,

$$h_{\text{Fal}}(X_\Phi) = -Z(A_{E,\Phi}^0, 0) - \frac{1}{2}\mu_{\text{Art}}(A_{E,\Phi}^0),$$

where in the above

- ▶ X_Φ is an **abelian variety** over $\overline{\mathbb{Q}}$ with **complex multiplication** by \mathcal{O}_E , and **CM type** for E (....)
- ▶ $h_{\text{Fal}}(X_\Phi)$ is the **Faltings height** of X_Φ , which in fact only depends on Φ ;
- ▶ The quantity on the right is defined in terms of logarithmic derivatives of Artin L -functions associated to characters defined in terms of the representation theory of $\text{Gal}(\mathbb{Q}^{\text{CM}}/\mathbb{Q})$.
- ▶ I could explain all this in more detail, but the margins of these slides are too small.

Suppose we have the following:

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);
- ▶ Then $\text{Gal}(E^c/\mathbb{Q}) \subseteq C_2 \wr G$, where

$$C_2 \wr G := C_2^d \rtimes G.$$

G -Weyl CM Fields

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);
- ▶ Then $\text{Gal}(E^c/\mathbb{Q}) \subseteq C_2 \wr G$, where

$$C_2 \wr G := C_2^d \rtimes G.$$

Definition

If $\text{Gal}(E^c/\mathbb{Q}) = C_2 \wr G$, we call E a G -Weyl CM field.

G-Weyl CM Fields

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);
- ▶ Then $\text{Gal}(E^c/\mathbb{Q}) \subseteq C_2 \wr G$, where

$$C_2 \wr G := C_2^d \rtimes G.$$

Definition

If $\text{Gal}(E^c/\mathbb{Q}) = C_2 \wr G$, we call E a G -Weyl CM field.

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Fields Satisfying the Colmez Conjecture

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Follows without too much difficulty from:

Fields Satisfying the Colmez Conjecture

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Follows without too much difficulty from:

- ▶ An **averaged version** of the Colmez conjecture, proved (independently) by Andreatta, Goren, Howard, and Madapusi Pera; and by Yuan-Zhang;

Fields Satisfying the Colmez Conjecture

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Follows without too much difficulty from:

- ▶ An **averaged version** of the Colmez conjecture, proved (independently) by Andreatta, Goren, Howard, and Madapusi Pera; and by Yuan-Zhang;
- ▶ Previous work of my coauthors.

A reduction of the main theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, 100% of CM fields of any fixed degree are G -Weyl CM fields, when ordered by discriminant.

A reduction of the main theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, 100% of CM fields of any fixed degree are G -Weyl CM fields, when ordered by discriminant.

That is: 100% of CM fields have Galois group as big as it can be.

The Weak Malle Conjecture for (d, G)

Choose quantities $M(G)$ and δ_G so that:

The Weak Malle Conjecture for (d, G)

Choose quantities $M(G)$ and δ_G so that:

- ▶ We have

$$N_d(X, G) \ll_G X^{M(G)}.$$

$M(G) \leq 1$ conjectured for each G ; proved in various cases.

The Weak Malle Conjecture for (d, G)

Choose quantities $M(G)$ and δ_G so that:

- ▶ We have

$$N_d(X, G) \ll_G X^{M(G)}.$$

$M(G) \leq 1$ conjectured for each G ; proved in various cases.

- ▶ For each K counted by $N_d(X, G)$ we have

$$|\mathrm{Cl}(K)[2]| \ll_{\epsilon, G} |\mathrm{Disc}(K)|^{\delta_G + \epsilon}.$$

$\delta_G = 0$ is believed; $\delta_G = \frac{1}{2}$ is known; sometimes can do better.

The Weak Malle Conjecture for (d, G)

Choose quantities $M(G)$ and δ_G so that:

- ▶ We have

$$N_d(X, G) \ll_G X^{M(G)}.$$

$M(G) \leq 1$ conjectured for each G ; proved in various cases.

- ▶ For each K counted by $N_d(X, G)$ we have

$$|\mathrm{Cl}(K)[2]| \ll_{\epsilon, G} |\mathrm{Disc}(K)|^{\delta_G + \epsilon}.$$

$\delta_G = 0$ is believed; $\delta_G = \frac{1}{2}$ is known; sometimes can do better.

Say the **Weak Malle Conjecture** holds for (d, G) if

$$M(G) + \delta_G < 2.$$

The Basic Strategy

- ▶ To count degree $2d$ CM fields E , count pairs (E, F) , where F is the maximal totally real subfield.

The Basic Strategy

- ▶ To count degree $2d$ CM fields E , count pairs (E, F) , where F is the maximal totally real subfield.
- ▶ Use transitivity of the discriminant:

$$\mathrm{Disc}(E) = \mathrm{Disc}(F)^2 \mathcal{N}_{F/\mathbb{Q}}(\mathfrak{d}(E/F)).$$

The Basic Strategy

- ▶ To count **degree $2d$ CM fields E** , count **pairs (E, F)** , where F is the maximal totally real subfield.
- ▶ Use **transitivity of the discriminant**:

$$\mathrm{Disc}(E) = \mathrm{Disc}(F)^2 \mathcal{N}_{F/\mathbb{Q}}(\mathfrak{d}(E/F)).$$

- ▶ Adapting work of Klüners, use the **abelian method** to count the number of F for each E .

Step 1: Counting quadratic extensions E/F

Define a Dirichlet series

$$D_{F,C_2}^-(s) := \sum_{[E:F]=2} \frac{1}{\mathcal{N}_{F/\mathbb{Q}}(\mathfrak{d}_{E/F})^s}$$

where the sum is over totally imaginary quadratic extensions E/F .

Counting quadratic extensions

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, \mathfrak{c}_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{\mathfrak{c}_\infty \subset \mathfrak{m}_\infty} \sum_{\mathfrak{c} | 2} \frac{(-1)^{|\mathfrak{c}_\infty|}}{2^{|\mathfrak{c}_\infty|}} \mathcal{N}_{F/\mathbb{Q}}(2/\mathfrak{c})^{1-2s} \sum_{\chi \in Q(\operatorname{Cl}_{\mathfrak{c}^2 \mathfrak{c}_\infty}(F))} L_F(\chi, s),$$

where \mathfrak{c} runs over all integral ideals of F dividing 2, \mathfrak{c}_∞ runs over all subsets of the set of real places \mathfrak{m}_∞ of F , χ runs over all quadratic characters $Q(\operatorname{Cl}_{\mathfrak{c}^2 \mathfrak{c}_\infty}(F))$ of the ray class group $\operatorname{Cl}_{\mathfrak{c}^2 \mathfrak{c}_\infty}(F)$ modulo $\mathfrak{c}^2 \mathfrak{c}_\infty$, and $L_F(\chi, s)$ is the L -function of χ .

Counting quadratic extensions: morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s).$$

Counting quadratic extensions: morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s).$$

We obtain

$$\#\{E : \mathcal{N}_{F/\mathbb{Q}}(\mathfrak{d}_{E/F}) \leq Y\} \asymp Y + \#\operatorname{Cl}(F)[2] \cdot o(Y),$$

Counting quadratic extensions: morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{\substack{c : \text{finite}}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s).$$

We obtain

$$\#\{E : \mathcal{N}_{F/\mathbb{Q}}(\mathfrak{d}_{E/F}) \leq Y\} \asymp Y + \#\operatorname{Cl}(F)[2] \cdot o(Y),$$

$$\sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F, C_2}(s)^{-s}.$$

Putting it all together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Putting it all together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

Putting it all together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

- ▶ $\xi(s)$ converges absolutely in $\Re(s) > 1$;

Putting it all together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

- ▶ $\xi(s)$ converges absolutely in $\Re(s) > 1$;
- ▶ $\xi(s)$ has meromorphic continuation to a half-plane $\Re(s) > \alpha$, with $\alpha < 1$; it has a simple pole at $s = 1$, with residue

$$\sum_F \frac{\operatorname{Res}_{s=1} \zeta_F(s)}{2^d d_F^2 \zeta_F(2)} \text{ “} \approx \text{” } \sum_F \frac{1}{d_F^2}.$$

Putting it all together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F, C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

- ▶ $\xi(s)$ converges absolutely in $\Re(s) > 1$;
- ▶ $\xi(s)$ has meromorphic continuation to a half-plane $\Re(s) > \alpha$, with $\alpha < 1$; it has a simple pole at $s = 1$, with residue

$$\sum_F \frac{\operatorname{Res}_{s=1} \zeta_F(s)}{2^d d_F^2 \zeta_F(2)} \text{ “} \approx \text{” } \sum_F \frac{1}{d_F^2}.$$

- ▶ $\xi(s)$ is polynomially bounded in vertical strips.

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension:

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

Lemma (Klüners)

Given the above, suppose that p is any prime unramified in F but ramified in E .

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

Lemma (Klüners)

Given the above, suppose that p is any prime unramified in F but ramified in E . Then $p^2 \mid d_L$.

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

Lemma (Klüners)

Given the above, suppose that p is any prime unramified in F but ramified in E . Then $p^2 \mid d_L$.

Application: Look at the Dirichlet series again: no contribution to the residue.

Table: Values of $r_d(G)$ for $d \leq 5$

d	G	Number of fields	Minimal discriminant	Residue	Proportion
2	C_2	100,000	5	0.009856	-
3		25,000	49	3.30×10^{-5}	-
	C_3	107	49	2.29×10^{-5}	0.69
	S_3	24,893	148	1.01×10^{-5}	0.31
4		25,000	725	1.24×10^{-7}	-
	C_4	75	1125	2.41×10^{-8}	0.19
	V_4	289	1600	1.56×10^{-8}	0.13
	D_4	8147	725	5.9×10^{-8}	0.48
	A_4	45	26569	9.3×10^{-11}	0.0008
	S_4	16,444	1957	2.5×10^{-8}	0.20
5		25,000	14641	1.05×10^{-10}	-
	C_5	5	14641	3.08×10^{-11}	0.29
	D_5	28	160801	4.24×10^{-13}	0.003
	F_5	15	2382032	9×10^{-15}	0.00009
	A_5	21	3104644	5×10^{-15}	0.00005
	S_5	24,931	24217	7.4×10^{-11}	0.70

Connection 2: Sato-Shintani Zeta Functions and Automorphy

Binary cubic forms I: Definitions

The lattice of *binary cubic forms* is

$$V(\mathbb{Z}) := \{au^3 + bu^2v + cuv^2 + dv^3 : a, b, c, d \in \mathbb{Z}\}.$$

Binary cubic forms I: Definitions

The lattice of *binary cubic forms* is

$$V(\mathbb{Z}) := \{au^3 + bu^2v + cuv^2 + dv^3 : a, b, c, d \in \mathbb{Z}\}.$$

There is an action of $GL(2)$, given by

$$(g \circ f)(u, v) = \frac{1}{\det g} f((u, v)g),$$

Binary cubic forms I: Definitions

The lattice of *binary cubic forms* is

$$V(\mathbb{Z}) := \{au^3 + bu^2v + cuv^2 + dv^3 : a, b, c, d \in \mathbb{Z}\}.$$

There is an action of $GL(2)$, given by

$$(g \circ f)(u, v) = \frac{1}{\det g} f((u, v)g),$$

which satisfies

$$\text{Disc}(g \circ f) = (\det g)^2 \text{Disc}(f),$$

$$\text{Disc}(f) = b^2c^2 - 4ac^3 - 4b^3d - 27a^2d^2 + 18abcd.$$

Binary cubic forms II: Parametrization

Theorem (Levi 1914, Delone-Faddeev 1940, Gan-Gross-Savin 2002)

*There is an explicit, discriminant-preserving bijection between the set of $\mathrm{GL}_2(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ and the set of **cubic rings**.*

A Sato-Shintani zeta function

Definition

The (*cubic*) *Sato-Shintani zeta function* associated to this (G, V) is

$$\begin{aligned}\xi^{\pm}(s) &:= \sum_{\substack{x \in \mathrm{GL}_2(\mathbb{Z}) \setminus V(\mathbb{Z}) \\ \pm \mathrm{Disc}(x) > 0}} \frac{1}{|\mathrm{Stab}(x)|} |\mathrm{Disc}(x)|^{-s} \\ &= \sum_{\pm \mathrm{Disc}(\mathcal{O}) > 0} |\mathrm{Disc}(\mathcal{O})|^{-s},\end{aligned}$$

where \mathcal{O} ranges over orders in cubic fields.

A Sato-Shintani zeta function

Definition

The (*cubic*) *Sato-Shintani zeta function* associated to this (G, V) is

$$\begin{aligned}\xi^{\pm}(s) &:= \sum_{\substack{x \in \mathrm{GL}_2(\mathbb{Z}) \setminus V(\mathbb{Z}) \\ \pm \mathrm{Disc}(x) > 0}} \frac{1}{|\mathrm{Stab}(x)|} |\mathrm{Disc}(x)|^{-s} \\ &= \sum_{\pm \mathrm{Disc}(\mathcal{O}) > 0} |\mathrm{Disc}(\mathcal{O})|^{-s},\end{aligned}$$

where \mathcal{O} ranges over orders in *étale cubic algebras*.

You can ignore this if it looks scary.

How to extract arithmetic density results

Let $\xi(s) := \sum_n a(n)n^{-s}$. Then **Perron's formula** states that

$$\sum_{n < X} a(n) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \xi(s) X^s \frac{ds}{s},$$

subject **(only)** to absolute convergence.

How to extract arithmetic density results

Let $\xi(s) := \sum_n a(n)n^{-s}$. Then **Perron's formula** states that

$$\sum_{n < X} a(n) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \xi(s) X^s \frac{ds}{s},$$

subject **(only)** to absolute convergence.

If $\xi(s)$ is a 'nice zeta function', we can hope for asymptotics with explicit error terms.

The functional equation

Theorem (Shintani, 1971)

The above zeta functions converge absolutely for $\Re(s) > 1$. They continue to functions holomorphic in the plane except for simple poles at 1 and $5/6$, and satisfy the functional equation

$$\begin{pmatrix} \xi^+(1-s) \\ \xi^-(1-s) \end{pmatrix} = \Gamma\left(s - \frac{1}{6}\right) \Gamma(s)^2 \Gamma\left(s + \frac{1}{6}\right) 2^{-1} 3^{6s-2} \pi^{-4s} \times \\ \begin{pmatrix} \sin 2\pi s & \sin \pi s \\ 3 \sin \pi s & \sin 2\pi s \end{pmatrix} \begin{pmatrix} \widehat{\xi}^+(s) \\ \widehat{\xi}^-(s) \end{pmatrix}.$$

Moreover,

$$\operatorname{Res}_{s=1} \xi^\pm(s) = \frac{\pi^2(3 + C^\pm)}{36}, \quad \operatorname{Res}_{s=5/6} \xi^\pm(s) = K^\pm \frac{\zeta(1/3) \Gamma(1/3)^3}{4\sqrt{3}\pi}.$$

Sato and Shintani's general theorem

Theorem (Sato-Shintani, 1974)

Let G be a *reductive group* acting *prehomogeneously* on a finite dimensional vector space V , with *irreducible singular set*.

Sato and Shintani's general theorem

Theorem (Sato-Shintani, 1974)

Let G be a *reductive group* acting *prehomogeneously* on a finite dimensional vector space V , with *irreducible singular set*.

Then, the associated Sato-Shintani zeta functions

$$\xi^{(i)}(s) := \sum_{\substack{x \in G(\mathbb{Z}) \setminus V^{(i)}(\mathbb{Z}) \\ \pm \text{Disc}(x) > 0}} \frac{1}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}$$

enjoy analytic continuations and functional equations.

Sato-Shintani: the fine print

$$\xi^{(i)}(s) := \sum_{x \in G(\mathbb{Z}) \setminus V^{(i)}(\mathbb{Z})} \frac{1}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}$$

$$\xi^{(i)}(s) := \sum_{x \in G(\mathbb{Z}) \setminus V^{(i)}(\mathbb{Z})} \frac{1}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}$$

- Here $V^{(i)}(\mathbb{Z}) := V(\mathbb{Z}) \cap V^{(i)}(\mathbb{R})$, for any of the **finitely many** nonsingular $G(\mathbb{R})$ -orbits on $V(\mathbb{R})$.

$$\xi^{(i)}(s) := \sum_{x \in G(\mathbb{Z}) \setminus V^{(i)}(\mathbb{Z})} \frac{1}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}$$

- ▶ Here $V^{(i)}(\mathbb{Z}) := V(\mathbb{Z}) \cap V^{(i)}(\mathbb{R})$, for any of the **finitely many** nonsingular $G(\mathbb{R})$ -orbits on $V(\mathbb{R})$.
- ▶ If $|\text{Stab}(x)|$ isn't finite, replace it with a **regulator**.

Sato-Shintani: the fine print

$$\xi^{(i)}(s) := \sum_{x \in G(\mathbb{Z}) \setminus V^{(i)}(\mathbb{Z})} \frac{1}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}$$

- ▶ Here $V^{(i)}(\mathbb{Z}) := V(\mathbb{Z}) \cap V^{(i)}(\mathbb{R})$, for any of the **finitely many** nonsingular $G(\mathbb{R})$ -orbits on $V(\mathbb{R})$.
- ▶ If $|\text{Stab}(x)|$ isn't finite, replace it with a **regulator**.
- ▶ Sato-Shintani don't give a recipe for writing down the **explicit** functional equation.

Sato-Shintani: the fine print

$$\xi^{(i)}(s) := \sum_{x \in G(\mathbb{Z}) \setminus V^{(i)}(\mathbb{Z})} \frac{1}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}$$

- ▶ Here $V^{(i)}(\mathbb{Z}) := V(\mathbb{Z}) \cap V^{(i)}(\mathbb{R})$, for any of the **finitely many** nonsingular $G(\mathbb{R})$ -orbits on $V(\mathbb{R})$.
- ▶ If $|\text{Stab}(x)|$ isn't finite, replace it with a **regulator**.
- ▶ Sato-Shintani don't give a recipe for writing down the **explicit** functional equation.
- ▶ The **residues** (and possibly even the **location and multiplicity of the poles**) are nontrivial to compute.

Question: Are Shintani zeta functions automorphic?

Question: Are Shintani zeta functions automorphic?

Ibukiyama and Saito (1995) considered the zeta functions associated to the space V of $n \times n$ symmetric matrices, with the action of $G = \mathrm{GL}(n)$

$$g \circ M = gMg^T.$$

Question: Are Shintani zeta functions automorphic?

Ibukiyama and Saito (1995) considered the zeta functions associated to the space V of $n \times n$ symmetric matrices, with the action of $G = \mathrm{GL}(n)$

$$g \circ M = gMg^T.$$

They proved:

Question: Are Shintani zeta functions automorphic?

Ibukiyama and Saito (1995) considered the zeta functions associated to the space V of $n \times n$ symmetric matrices, with the action of $G = \mathrm{GL}(n)$

$$g \circ M = gMg^T.$$

They proved:

- ▶ An explicit description of these zeta functions: “They are essentially sums of two Dirichlet series, which are products of shifted Riemann zeta functions or of Dirichlet series attached to some Eisenstein series of one variable of half integral weight”.

Question: Are Shintani zeta functions automorphic?

Ibukiyama and Saito (1995) considered the zeta functions associated to the space V of $n \times n$ symmetric matrices, with the action of $G = \mathrm{GL}(n)$

$$g \circ M = gMg^T.$$

They proved:

- ▶ An explicit description of these zeta functions: “They are essentially sums of two Dirichlet series, which are products of shifted Riemann zeta functions or of Dirichlet series attached to some Eisenstein series of one variable of half integral weight”.
- ▶ Explicit determination of the functional equations, poles, and residues. (Also determined previously by Shintani.)

Question: Are Shintani zeta functions automorphic?

Ibukiyama and Saito (1995) considered the zeta functions associated to the space V of $n \times n$ symmetric matrices, with the action of $G = \mathrm{GL}(n)$

$$g \circ M = gMg^T.$$

They proved:

- ▶ An explicit description of these zeta functions: “They are essentially sums of two Dirichlet series, which are products of shifted Riemann zeta functions or of Dirichlet series attached to some Eisenstein series of one variable of half integral weight”.
- ▶ Explicit determination of the functional equations, poles, and residues. (Also determined previously by Shintani.)
- ▶ Much else besides.

A negative theorem

Theorem (T.)

*The Sato-Shintani zeta function associated to the space of **binary cubic forms** is **not** a **finite sum of Euler products**.*

The abelian method again

Theorem (Cohen, Morra, T.)

Let $D \neq 0, 1$ be a fundamental discriminant. Then,

$$\sum_{\substack{[K:\mathbb{Q}]=3 \\ \text{Disc}(K)=Dn^2}} |\text{Disc}(K)|^{-s} = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{2}{3^{2s}}\right) \prod_{\left(\frac{-3D}{p}\right)=1} \left(1 + \frac{2}{p^s}\right) +$$

$$\sum_{L \in \mathcal{L}_3(D)} M_{3,2,L}(s) \prod_{\left(\frac{-3D}{p}\right)=1} \left(1 + \frac{\omega_L(p)}{p^s}\right),$$

$$M_{3,2,L}(s) := \begin{cases} 1 - 3^{-2s} & : \text{Disc}(L) = -27D \\ 1 + 2 \cdot 3^{-2s} & : \text{Disc}(L) = -3D, \end{cases}$$

The abelian method again

Theorem (Cohen, Morra, T.)

Let $D \neq 0, 1$ be a fundamental discriminant. Then,

$$\sum_{\substack{[K:\mathbb{Q}]=3 \\ \text{Disc}(K)=Dn^2}} |\text{Disc}(K)|^{-s} = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{2}{3^{2s}}\right) \prod_{\left(\frac{-3D}{p}\right)=1} \left(1 + \frac{2}{p^s}\right) +$$

$$\sum_{L \in \mathcal{L}_3(D)} M_{3,2,L}(s) \prod_{\left(\frac{-3D}{p}\right)=1} \left(1 + \frac{\omega_L(p)}{p^s}\right),$$

$$M_{3,2,L}(s) := \begin{cases} 1 - 3^{-2s} & : \text{Disc}(L) = -27D \\ 1 + 2 \cdot 3^{-2s} & : \text{Disc}(L) = -3D, \end{cases}$$

- $\mathcal{L}_3(D)$ is the set of cubic fields of discriminant $-3D$ or $-27D$;

The abelian method again

Theorem (Cohen, Morra, T.)

Let $D \neq 0, 1$ be a fundamental discriminant. Then,

$$\sum_{\substack{[K:\mathbb{Q}]=3 \\ \text{Disc}(K)=Dn^2}} |\text{Disc}(K)|^{-s} = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{2}{3^{2s}}\right) \prod_{\left(\frac{-3D}{p}\right)=1} \left(1 + \frac{2}{p^s}\right) +$$

$$\sum_{L \in \mathcal{L}_3(D)} M_{3,2,L}(s) \prod_{\left(\frac{-3D}{p}\right)=1} \left(1 + \frac{\omega_L(p)}{p^s}\right),$$

$$M_{3,2,L}(s) := \begin{cases} 1 - 3^{-2s} & : \text{Disc}(L) = -27D \\ 1 + 2 \cdot 3^{-2s} & : \text{Disc}(L) = -3D, \end{cases}$$

- ▶ $\mathcal{L}_3(D)$ is the set of cubic fields of discriminant $-3D$ or $-27D$;
- ▶ $\omega_L(p) = 2$ if p splits completely in L , and $\omega_L(p) = -1$ otherwise.

Connection 3: Twisting Sato-Shintani Zeta Functions by
Automorphic Forms
(work of [Bob Hough](#))

Sato-Shintani zeta functions reimagined

The Sato-Shintani zeta function associated to the space of binary cubic forms:

$$\xi^{\pm}(s) := \sum_{\substack{x \in \mathrm{GL}_2(\mathbb{Z}) \setminus \mathcal{V}(\mathbb{Z}) \\ \pm \mathrm{Disc}(x) > 0}} \frac{1}{|\mathrm{Stab}(x)|} |\mathrm{Disc}(x)|^{-s}.$$

Sato-Shintani zeta functions reimagined

The Sato-Shintani zeta function associated to the space of binary cubic forms:

$$\xi^{\pm}(s) := \sum_{\substack{x \in \mathrm{GL}_2(\mathbb{Z}) \setminus V(\mathbb{Z}) \\ \pm \mathrm{Disc}(x) > 0}} \frac{1}{|\mathrm{Stab}(x)|} |\mathrm{Disc}(x)|^{-s}.$$

For simplicity, consider the positive zeta function only:

$$\xi^{+}(s) := \sum_{\substack{x \in \mathrm{GL}_2(\mathbb{Z}) \setminus V^{+}(\mathbb{Z}) \\ (\mathrm{Disc}(x) > 0)}} \frac{1}{|\mathrm{Stab}(x)|} |\mathrm{Disc}(x)|^{-s}.$$

The lattice reimagined

Choose (for example) $v_+ = uv(u + v) \in V^+(\mathbb{R})$.

The lattice reimagined

Choose (for example) $v_+ = uv(u + v) \in V^+(\mathbb{R})$. The map

$$\begin{aligned}\mathrm{GL}_2(\mathbb{R}) &\longrightarrow V^+(\mathbb{R}) \\ g &\longrightarrow g \cdot v_+\end{aligned}$$

is surjective and exactly 6-to-1.

The lattice reimagined

Choose (for example) $v_+ = uv(u + v) \in V^+(\mathbb{R})$. The map

$$\begin{aligned}\mathrm{GL}_2(\mathbb{R}) &\longrightarrow V^+(\mathbb{R}) \\ g &\longrightarrow g \cdot v_+\end{aligned}$$

is surjective and exactly 6-to-1.

Pretend for today: This map is a bijection.

Twisting by cusp forms

Let ϕ be a Maass cusp form on $SL_2(\mathbb{R})/SL_2(\mathbb{Z})$:

Twisting by cusp forms

Let ϕ be a Maass cusp form on $SL_2(\mathbb{R})/SL_2(\mathbb{Z})$:

- ▶ Extend ϕ to a function on $GL_2(\mathbb{R})/GL_2(\mathbb{Z})$, projecting by a diagonal matrix.

Twisting by cusp forms

Let ϕ be a Maass cusp form on $SL_2(\mathbb{R})/SL_2(\mathbb{Z})$:

- ▶ Extend ϕ to a function on $GL_2(\mathbb{R})/GL_2(\mathbb{Z})$, projecting by a diagonal matrix.
- ▶ Pushing forward, identify ϕ as a function on $GL_2(\mathbb{Z}) \backslash V^+(\mathbb{R})$.

Twisting by cusp forms

Let ϕ be a Maass cusp form on $SL_2(\mathbb{R})/SL_2(\mathbb{Z})$:

- ▶ Extend ϕ to a function on $GL_2(\mathbb{R})/GL_2(\mathbb{Z})$, projecting by a diagonal matrix.
- ▶ Pushing forward, identify ϕ as a function on $GL_2(\mathbb{Z}) \backslash V^+(\mathbb{R})$.
- ▶ Define the ϕ -twisted Shintani \mathcal{L} -function

$$\xi^+(s, \phi) := \sum_{x \in GL_2(\mathbb{Z}) \backslash V^+(\mathbb{Z})} \frac{\phi(x)}{|\text{Stab}(x)|} |\text{Disc}(x)|^{-s}.$$

Theorem (Hough)

*The function $\xi^+(s, \phi)$, **when properly defined**, has analytic continuation to $\Re(s) > \frac{1}{8}$.*

The shape of a number field

Given a number field K with real embeddings $\sigma_1, \dots, \sigma_r$ and complex embeddings $\tau_1, \tau_1', \dots, \tau_s, \tau_s'$, its **shape** is the quadratic form

$$q(x) := \sigma_1(x)^2 + \dots + \sigma_r(x)^2 + 2|\tau_1(x)|^2 + \dots + 2|\tau_s(x)|^2,$$

restricted to the $(n-1)$ -dimensional lattice

$$\{x \in \mathbb{Z} + n\mathcal{O}_K : \mathrm{Tr}(x) = 0\},$$

which we may consider as an element of

$$\mathrm{GL}_{n-1}(\mathbb{Z}) \backslash \mathrm{GL}_{n-1}(\mathbb{R}) / \mathrm{GO}_{n-1}(\mathbb{R}).$$

Equidistribution of shapes

Theorem (Terr, Bhargava-Harron)

Let $n = 3, 4, 5$. The shapes of number fields counted by $N_n(X, S_n)$ become equidistributed as $X \rightarrow \infty$.

Theorem (Hough)

Let ϕ be a cuspidal automorphic form on $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})$ (satisfying suitable technical hypotheses). Then, for any smooth, compactly supported test function $F : (0, \infty) \rightarrow \mathbb{R}$ we have

$$\sum_{[K:\mathbb{Q}]=3} \phi(\mathcal{O}_K) F\left(\frac{\pm \mathrm{Disc}(K)}{X}\right) \ll X^{3/4+\epsilon}.$$

Equidistribution of shapes

Theorem (Terr, Bhargava-Harron)

Let $n = 3, 4, 5$. The shapes of number fields counted by $N_n(X, S_n)$ become equidistributed as $X \rightarrow \infty$.

Theorem (Hough)

Let ϕ be a cuspidal automorphic form on $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})$ (satisfying suitable technical hypotheses). Then, for any smooth, compactly supported test function $F : (0, \infty) \rightarrow \mathbb{R}$ we have

$$\sum_{[K:\mathbb{Q}]=3} \phi(\mathcal{O}_K) F\left(\frac{\pm \mathrm{Disc}(K)}{X}\right) \ll X^{3/4+\epsilon}.$$

Equidistribution of shapes

Theorem (Terr, Bhargava-Harron)

Let $n = 3, 4, 5$. The shapes of number fields counted by $N_n(X, S_n)$ become equidistributed as $X \rightarrow \infty$.

Theorem (Hough)

Let ϕ be a cuspidal automorphic form on $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})$ (satisfying suitable technical hypotheses). Then, for any smooth, compactly supported test function $F : (0, \infty) \rightarrow \mathbb{R}$ we have

$$\sum_{[K:\mathbb{Q}]=3} \phi(\mathcal{O}_K) F\left(\frac{\pm \mathrm{Disc}(K)}{X}\right) \ll X^{3/4+\epsilon}.$$

- Recover shape equidistribution via spectral decomposition of $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})$.

Equidistribution of shapes

Theorem (Terr, Bhargava-Harron)

Let $n = 3, 4, 5$. The shapes of number fields counted by $N_n(X, S_n)$ become equidistributed as $X \rightarrow \infty$.

Theorem (Hough)

Let ϕ be a cuspidal automorphic form on $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})$ (satisfying suitable technical hypotheses). Then, for any smooth, compactly supported test function $F : (0, \infty) \rightarrow \mathbb{R}$ we have

$$\sum_{[K:\mathbb{Q}]=3} \phi(\mathcal{O}_K) F\left(\frac{\pm \mathrm{Disc}(K)}{X}\right) \ll X^{3/4+\epsilon}.$$

- ▶ Recover shape equidistribution via spectral decomposition of $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})$.
- ▶ Results for $n = 4$ too.