

The Distribution of G -Weyl CM Fields and the Colmez Conjecture

Frank Thorne (with Adrian Barquero-Sanchez and Riad Masri)

University of South Carolina

Automorphic Forms Workshop 2018

The Main Theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, the Colmez conjecture is true for 100% of CM fields of any fixed degree, when ordered by discriminant.

The Main Theorem

What do these words mean???

Theorem

Assume a weak form of *Malle's Conjecture*.

Then, *the Colmez conjecture* is true for 100% of *CM fields* of any fixed degree, when ordered by discriminant.

The Colmez Conjecture

Conjecture (Colmez '93)

Let E be a CM field, and let X_Φ be a CM abelian variety of type (\mathcal{O}_E, Φ) . Then,

$$h_{\text{Fal}}(X_\Phi) = -Z(A_{E,\Phi}^0, 0) - \frac{1}{2}\mu_{\text{Art}}(A_{E,\Phi}^0),$$

where in the above

The Colmez Conjecture

Conjecture (Colmez '93)

Let E be a CM field, and let X_Φ be a CM abelian variety of type (\mathcal{O}_E, Φ) . Then,

$$h_{\text{Fal}}(X_\Phi) = -Z(A_{E,\Phi}^0, 0) - \frac{1}{2}\mu_{\text{Art}}(A_{E,\Phi}^0),$$

where in the above

- ▶ X_Φ is an **abelian variety** over $\overline{\mathbb{Q}}$ with **complex multiplication** by \mathcal{O}_E , and **CM type** for E (....)
- ▶ $h_{\text{Fal}}(X_\Phi)$ is the **Faltings height** of X_Φ , which in fact only depends on Φ ;
- ▶ The quantity on the right is defined in terms of logarithmic derivatives of Artin L -functions associated to characters defined in terms of the representation theory of $\text{Gal}(\mathbb{Q}^{\text{CM}}/\mathbb{Q})$.
- ▶ I could explain all this in more detail, but the margins of these slides are too small.

Suppose we have the following:

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);
- ▶ Then $\text{Gal}(E^c/\mathbb{Q}) \subseteq C_2 \wr G$, where

$$C_2 \wr G := C_2^d \rtimes G.$$

G-Weyl CM Fields

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);
- ▶ Then $\text{Gal}(E^c/\mathbb{Q}) \subseteq C_2 \wr G$, where

$$C_2 \wr G := C_2^d \rtimes G.$$

Definition

If $\text{Gal}(E^c/\mathbb{Q}) = C_2 \wr G$, we call E a G -Weyl CM field.

G-Weyl CM Fields

Suppose we have the following:

- ▶ A *totally real* number field F of degree $[F : \mathbb{Q}] = d$;
- ▶ $G := \text{Gal}(F^c/\mathbb{Q})$ is a transitive subgroup of S_d ;
- ▶ A *totally imaginary* quadratic extension E/F (a **CM field**);
- ▶ Then $\text{Gal}(E^c/\mathbb{Q}) \subseteq C_2 \wr G$, where

$$C_2 \wr G := C_2^d \rtimes G.$$

Definition

If $\text{Gal}(E^c/\mathbb{Q}) = C_2 \wr G$, we call E a G -Weyl CM field.

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Fields Satisfying the Colmez Conjecture

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Follows without too much difficulty from:

Fields Satisfying the Colmez Conjecture

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Follows without too much difficulty from:

- ▶ An **averaged version** of the Colmez conjecture, proved (independently) by Andreatta, Goren, Howard, and Madapusi Pera; and by Yuan-Zhang;

Fields Satisfying the Colmez Conjecture

Theorem

If E is a G -Weyl CM field, then the Colmez conjecture is true for E .

Follows without too much difficulty from:

- ▶ An **averaged version** of the Colmez conjecture, proved (independently) by Andreatta, Goren, Howard, and Madapusi Pera; and by Yuan-Zhang;
- ▶ Previous work of my coauthors.

A reduction of the main theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, 100% of CM fields of any fixed degree are G -Weyl CM fields, when ordered by discriminant.

A reduction of the main theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, 100% of CM fields of any fixed degree are G -Weyl CM fields, when ordered by discriminant.

That is: 100% of CM fields have G -Weyl CM fields as big as it can be.

A reduction of the main theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, 100% of CM fields of any fixed degree are G -Weyl CM fields, when ordered by discriminant.

That is: 100% of CM fields have Galois group as big as it can be.

To count degree $2d$ CM fields E , count pairs (E, F) , where F is the maximal totally real subfield.

A reduction of the main theorem

Theorem

Assume a weak form of Malle's Conjecture.

Then, 100% of CM fields of any fixed degree are G -Weyl CM fields, when ordered by discriminant.

That is: 100% of CM fields have Galois group as big as it can be.

To count degree $2d$ CM fields E , count pairs (E, F) , where F is the maximal totally real subfield.

An adaptation of work of Klüners.

Step 1: Counting quadratic extensions E/F

Step 1: For each totally real F , enumerate all totally imaginary quadratic extensions E/F .

Step 1: Counting quadratic extensions E/F

Step 1: For each totally real F , enumerate all totally imaginary quadratic extensions E/F .

To do this, define a Dirichlet series

$$D_{F, C_2}^-(s) := \sum_{[E:F]=2} \frac{1}{\mathcal{N}_{F/\mathbb{Q}}(\mathfrak{D}_{E/F})^s}$$

where the sum is over all totally imaginary quadratic extensions E/F , and $\mathfrak{D}_{E/F}$ is the relative discriminant.

Counting quadratic extensions

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, \mathfrak{c}_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{\mathfrak{c}_\infty \subset \mathfrak{m}_\infty} \sum_{\mathfrak{c} | 2} \frac{(-1)^{|\mathfrak{c}_\infty|}}{2^{|\mathfrak{c}_\infty|}} \mathcal{N}_{F/\mathbb{Q}}(2/\mathfrak{c})^{1-2s} \sum_{\chi \in Q(\operatorname{Cl}_{\mathfrak{c}^2 \mathfrak{c}_\infty}(F))} L_F(\chi, s),$$

where \mathfrak{c} runs over all integral ideals of F dividing 2, \mathfrak{c}_∞ runs over all subsets of the set of real places \mathfrak{m}_∞ of F , χ runs over all quadratic characters $Q(\operatorname{Cl}_{\mathfrak{c}^2 \mathfrak{c}_\infty}(F))$ of the ray class group $\operatorname{Cl}_{\mathfrak{c}^2 \mathfrak{c}_\infty}(F)$ modulo $\mathfrak{c}^2 \mathfrak{c}_\infty$, and $L_F(\chi, s)$ is the L -function of χ .

Counting quadratic extensions, morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s),$$

where χ ranges (more or less) over *quadratic* characters of $\operatorname{Cl}(F)$.

Counting quadratic extensions, morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s),$$

where χ ranges (more or less) over *quadratic* characters of $\operatorname{Cl}(F)$.

What this says:

Counting quadratic extensions, morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s),$$

where χ ranges (more or less) over *quadratic* characters of $\operatorname{Cl}(F)$.

What this says:

- Wild ramification is annoying.

Counting quadratic extensions, morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s),$$

where χ ranges (more or less) over *quadratic* characters of $\operatorname{Cl}(F)$.

What this says:

- ▶ Wild ramification is annoying.
- ▶ There are $|\operatorname{Cl}(K)[2]|$ characters in the sum, and $|\operatorname{Cl}_{4\infty}(K)[2]|$ characters in the 'actual sum'.

Counting quadratic extensions, morally

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s),$$

where χ ranges (more or less) over *quadratic* characters of $\operatorname{Cl}(F)$.

What this says:

- ▶ Wild ramification is annoying.
- ▶ There are $|\operatorname{Cl}(K)[2]|$ characters in the sum, and $|\operatorname{Cl}_{4m_\infty}(K)[2]|$ characters in the 'actual sum'.
- ▶ The quadratic extensions ramified only at 2_∞ control the rest.

Counting quadratic extensions: the upshot

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s).$$

Counting quadratic extensions: the upshot

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s).$$

We obtain

$$\#\{E : [E : F] = 2, \text{ t.i.}, \mathcal{N}_{F/\mathbb{Q}}(\mathfrak{D}_{E/F}) \leq X\} \asymp X + \#\operatorname{Cl}(F)[2] \cdot o(X),$$

Counting quadratic extensions: the upshot

Theorem (Cohen-Diaz-Olivier '02)

For $\operatorname{Re}(s) > 1$ we have

$$D_{F, C_2}^-(s) = \frac{1}{\zeta_F(2s)} \sum_{c : \text{finite}} \left(\begin{array}{c} \text{some 2-adic} \\ \text{mumbo jumbo} \end{array} \right) \sum_{\chi} L_F(\chi, s).$$

We obtain

$$\#\{E : [E : F] = 2, \text{ t.i., } \mathcal{N}_{F/\mathbb{Q}}(\mathfrak{D}_{E/F}) \leq X\} \asymp X + \#\operatorname{Cl}(F)[2] \cdot o(X),$$

$$\sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F, C_2}(s)^{-s}.$$

Controlling the quadratic extensions

Definition (The constant δ_d)

For each d , choose $\delta_d \geq 0$ so that

$$|\mathrm{Cl}(K)[2]| \ll_{\epsilon, d} |d_K|^{\delta_d + \epsilon},$$

for all number fields K of degree d .

Controlling the quadratic extensions

Definition (The constant δ_d)

For each d , choose $\delta_d \geq 0$ so that

$$|\mathrm{Cl}(K)[2]| \ll_{\epsilon, d} |d_K|^{\delta_d + \epsilon},$$

for all number fields K of degree d .

- The 'trivial bound': $\delta_d = \frac{1}{2}$.

Controlling the quadratic extensions

Definition (The constant δ_d)

For each d , choose $\delta_d \geq 0$ so that

$$|\mathrm{Cl}(K)[2]| \ll_{\epsilon, d} |d_K|^{\delta_d + \epsilon},$$

for all number fields K of degree d .

- ▶ The 'trivial bound': $\delta_d = \frac{1}{2}$.
- ▶ The conjectured truth: $\delta_d = 0$.

Controlling the quadratic extensions

Definition (The constant δ_d)

For each d , choose $\delta_d \geq 0$ so that

$$|\mathrm{Cl}(K)[2]| \ll_{\epsilon, d} |d_K|^{\delta_d + \epsilon},$$

for all number fields K of degree d .

- ▶ The 'trivial bound': $\delta_d = \frac{1}{2}$.
- ▶ The conjectured truth: $\delta_d = 0$.
- ▶ A little better (BSTTTZ): $\delta_3 = .2784 \dots$ and $\delta_n = \frac{1}{2} - \frac{1}{2n}$ for $n \geq 4$.

Step 2: Counting the base fields

Notation: For $d \geq 1$ and $G \subseteq S_d$, write

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, \operatorname{Gal}(K^c/\mathbb{Q}) = G\}.$$

Step 2: Counting the base fields

Notation: For $d \geq 1$ and $G \subseteq S_d$, write

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, \operatorname{Gal}(K^c/\mathbb{Q}) = G\}.$$

Definition (The constant $M(G)$)

For each G , choose $M(G) > 0$ so that

$$N_d(X, G) \ll X^{M(G)}.$$

Step 2: Counting the base fields

Notation: For $d \geq 1$ and $G \subseteq S_d$, write

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, \operatorname{Gal}(K^c/\mathbb{Q}) = G\}.$$

Definition (The constant $M(G)$)

For each G , choose $M(G) > 0$ so that

$$N_d(X, G) \ll X^{M(G)}.$$

We can restrict the above to **totally real** K if we like, but this doesn't seem to help us.

Conjectures on $M(G)$

Conjectures:

Conjectures on $M(G)$

Conjectures:

- ▶ (Folklore/Linnik) $M(G) = 1$ always;

Conjectures on $M(G)$

Conjectures:

- ▶ (Folklore/Linnik) $M(G) = 1$ always;
- ▶ (Malle's Conjecture)

$$X^{a(G)} \ll N_d(X, G) \ll X^{a(G)+\epsilon},$$

for $a(G) \leq 1$ determined by the group theory of G .

Conjectures on $M(G)$

Conjectures:

- ▶ (Folklore/Linnik) $M(G) = 1$ always;
- ▶ (Malle's Conjecture)

$$X^{a(G)} \ll N_d(X, G) \ll X^{a(G)+\epsilon},$$

for $a(G) \leq 1$ determined by the group theory of G .

Conjecture (Our Weak Malle Conjecture)

We always have

$$M(G) + \delta_d < 2.$$

Conjectures on $M(G)$

Conjectures:

- ▶ (Folklore/Linnik) $M(G) = 1$ always;
- ▶ (Malle's Conjecture)

$$X^{a(G)} \ll N_d(X, G) \ll X^{a(G)+\epsilon},$$

for $a(G) \leq 1$ determined by the group theory of G .

Conjecture (Our Weak Malle Conjecture)

We always have

$$M(G) + \delta_d < 2.$$

Enough if $M(G) < \frac{3}{2} + \frac{1}{2n}$.

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);
- ▶ $d = \ell$ prime, $G = D_\ell$ (Klüners);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);
- ▶ $d = \ell$ prime, $G = D_\ell$ (Klüners);
- ▶ $d = 2|H|$, $G = C_2 \rtimes H$ in many cases (Klüners);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);
- ▶ $d = \ell$ prime, $G = D_\ell$ (Klüners);
- ▶ $d = 2|H|$, $G = C_2 \rtimes H$ in many cases (Klüners);
- ▶ $d = 3|A|, 4|A|$, $G = S_3 \times A, S_4 \times A$, A abelian (Wang);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);
- ▶ $d = \ell$ prime, $G = D_\ell$ (Klüners);
- ▶ $d = 2|H|$, $G = C_2 \rtimes H$ in many cases (Klüners);
- ▶ $d = 3|A|, 4|A|$, $G = S_3 \times A, S_4 \times A$, A abelian (Wang);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);
- ▶ $d = \ell$ prime, $G = D_\ell$ (Klüners);
- ▶ $d = 2|H|$, $G = C_2 \rtimes H$ in many cases (Klüners);
- ▶ $d = 3|A|, 4|A|$, $G = S_3 \times A, S_4 \times A$, A abelian (Wang);
- ▶ Any d , $|G| = d$ (Ellenberg-Venkatesh);
- ▶ Frobenius groups (Mehta);

Known Cases of Weak Malle

Theorem

The weak Malle conjecture is known for:

- ▶ $d \leq 5$, any G (Bhargava, Cohen-Diaz-Olivier);
- ▶ Any d , G abelian (Mäki);
- ▶ Any d , G a p -group (Klüners-Malle);
- ▶ $d = \ell$ prime, $G = D_\ell$ (Klüners);
- ▶ $d = 2|H|$, $G = C_2 \rtimes H$ in many cases (Klüners);
- ▶ $d = 3|A|, 4|A|$, $G = S_3 \times A, S_4 \times A$, A abelian (Wang);
- ▶ Any d , $|G| = d$ (Ellenberg-Venkatesh);
- ▶ Frobenius groups (Mehta);
- ▶ (....)

Step 3: Putting it together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Step 3: Putting it together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

Step 3: Putting it together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F,C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

- ▶ $\xi(s)$ converges absolutely in $\Re(s) > 1$;

Step 3: Putting it together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F, C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

- ▶ $\xi(s)$ converges absolutely in $\Re(s) > 1$;
- ▶ $\xi(s)$ has meromorphic continuation to a half-plane $\Re(s) > \alpha$, with $\alpha < 1$; it has a simple pole at $s = 1$, with residue

$$\sum_F \frac{\text{Res}_{s=1} \zeta_F(s)}{2^d d_F^2 \zeta_F(2)} \text{ " } \approx \text{ " } \sum_F \frac{1}{d_F^2}.$$

Step 3: Putting it together

Incorporating everything above, we have

$$\xi(s) := \sum_E d_E^{-s} = \sum_F \frac{1}{d_F^{2s}} D_{F, C_2}(s)^{-s}.$$

Proposition

If the Weak Malle Conjecture is true for **whatever family of fields we're summing F over**, then:

- ▶ $\xi(s)$ converges absolutely in $\Re(s) > 1$;
- ▶ $\xi(s)$ has meromorphic continuation to a half-plane $\Re(s) > \alpha$, with $\alpha < 1$; it has a simple pole at $s = 1$, with residue

$$\sum_F \frac{\operatorname{Res}_{s=1} \zeta_F(s)}{2^d d_F^2 \zeta_F(2)} \text{ " } \approx \text{ " } \sum_F \frac{1}{d_F^2}.$$

- ▶ $\xi(s)$ is polynomially bounded in vertical strips.

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension:

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

Lemma (Klüners)

Given the above, suppose that p is any prime unramified in F but ramified in E .

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

Lemma (Klüners)

Given the above, suppose that p is any prime unramified in F but ramified in E . Then $p^2 \mid d_L$.

The Significance of G -Weylness

Suppose that E is **not** a G -Weyl extension: $\text{Gal}(F^c/\mathbb{Q}) = G$ but $\text{Gal}(E^c/\mathbb{Q}) \neq C_2 \wr G$.

Lemma (Klüners)

Given the above, suppose that p is any prime unramified in F but ramified in E . Then $p^2 \mid d_L$.

Application: Look at the Dirichlet series again: no contribution to the residue.

Table: Values of $r_d(G)$ for $d \leq 5$

d	G	Number of fields	Minimal discriminant	Residue	Proportion
2	C_2	100,000	5	0.009856	-
3		25,000	49	3.30×10^{-5}	-
	C_3	107	49	2.29×10^{-5}	0.69
	S_3	24,893	148	1.01×10^{-5}	0.31
4		25,000	725	1.24×10^{-7}	-
	C_4	75	1125	2.41×10^{-8}	0.19
	V_4	289	1600	1.56×10^{-8}	0.13
	D_4	8147	725	5.9×10^{-8}	0.48
	A_4	45	26569	9.3×10^{-11}	0.0008
	S_4	16,444	1957	2.5×10^{-8}	0.20
5		25,000	14641	1.05×10^{-10}	-
	C_5	5	14641	3.08×10^{-11}	0.29
	D_5	28	160801	4.24×10^{-13}	0.003
	F_5	15	2382032	9×10^{-15}	0.00009
	A_5	21	3104644	5×10^{-15}	0.00005
	S_5	24,931	24217	7.4×10^{-11}	0.70