

# State High School Mathematics Tournament

University of South Carolina

Round 2 – April 22, 2023

# Tournament Round 2 – Rules

# Tournament Round 2 – Rules

- ▶ You will be asked a series of questions. Each correct answer earns you a point. First to three points wins!

# Tournament Round 2 – Rules

- ▶ You will be asked a series of questions. Each correct answer earns you a point. First to three points wins!
- ▶ You have 90 seconds to answer each question. The clock will be started once I've finished reading the question.

# Tournament Round 2 – Rules

- ▶ You will be asked a series of questions. Each correct answer earns you a point. First to three points wins!
- ▶ You have 90 seconds to answer each question. The clock will be started once I've finished reading the question.
- ▶ To answer, buzz in at any time. (You do not have to wait for me to finish reading the question.) This will stop the clock. If your answer is correct, you win the point and we move to the next question.

# Tournament Round 2 – Rules

- ▶ You will be asked a series of questions. Each correct answer earns you a point. First to three points wins!
- ▶ You have 90 seconds to answer each question. The clock will be started once I've finished reading the question.
- ▶ To answer, buzz in at any time. (You do not have to wait for me to finish reading the question.) This will stop the clock. If your answer is correct, you win the point and we move to the next question.
- ▶ If your answer is wrong, the clock will be restarted. If your opponent doesn't buzz in, they may answer *immediately* after time is called.

## Question 2-1

How many times does the graph of  $y = x^6 + 6x^4 + 11x^2 + 6$  cross the  $x$ -axis?

## Solution 2-1

**Answer.** 0.



## Solution 2-1

**Answer.** 0.

$$x^6 + 6x^4 + 11x^2 + 6 \geq 0 + 0 + 0 + 6$$

## Question 2-2

If you expand out  
  
and simplify, how many terms will the resulting polynomial have?

## Question 2-2

If you expand out

$$(x + y)^{10} + (x - y)^{10}$$

and simplify, how many terms will the resulting polynomial have?

**Answer. 6.**

**Answer.** 6.

$$(x + y)^{10} = x^{10} + 10x^9y + 55x^8y^2 + 120x^7y^3 + \cdots + y^{10},$$

**Answer.** 6.

$$(x + y)^{10} = x^{10} + 10x^9y + 55x^8y^2 + 120x^7y^3 + \cdots + y^{10},$$

$$(x - y)^{10} = x^{10} - 10x^9y + 55x^8y^2 - 120x^7y^3 + \cdots + y^{10}.$$

**Answer.** 6.

$$(x + y)^{10} = x^{10} + 10x^9y + 55x^8y^2 + 120x^7y^3 + \cdots + y^{10},$$

$$(x - y)^{10} = x^{10} - 10x^9y + 55x^8y^2 - 120x^7y^3 + \cdots + y^{10}.$$

The odd terms cancel and the even terms remain.

## Question 2-3

How many zeroes does  $2023!$  end in?



## Solution 2-3

**Answer.** 503.

**Answer.** 503.

**Solution.** The answer is the number of factors of 5 in  $2023!$ .

**Answer.** 503.

**Solution.** The answer is the number of factors of 5 in  $2023!$ .

►  $\lfloor \frac{2023}{5} \rfloor = 404$  integers  $n \leq 2023$  are divisible by 5.

**Answer.** 503.

**Solution.** The answer is the number of factors of 5 in  $2023!$ .

- ▶  $\lfloor \frac{2023}{5} \rfloor = 404$  integers  $n \leq 2023$  are divisible by 5.
- ▶ 80 integers  $n \leq 2023$  are divisible by  $5^2$ .

**Answer.** 503.

**Solution.** The answer is the number of factors of 5 in  $2023!$ .

- ▶  $\lfloor \frac{2023}{5} \rfloor = 404$  integers  $n \leq 2023$  are divisible by 5.
- ▶ 80 integers  $n \leq 2023$  are divisible by  $5^2$ .
- ▶ 16 integers  $n \leq 2023$  are divisible by  $5^3$ .

## Solution 2-3

**Answer.** 503.

**Solution.** The answer is the number of factors of 5 in  $2023!$ .

- ▶  $\lfloor \frac{2023}{5} \rfloor = 404$  integers  $n \leq 2023$  are divisible by 5.
- ▶ 80 integers  $n \leq 2023$  are divisible by  $5^2$ .
- ▶ 16 integers  $n \leq 2023$  are divisible by  $5^3$ .
- ▶ 3 integers  $n \leq 2023$  are divisible by  $5^4$ .

**Answer.** 503.

**Solution.** The answer is the number of factors of 5 in  $2023!$ .

- ▶  $\lfloor \frac{2023}{5} \rfloor = 404$  integers  $n \leq 2023$  are divisible by 5.
- ▶ 80 integers  $n \leq 2023$  are divisible by  $5^2$ .
- ▶ 16 integers  $n \leq 2023$  are divisible by  $5^3$ .
- ▶ 3 integers  $n \leq 2023$  are divisible by  $5^4$ .

$$404 + 80 + 16 + 3 = 503.$$

## Question 2-4

What is the sum of the real number solutions to  $x^6 - 7x^3 - 8 = 0$ ?



**Answer. 1.**

**Answer. 1.**

$$x^6 - 7x^3 - 8 = (x^3 - 8)(x^3 + 1)$$

**Answer. 1.**

$$x^6 - 7x^3 - 8 = (x^3 - 8)(x^3 + 1)$$

The two factors have unique roots  $x = 2$  and  $x = -1$  respectively.

## Question 2-5

	O	X
	X	
X		O

## Question 2-5

	O	X
	X	
X		O

The above shows a Tic-Tac-Toe board, where X has won after five moves.

## Question 2-5

	O	X
	X	
X		O

The above shows a Tic-Tac-Toe board, where X has won after five moves.

How many such Tic-Tac-Toe boards are there?

## Solution 2-5

**Answer.** 120.

**Answer.** 120.

- ▶ 8 possible configurations of Xs: three rows, three columns, two diagonals.



**Answer.** 120.

- ▶ 8 possible configurations of Xs: three rows, three columns, two diagonals.
- ▶ For each,  $\binom{6}{2} = 15$  ways to place the Os.

**Answer.** 120.

- ▶ 8 possible configurations of Xs: three rows, three columns, two diagonals.
- ▶ For each,  $\binom{6}{2} = 15$  ways to place the Os.
- ▶  $8 \times 15 = 120$ .

## Question 2-6

What is the last digit of  $2023^{2023}$ ?

**Answer. 7.**

**Answer.** 7.

**Solution.** The last digit of  $2023^{2023}$  equals the last digit of  $3^{2023}$ .

## Solution 2-6

**Answer.** 7.

**Solution.** The last digit of  $2023^{2023}$  equals the last digit of  $3^{2023}$ .

$3^4 = 81$  and  $2023 = 4 \cdot 505 + 3$ , so

## Solution 2-6

**Answer.** 7.

**Solution.** The last digit of  $2023^{2023}$  equals the last digit of  $3^{2023}$ .

$3^4 = 81$  and  $2023 = 4 \cdot 505 + 3$ , so

$$3^{2023} = 3^{4 \cdot 505 + 3} = (81)^{505} \cdot 3^3 = (\dots??1) \cdot 27,$$

which ends in 7.


## Question 7

If  $\triangle ABC$  is an isosceles triangle with  $AB = BC = 1$ , what should the length of  $AC$  be to maximize the triangle's area?



## Solution 7


**Answer.**  $\sqrt{2}$



$$\text{Area} = \sin(\theta) \cdot \cos(\theta) = \frac{1}{2} \sin(2\theta).$$

## Solution 7

**Answer.**  $\sqrt{2}$



$$\text{Area} = \sin(\theta) \cdot \cos(\theta) = \frac{1}{2} \sin(2\theta).$$


Maximize with  $\theta = \frac{\pi}{4}$ , so  $AC = \sqrt{2}$ .

## Question 8

The equation  $2^x = x^2$  has three real solutions. What is the nearest integer to their sum?

# Solution 8


**Answer. 5**



$x = 2$ ,  $x = 4$ , and  $x \approx -0.76 \dots$

# Solution 8

**Answer. 5**



$x = 2$ ,  $x = 4$ , and  $x = -1$ ...

For the negative solution, note that  $2^{-\frac{1}{2}} > (-\frac{1}{2})^2$ , so  $x < -\frac{1}{2}$ .

## Question 9

What is

$$1 - 2 + 3 - 4 + 5 - \cdots + 2021 - 2022 + 2023?$$

# Solution 9

**Answer.** 1012.

# Solution 9

**Answer.** 1012.

Write it as

$$(1-2)+(3-4)+\cdots+1012+\cdots+(-2020+2021)+(-2022+2023).$$

We have, e.g.,

$$1 - 2 - 2022 + 2023 = 0.$$



## Question 10

How many positive integers  $n \leq 10$  satisfy  $\cos(n) > 0$ ?  
(Assume radian measure.)

# Solution 10

**Answer. 4.**

**Answer. 4.**

$$n \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{3\pi}{2}, \frac{5\pi}{2}\right)$$

**Answer. 4.**

$$n \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{3\pi}{2}, \frac{5\pi}{2}\right)$$

$$n \in (0, 1.57 \dots) \cup (4.71 \dots, 7.85 \dots)$$

**Answer. 4.**

$$n \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{3\pi}{2}, \frac{5\pi}{2}\right)$$

$$n \in (0, 1.57\dots) \cup (4.71\dots, 7.85\dots)$$

$$n \in \{1, 5, 6, 7\}$$

## Question 11

There are unique integers  $a$  and  $b$  for which

## Question 11

There are unique integers  $a$  and  $b$  for which

$$(2 - \sqrt{3})^3 = a + b\sqrt{3}.$$

What is  $a + b$ ?

# Solution 11

**Answer. 11.**



# Solution 11

**Answer.** 11. We have

$$(2 - \sqrt{3})^3 = 8 - 12\sqrt{3} + 6(\sqrt{3})^2 - (\sqrt{3})^3 = 26 - 15\sqrt{3}.$$

## Question 12

Simplify:

## Question 12

Simplify:

$$\frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5}}}}}$$

# Solution 12

**Answer.**  $\frac{17}{28}$ .

## Solution 12

**Answer.**  $\frac{17}{28}$ .

$$\blacktriangleright 1 + \frac{1}{5} = \frac{6}{5}$$

# Solution 12

**Answer.**  $\frac{17}{28}$ .

▶  $1 + \frac{1}{5} = \frac{6}{5}$

▶  $1 + \frac{1}{1 + \frac{1}{5}} = \frac{11}{6}$

# Solution 12

**Answer.**  $\frac{17}{28}$ .

$$\blacktriangleright 1 + \frac{1}{5} = \frac{6}{5}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{5}} = \frac{11}{6}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5}}} = \frac{17}{11}$$

# Solution 12

**Answer.**  $\frac{17}{28}$ .

$$\blacktriangleright 1 + \frac{1}{5} = \frac{6}{5}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{5}} = \frac{11}{6}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5}}} = \frac{17}{11}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5}}}} = \frac{28}{17}$$



# Solution 12

**Answer.**  $\frac{17}{28}$ .

$$\blacktriangleright 1 + \frac{1}{5} = \frac{6}{5}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{5}} = \frac{11}{6}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5}}} = \frac{17}{11}$$

$$\blacktriangleright 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5}}}} = \frac{28}{17}$$

Notice the pattern:  $\frac{6}{5}, \frac{11}{6}, \frac{17}{11}, \frac{28}{17}$

## Question 2-1

In 2019, Andrew Booker found the first known integer solution to  $a^3 + b^3 + c^3 = 33$ .

## Question 2-1

In 2019, Andrew Booker found the first known integer solution to  $a^3 + b^3 + c^3 = 33$ .

One digit in his solution has been replaced by a question mark.  
What is the missing digit?

## Question 2-1

In 2019, Andrew Booker found the first known integer solution to  $a^3 + b^3 + c^3 = 33$ .

One digit in his solution has been replaced by a question mark.  
What is the missing digit?

$$\begin{aligned} 33 &= 8866128975287528^3 \\ &\quad + (-877840544286223?)^3 \\ &\quad + (-2736111468807040)^3. \end{aligned}$$

**Answer. 9.**

## Solution 2-1

**Answer.** 9.

Modulo 10, we have

$$3 \equiv 8^3 + (-?)^3 + 0^3 \equiv 512 + (-?)^3,$$

## Solution 2-1

**Answer.** 9.

Modulo 10, we have

$$3 \equiv 8^3 + (-?)^3 + 0^3 \equiv 512 + (-?)^3,$$

so

$$1 \equiv (-?)^3,$$

**Answer.** 9.

Modulo 10, we have

$$3 \equiv 8^3 + (-?)^3 + 0^3 \equiv 512 + (-?)^3,$$

so

$$1 \equiv (-?)^3,$$

for which  $-? \equiv 1$  is the unique solution. So  $? = 9$ .



## Question 2-1

Solve for  $x$ :

$$\log_3(9x) + \log_9(3x) = 7$$

**Answer.** 27.

**Answer.** 27.

Rewrite the equation as

$$\log_3(9) + \log_3(x) + \log_9(3) + \frac{1}{2} \log_3(x) = 7,$$

## Solution 2-2

**Answer.** 27.

Rewrite the equation as

$$\log_3(9) + \log_3(x) + \log_9(3) + \frac{1}{2}\log_3(x) = 7,$$

or

$$\frac{5}{2} + \frac{3}{2}\log_3(x) = 7.$$

**Answer.** 27.

Rewrite the equation as

$$\log_3(9) + \log_3(x) + \log_9(3) + \frac{1}{2}\log_3(x) = 7,$$

or

$$\frac{5}{2} + \frac{3}{2}\log_3(x) = 7.$$

So

$$\frac{3}{2}\log_3(x) = 7 - \frac{5}{2} = \frac{9}{2},$$

and  $\log_3(x) = 3$ , so  $x = 27$ .

## Question 2-3

What is the smallest value of  $r$  for which the following is true?

## Question 2-3

What is the smallest value of  $r$  for which the following is true?

The parabola  $y = x^2$  intersects (in at least one point) the circle with center  $(0, 1)$  and radius  $r$ .

## Solution 2-3

**Answer.**  $\frac{\sqrt{3}}{2}$ .



## Solution 2-3

**Answer.**  $\frac{\sqrt{3}}{2}$ .

Solving  $y = x^2$  and  $x^2 + (y - 1)^2 = r^2$  yields

$$y^2 - y + (1 - r^2) = 0,$$

## Solution 2-3

**Answer.**  $\frac{\sqrt{3}}{2}$ .

Solving  $y = x^2$  and  $x^2 + (y - 1)^2 = r^2$  yields

$$y^2 - y + (1 - r^2) = 0,$$

$$y = \frac{1 \pm \sqrt{1 - 4(1 - r^2)}}{2},$$

## Solution 2-3

**Answer.**  $\frac{\sqrt{3}}{2}$ .

Solving  $y = x^2$  and  $x^2 + (y - 1)^2 = r^2$  yields

$$y^2 - y + (1 - r^2) = 0,$$

$$y = \frac{1 \pm \sqrt{1 - 4(1 - r^2)}}{2},$$

$$y = \frac{1 \pm \sqrt{-3 + 4r^2}}{2}.$$

## Solution 2-3

**Answer.**  $\frac{\sqrt{3}}{2}$ .

Solving  $y = x^2$  and  $x^2 + (y - 1)^2 = r^2$  yields

$$y^2 - y + (1 - r^2) = 0,$$

$$y = \frac{1 \pm \sqrt{1 - 4(1 - r^2)}}{2},$$

$$y = \frac{1 \pm \sqrt{-3 + 4r^2}}{2}.$$

This has a nonnegative solution when  $-3 + 4r^2 \geq 0$ , so when  $r^2 \geq \frac{3}{4}$ .

## Question 2-4

Your friend rolls two ordinary dice and you roll one.

## Question 2-4

Your friend rolls two ordinary dice and you roll one.  
What is the probability that your die roll exceeds the total of hers?

## Solution 2-4

**Answer.**  $\frac{5}{54}$ .

Depending on whether you roll 1, 2, 3, 4, 5, 6, the probability that her total is lower is respectively

$$0, 0, \frac{1}{36}, \frac{3}{36}, \frac{6}{36}, \frac{10}{36}.$$

So the overall probability is

$$\frac{1}{6} \left( \frac{1}{36} + \frac{3}{36} + \frac{6}{36} + \frac{10}{36} \right) = \frac{1}{6} \cdot \frac{20}{36} = \frac{20}{216} = \frac{5}{54}.$$

## Question 2-5

If

$$2 \cos^2(x) - \sin^2(x) = \frac{1}{2}$$

and  $0 < x < \frac{\pi}{2}$ , what is  $x$ ?



## Solution 2-5

**Answer.**  $\frac{\pi}{4}$ .

## Solution 2-5

**Answer.**  $\frac{\pi}{4}$ .

We have

$$3 \cos^2(x) = (2 \cos^2(x) - \sin^2(x)) + (\cos^2(x) + \sin^2(x)) = \frac{1}{2} + 1 = \frac{3}{2},$$

## Solution 2-5

**Answer.**  $\frac{\pi}{4}$ .

We have

$$3 \cos^2(x) = (2 \cos^2(x) - \sin^2(x)) + (\cos^2(x) + \sin^2(x)) = \frac{1}{2} + 1 = \frac{3}{2},$$

$$\text{so } \cos^2(x) = \frac{1}{2} \text{ and } \cos(x) = \frac{\sqrt{2}}{2}.$$

$$\text{So } x = \frac{\pi}{4}.$$

## Question 6

You eat a bunch of cupcakes. On the first day, you eat one cupcake; on the second day, you eat two cupcakes; on the third day, you eat three; and so on.

## Question 6

You eat a bunch of cupcakes. On the first day, you eat one cupcake; on the second day, you eat two cupcakes; on the third day, you eat three; and so on.

On what day will you eat your five thousandth cupcake?

# Solution 6

**Answer.** 100.

## Solution 6

**Answer.** 100.

After  $n$  days, you will have eaten

$$1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

cupcakes. So what is the minimal  $n$  for which

$$\frac{n(n+1)}{2} \geq 5000, \quad \text{or} \quad n(n+1) \geq 10000?$$

## Solution 6

**Answer.** 100.

After  $n$  days, you will have eaten

$$1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

cupcakes. So what is the minimal  $n$  for which

$$\frac{n(n+1)}{2} \geq 5000, \text{ or } n(n+1) \geq 10000?$$

Since  $10000 = 100^2$ , we have  $n = 100$ .



## Question 7

This is the logo of the Mathematical Association of America, a national organization that sponsors competitions and publishes excellent journals and books.



## Question 7

This is the logo of the Mathematical Association of America, a national organization that sponsors competitions and publishes excellent journals and books.



It depicts an icosahedron: a regular solid with twenty faces, each of which is an identical equilateral triangle.

## Question 7

This is the logo of the Mathematical Association of America, a national organization that sponsors competitions and publishes excellent journals and books.



It depicts an icosahedron: a regular solid with twenty faces, each of which is an identical equilateral triangle.

How many edges are not visible in the logo?

# Solution 7

**Answer.** 12.

# Solution 7

**Answer.** 12.

An icosahedron has 30 edges: 20 triangles times 3 edges per triangle, divided by 2 since each edge is shared between two triangles.

# Solution 7

**Answer.** 12.

An icosahedron has 30 edges: 20 triangles times 3 edges per triangle, divided by 2 since each edge is shared between two triangles.

You can count that 18 edges are visible in the picture, and  $30 - 18 = 12$ .

## Question 8

What is the smallest integer larger than

## Question 8

What is the smallest integer larger than

$$\log_2(33) + \log_{33}(2)?$$



# Solution 8

**Answer: 6.**

# Solution 8

**Answer:** 6.

We have

$$2^5 = 32, \quad 2^6 = 64,$$

so that  $\log_2(33)$  is slightly bigger than 5.

# Solution 8

**Answer:** 6.

We have

$$2^5 = 32, \quad 2^6 = 64,$$

so that  $\log_2(33)$  is slightly bigger than 5.

We have

$$\log_{33}(2) = \frac{1}{\log_2(33)} < \frac{1}{5}.$$

# Solution 8

**Answer:** 6.

We have

$$2^5 = 32, \quad 2^6 = 64,$$

so that  $\log_2(33)$  is slightly bigger than 5.

We have

$$\log_{33}(2) = \frac{1}{\log_2(33)} < \frac{1}{5}.$$

The sum of these numbers is less than 6.

## Question 9

How many real solutions  $x$  are there to the equation

## Question 9

How many real solutions  $x$  are there to the equation


$$x^5 + \sin(x) = 0?$$

# Solution 9

**Answer:** 1.

# Solution 9

**Answer:** 1.




The graphs of  $y = x^5$  and  $y = -\sin(x)$  don't intersect in  $(0, \pi)$  because of opposite signs, or in  $[\pi, \infty)$  because  $x^5 > 1$ . Similarly, there are no intersection points with  $x < 0$ .



## Solution 9

**Answer:** 1.



The graphs of  $y = x^5$  and  $y = -\sin(x)$  don't intersect in  $(0, \pi)$  because of opposite signs, or in  $[\pi, \infty)$  because  $x^5 > 1$ . Similarly, there are no intersection points with  $x < 0$ . So  $x = 0$  is the only intersection point.

## Question 10

You toss four coins. What is the probability that at least three of them come up heads?

# Solution 10

**Answer.**  $\frac{5}{16}$ .

## Solution 10

**Answer.**  $\frac{5}{16}$ .

There are  $2^4 = 16$  total ways to flip four coins.  
The total number with at least three heads is

$$\binom{4}{3} + \binom{4}{4} = 4 + 1 = 5,$$

## Solution 10


**Answer.**  $\frac{5}{16}$ .

There are  $2^4 = 16$  total ways to flip four coins.  
The total number with at least three heads is

$$\binom{4}{3} + \binom{4}{4} = 4 + 1 = 5,$$

*HHHH, HHHT, HHTH, HTHH, THHH.*

## Question 11



Given rectangle  $ABCD$  as above. If  $\angle AEB = 90^\circ$ , what is  $AE$ ?

# Solution 11

**Answer.**  $\frac{12}{5}$ .

# Solution 11

**Answer.**  $\frac{12}{5}$ .

$BD = 5$ , and  $\triangle ABE \sim \triangle BDC$ . So

$$\frac{AE}{AB} = \frac{BC}{BD} = \frac{3}{5}$$

and

$$AE = \frac{3}{5} \cdot AB = \frac{3}{5} \cdot 4 = \frac{12}{5}.$$



## Question 12

How many pairs of positive prime numbers  $p, q$  are there with

$$p - q = 21?$$

# Solution 12

**Answer. 1.**

# Solution 12

**Answer.** 1.

All prime numbers other than 2 are odd. The difference of two odd numbers is even. Therefore  $q$  must be 2. Since  $2 + 21 = 23$  is prime, there is one solution.

## Question 13

*In baseball, an **at bat** results in either a **hit** or an **out**. A player's **batting average** is their total number of hits divided by at bats, rounded off to the nearest thousandth.*

## Question 13

*In baseball, an **at bat** results in either a **hit** or an **out**. A player's **batting average** is their total number of hits divided by at bats, rounded off to the nearest thousandth.*

Five games into the baseball season, Cocky Gamecock has a batting average of .435. In his sixth game, he has five at bats and gets hits in all of them.

## Question 13

*In baseball, an **at bat** results in either a **hit** or an **out**. A player's **batting average** is their total number of hits divided by at bats, rounded off to the nearest thousandth.*

Five games into the baseball season, Cocky Gamecock has a batting average of .435. In his sixth game, he has five at bats and gets hits in all of them.

If this raises his batting average to .536, how many at bats does he have through his first six games?

# Solution 13

**Answer.** 28.

# Solution 13

**Answer.** 28.

Let  $x$  be the number of hits through 6 games, and  $y$  the number of at bats. Within a small roundoff error,

$$\frac{x - 5}{y - 5} = .435, \quad \frac{x}{y} = .536.$$



## Solution 13

**Answer.** 28.

Let  $x$  be the number of hits through 6 games, and  $y$  the number of at bats. Within a small roundoff error,

$$\frac{x - 5}{y - 5} = .435, \quad \frac{x}{y} = .536.$$

We have

$$x - 5 = .435(y - 5), \quad x = .435y + 5 - 2.175 = .435y + 2.825.$$

## Solution 13

**Answer.** 28.

Let  $x$  be the number of hits through 6 games, and  $y$  the number of at bats. Within a small roundoff error,

$$\frac{x - 5}{y - 5} = .435, \quad \frac{x}{y} = .536.$$

We have

$$x - 5 = .435(y - 5), \quad x = .435y + 5 - 2.175 = .435y + 2.825.$$

We thus have

$$x = .536y, \quad .101y = 2.825.$$

## Solution 13

**Answer.** 28.

Let  $x$  be the number of hits through 6 games, and  $y$  the number of at bats. Within a small roundoff error,

$$\frac{x - 5}{y - 5} = .435, \quad \frac{x}{y} = .536.$$

We have

$$x - 5 = .435(y - 5), \quad x = .435y + 5 - 2.175 = .435y + 2.825.$$

We thus have

$$x = .536y, \quad .101y = 2.825.$$

Thus, we have

$$y = \frac{2.825}{.101},$$

or  $y = 28$  up to the roundoff error.

## Question 14

If you write  $\frac{1}{2020}$  as an infinite repeating decimal,

## Question 14

If you write  $\frac{1}{2020}$  as an infinite repeating decimal, what is the sum of the first six digits after the decimal place?

# Solution 14

**Answer.** 18.

$$\frac{1}{2020} = 0.00049504950 \dots$$

# Solution 14

**Answer.** 18.

$$\frac{1}{2020} = 0.00049504950 \dots$$

Note that

$$\frac{1}{101} = .009900990099 \dots,$$

so

$$\frac{1}{1010} = .0009900990099 \dots,$$

$$\frac{1}{2020} = .0004950495049 \dots,$$