

Error Terms in Arithmetic Statistics


Frank Thorne


University of South Carolina


Chennai Mathematical Institute, November 25, 2025


`thornef.github.io/cmi-2025.pdf`











Sample Theorem 1: Counting Cubic Fields

What is “arithmetic statistics”?

Sample Theorem 1: Counting Cubic Fields

What is “arithmetic statistics”?

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\mathrm{Disc}(K)| < X\}.$$

Sample Theorem 1: Counting Cubic Fields

What is “arithmetic statistics”?

For any integer $d \geq 1$, write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\mathrm{Disc}(K)| < X\}.$$

Theorem (Davenport-Heilbronn)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + o(X).$$

Sample Theorem 2: Counting Quartic and Quintic Fields

Theorem (Bhargava)

We have

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4}) X,$$

Sample Theorem 2: Counting Quartic and Quintic Fields

Theorem (Bhargava)

We have

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

$$N_5(X) \sim \frac{13}{130} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})X.$$

Sample Theorem 3: 3-torsion in Quadratic Class Groups

Theorem (Davenport-Heilbronn)

We have

$$\sum_{|D| < X} \#|\mathrm{Cl}(\mathbb{Q}(\sqrt{D}))[3]| = \frac{3 + 3 + 1 + 3}{\pi^2} X + o(X).$$

Sample Theorem 4: 2-Selmer Groups in Elliptic Curves

Theorem (Bhargava-Shankar)

*When elliptic curves E are ordered by **height**, the average size of their **2-Selmer groups** is **3**.*

Sample Theorem 4: 2-Selmer Groups in Elliptic Curves

Theorem (Bhargava-Shankar)

*When elliptic curves E are ordered by **height**, the average size of their **2-Selmer groups** is **3**.*

Corollary

*Their average **rank** is at most 1.5.*

Parametrization: The Basic Metatheorem

Theorem

There exists an explicit, “nice” bijection

$$\{ \text{Something nice} \} \longleftrightarrow G(\mathbb{Z}) \backslash V(\mathbb{Z})$$

where V is a f.d. representation of an algebraic group G .

Parametrization: The Basic Metatheorem

Theorem

There exists an explicit, “nice” bijection

$$\{ \text{Something nice} \} \longleftrightarrow G(\mathbb{Z}) \backslash V(\mathbb{Z})$$

where V is a f.d. representation of an algebraic group G .

Moreover, **certain arithmetic properties** on the left correspond to **congruence conditions** on the right.

Example: Binary Cubic Forms

Let V be the space of binary cubic forms:

$$V := \{x(u, v) = au^3 + bu^2v + cuv^2 + dv^3\}.$$

Example: Binary Cubic Forms

Let V be the space of binary cubic forms:

$$V := \{x(u, v) = au^3 + bu^2v + cuv^2 + dv^3\}.$$

$G = \mathrm{GL}_2$ acts on V by

$$(gx)(u, v) = \frac{1}{\det g} x((u, v)g).$$

Example: Binary Cubic Forms

Let V be the space of binary cubic forms:

$$V := \{x(u, v) = au^3 + bu^2v + cuv^2 + dv^3\}.$$

$G = \mathrm{GL}_2$ acts on V by

$$(gx)(u, v) = \frac{1}{\det g} x((u, v)g).$$

We have

$$\mathrm{Disc}(x) = b^2c^2 - 4ac^3 - 4b^3d - 27a^2d^2 + 18abcd,$$

for which we have

Example: Binary Cubic Forms

Let V be the space of binary cubic forms:

$$V := \{x(u, v) = au^3 + bu^2v + cuv^2 + dv^3\}.$$

$G = \mathrm{GL}_2$ acts on V by

$$(gx)(u, v) = \frac{1}{\det g} x((u, v)g).$$

We have

$$\mathrm{Disc}(x) = b^2c^2 - 4ac^3 - 4b^3d - 27a^2d^2 + 18abcd,$$

for which we have

$$\blacktriangleright \mathrm{Disc}(gx) = (\det g)^2 \mathrm{Disc}(x);$$

Example: Binary Cubic Forms

Let V be the space of binary cubic forms:

$$V := \{x(u, v) = au^3 + bu^2v + cuv^2 + dv^3\}.$$

$G = \mathrm{GL}_2$ acts on V by

$$(gx)(u, v) = \frac{1}{\det g} x((u, v)g).$$

We have

$$\mathrm{Disc}(x) = b^2c^2 - 4ac^3 - 4b^3d - 27a^2d^2 + 18abcd,$$

for which we have

- ▶ $\mathrm{Disc}(gx) = (\det g)^2 \mathrm{Disc}(x)$;
- ▶ $\mathrm{Disc}(x) = 0$ if and only if $x(u, v)$ has a repeated root.

Example: Binary Cubic Forms (2)

Theorem (Levi, Delone-Faddeev, Gan-Gross-Savin)

$G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ parametrize *cubic rings*. Further, if $v \leftrightarrow R$,

Example: Binary Cubic Forms (2)

Theorem (Levi, Delone-Faddeev, Gan-Gross-Savin)

$G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ parametrize *cubic rings*. Further, if $v \leftrightarrow R$,

- ▶ $\text{Stab}(v)$ is isomorphic to $\text{Aut}(R)$;

Example: Binary Cubic Forms (2)

Theorem (Levi, Delone-Faddeev, Gan-Gross-Savin)

$G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ parametrize *cubic rings*. Further, if $v \leftrightarrow R$,

- ▶ $\text{Stab}(v)$ is isomorphic to $\text{Aut}(R)$;
- ▶ $\text{Disc}(v) = \text{Disc}(R)$;

Example: Binary Cubic Forms (2)

Theorem (Levi, Delone-Faddeev, Gan-Gross-Savin)

$G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$ parametrize *cubic rings*. Further, if $v \leftrightarrow R$,

- ▶ $\text{Stab}(v)$ is isomorphic to $\text{Aut}(R)$;
- ▶ $\text{Disc}(v) = \text{Disc}(R)$;
- ▶ (Davenport-Heilbronn) R is *maximal* iff, for all primes p , v satisfies a certain congruence condition (mod p^2).

Theorem (DHBBPBSTTTBTT)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1+\sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{\frac{2}{3}}(\log X)^3).$$

Some parametrizations

Some parametrizations

- ▶ $V = \text{Sym}^3(2)$, $G = \text{GL}_2$: cubic rings; 3-torsion in class groups
(Levi, Delone-Faddeev, Davenport-Heilbronn)

Some parametrizations

- ▶ $V = \text{Sym}^3(2)$, $G = \text{GL}_2$: cubic rings; 3-torsion in class groups
(Levi, Delone-Faddeev, Davenport-Heilbronn)
- ▶ $V = 2 \otimes \text{Sym}^2(3)$, $G = \text{GL}_2 \times \text{GL}_3$.
quartic rings; 2-torsion in class groups of cubic fields
(Wright-Yukie; Bhargava)

Some parametrizations

- ▶ $V = \operatorname{Sym}^3(2)$, $G = \operatorname{GL}_2$: cubic rings; 3-torsion in class groups
(Levi, Delone-Faddeev, Davenport-Heilbronn)
- ▶ $V = 2 \otimes \operatorname{Sym}^2(3)$, $G = \operatorname{GL}_2 \times \operatorname{GL}_3$.
quartic rings; 2-torsion in class groups of cubic fields
(Wright-Yukie; Bhargava)
- ▶ $V = 4 \otimes \wedge^2(5)$, $G = \operatorname{GL}_4 \times \operatorname{SL}_5$
quintic rings (Wright-Yukie; Bhargava)

Some parametrizations

- ▶ $V = \text{Sym}^3(2)$, $G = \text{GL}_2$: cubic rings; 3-torsion in class groups
(Levi, Delone-Faddeev, Davenport-Heilbronn)
- ▶ $V = 2 \otimes \text{Sym}^2(3)$, $G = \text{GL}_2 \times \text{GL}_3$.
quartic rings; 2-torsion in class groups of cubic fields
(Wright-Yukie; Bhargava)
- ▶ $V = 4 \otimes \wedge^2(5)$, $G = \text{GL}_4 \times \text{SL}_5$
quintic rings (Wright-Yukie; Bhargava)
- ▶ $V = \text{Sym}^4(2)$, $G = \text{PGL}_2$
2-Selmer elements of elliptic curves
(Birch-Swinnerton-Dyer, Bhargava-Shankar)

Some parametrizations

- ▶ $V = \text{Sym}^3(2)$, $G = \text{GL}_2$: cubic rings; 3-torsion in class groups
(Levi, Delone-Faddeev, Davenport-Heilbronn)
- ▶ $V = 2 \otimes \text{Sym}^2(3)$, $G = \text{GL}_2 \times \text{GL}_3$.
quartic rings; 2-torsion in class groups of cubic fields
(Wright-Yukie; Bhargava)
- ▶ $V = 4 \otimes \wedge^2(5)$, $G = \text{GL}_4 \times \text{SL}_5$
quintic rings (Wright-Yukie; Bhargava)
- ▶ $V = \text{Sym}^4(2)$, $G = \text{PGL}_2$
2-Selmer elements of elliptic curves
(Birch-Swinnerton-Dyer, Bhargava-Shankar)
- ▶ ... and more!
(Bhargava, Ho, Shankar, Varma, X. Wang, Wood,

More Interesting Parametrizations

56

MANJUL BHARGAVA

Table 1: Summary of Higher Composition Laws

#	Lattice ($V_{\mathbb{Z}}$)	Group acting ($G_{\mathbb{Z}}$)	Parametrizes (\mathcal{C})	(k)	(n)	(H)
1.	$\{0\}$	-	Linear rings	0	0	A_0
2.	$\tilde{\mathbb{Z}}$	$\mathrm{SL}_1(\mathbb{Z})$	Quadratic rings	1	1	A_1
3.	$(\mathrm{Sym}^2 \mathbb{Z}^2)^*$ (GAUSS'S LAW)	$\mathrm{SL}_2(\mathbb{Z})$	Ideal classes in quadratic rings	2	3	B_2
4.	$\mathrm{Sym}^3 \mathbb{Z}^2$	$\mathrm{SL}_2(\mathbb{Z})$	Order 3 ideal classes in quadratic rings	4	4	G_2
5.	$\mathbb{Z}^2 \otimes \mathrm{Sym}^2 \mathbb{Z}^2$	$\mathrm{SL}_2(\mathbb{Z})^2$	Ideal classes in quadratic rings	4	6	B_3
6.	$\mathbb{Z}^2 \otimes \mathbb{Z}^2 \otimes \mathbb{Z}^2$	$\mathrm{SL}_2(\mathbb{Z})^3$	Pairs of ideal classes in quadratic rings	4	8	D_4
7.	$\mathbb{Z}^2 \otimes \wedge^2 \mathbb{Z}^4$	$\mathrm{SL}_2(\mathbb{Z}) \times \mathrm{SL}_4(\mathbb{Z})$	Ideal classes in quadratic rings	4	12	D_5
8.	$\wedge^3 \mathbb{Z}^6$	$\mathrm{SL}_6(\mathbb{Z})$	Quadratic rings	4	20	E_6
9.	$(\mathrm{Sym}^3 \mathbb{Z}^2)^*$	$\mathrm{GL}_2(\mathbb{Z})$	Cubic rings	4	4	G_2
10.	$\mathbb{Z}^2 \otimes \mathrm{Sym}^2 \mathbb{Z}^3$	$\mathrm{GL}_2(\mathbb{Z}) \times \mathrm{SL}_3(\mathbb{Z})$	Order 2 ideal classes in cubic rings	12	12	F_4
11.	$\mathbb{Z}^2 \otimes \mathbb{Z}^3 \otimes \mathbb{Z}^3$	$\mathrm{GL}_2(\mathbb{Z}) \times \mathrm{SL}_3(\mathbb{Z})^2$	Ideal classes in cubic rings	12	18	E_6
12.	$\mathbb{Z}^2 \otimes \wedge^2 \mathbb{Z}^6$	$\mathrm{GL}_2(\mathbb{Z}) \times \mathrm{SL}_6(\mathbb{Z})$	Cubic rings	12	30	E_7
13.	$(\mathbb{Z}^2 \otimes \mathrm{Sym}^2 \mathbb{Z}^3)^*$	$\mathrm{GL}_2(\mathbb{Z}) \times \mathrm{SL}_3(\mathbb{Z})$	Quartic rings	12	12	F_4
14.	$\mathbb{Z}^4 \otimes \wedge^2 \mathbb{Z}^5$	$\mathrm{GL}_4(\mathbb{Z}) \times \mathrm{SL}_5(\mathbb{Z})$	Quintic rings	40	40	E_8

Bhargava, *Higher composition laws IV*, Ann. Math., 2008



Still More Interesting Parametrizations

	Group (s.s.)	Representation	Geometric Data	Invariants	Dynkin	§
1.	SL_2	$\mathrm{Sym}^4(2)$	(C, L_2)	2, 3	$A_2^{(2)}$	4.1
2.	SL_2^2	$\mathrm{Sym}^2(2) \otimes \mathrm{Sym}^2(2)$	$(C, L_2, L'_2) \sim (C, L_2, P)$	2, 3, 4	$D_3^{(2)}$	6.1
3.	SL_2^4	$2 \otimes 2 \otimes 2 \otimes 2$	$(C, L_2, L'_2, L''_2) \sim (C, L_2, P, P')$	2, 4, 4, 6	$D_4^{(1)}$	6.2
4.	SL_2^3	$2 \otimes 2 \otimes \mathrm{Sym}^2(2)$	$(C, L_2, L'_2) \sim (C, L_2, P)$	2, 4, 6	$E_3^{(1)}$	6.3.1
5.	SL_2^2	$\mathrm{Sym}^2(2) \otimes \mathrm{Sym}^2(2)$	$(C, L_2, L'_2) \sim (C, L_2, P)$	2, 3, 4	$D_3^{(2)}$	6.3.3
6.	SL_2^2	$2 \otimes \mathrm{Sym}^3(2)$	(C, L_2, P_3)	2, 6	$G_2^{(1)}$	6.3.2
7.	SL_2	$\mathrm{Sym}^4(2)$	(C, L_2, P_3)	2, 3	$A_2^{(2)}$	6.3.4
8.	$\mathrm{SL}_2^2 \times \mathrm{GL}_4$	$2 \otimes 2 \otimes \wedge^2(4)$	$(C, L_2, M_{2,6})$	2, 4, 6, 8	$D_5^{(1)}$	6.6.1
9.	$\mathrm{SL}_2 \times \mathrm{SL}_6$	$2 \otimes \wedge^3(6)$	$(C, L_2, M_{3,6})$ with $L^{\otimes 3} \cong \det M$	2, 6, 8, 12	$E_6^{(1)}$	6.6.2
10.	$\mathrm{SL}_2 \times \mathrm{Sp}_6$	$2 \otimes \wedge_0^3(6)$	$(C, L_2, (M_{3,6}, \varphi))$ with $L^{\otimes 3} \cong \det M$	2, 6, 8, 12	$E_6^{(2)}$	6.6.3
11.	$\mathrm{SL}_2 \times \mathrm{Spin}_{12}$	$2 \otimes S^+(32)$	$(C \rightarrow \mathbb{P}^1(\mathcal{H}_3(\mathbb{H})), L_2)$	2, 6, 8, 12	$E_7^{(1)}$	6.6.3
12.	$\mathrm{SL}_2 \times E_7$	$2 \otimes 56$	$(C \rightarrow \mathbb{P}^1(\mathcal{H}_3(\mathbb{O})), L_2)$	2, 6, 8, 12	$E_8^{(1)}$	6.6.3
13.	SL_3	$\mathrm{Sym}^3(3)$	(C, L_3)	4, 6	$D_4^{(3)}$	4.2
14.	SL_3^3	$3 \otimes 3 \otimes 3$	$(C, L_3, L'_3) \sim (C, L_3, P)$	6, 9, 12	$E_6^{(1)}$	5.1
15.	SL_3^2	$3 \otimes \mathrm{Sym}^2(3)$	(C, L_3, P_2)	6, 12	$F_4^{(1)}$	5.2.1
16.	SL_3	$\mathrm{Sym}^3(3)$	(C, L_3, P_2)	4, 6	$D_4^{(3)}$	5.2.2
17.	$\mathrm{SL}_3 \times \mathrm{SL}_6$	$3 \otimes \wedge^2(6)$	$(C, L_3, M_{2,6})$ with $L^{\otimes 2} \cong \det M$	6, 12, 18	$E_7^{(1)}$	5.5
18.	$\mathrm{SL}_3 \times E_6$	$3 \otimes 27$	$(C \hookrightarrow \mathbb{P}^2(\mathbb{O}), L_3)$	6, 12, 18	$E_8^{(1)}$	5.4
19.	$\mathrm{SL}_2 \times \mathrm{SL}_4$	$2 \otimes \mathrm{Sym}^2(4)$	(C, L_4)	8, 12	$E_6^{(2)}$	4.3
20.	$\mathrm{SL}_5 \times \mathrm{SL}_5$	$\wedge^2(5) \otimes 5$	(C, L_5)	20, 30	$E_8^{(1)}$	4.4

Table 1: Table of coregular representations and their moduli interpretations

Bhargava and Ho, *Coregular spaces and genus one curves*, Camb. J. Math.

Problem: Want to count $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$, where:

Problem: Want to count $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$, where:

- ▶ (G, V) is a “nice” f.d. representation of an algebraic group;

Problem: Want to count $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$, where:

- ▶ (G, V) is a “nice” f.d. representation of an algebraic group;
- ▶ We require $|\text{Disc}(v)| < X$ for $v \in V(\mathbb{Z})$;

Problem: Want to count $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})$, where:

- ▶ (G, V) is a “nice” f.d. representation of an algebraic group;
- ▶ We require $|\text{Disc}(v)| < X$ for $v \in V(\mathbb{Z})$;
- ▶ We might impose congruence conditions as well.

Five Lattice Point Counting Methods

- ▶ Geometry of Numbers (Gauss, Davenport-Heilbronn, ...)

Five Lattice Point Counting Methods

- ▶ Geometry of Numbers (Gauss, Davenport-Heilbronn, ...) General and very widely applicable.

Five Lattice Point Counting Methods

- ▶ Geometry of Numbers (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.
- ▶ **“Simple Harmonic Analysis”** (Belabas-Fouvry, Taniguchi-T., ...)

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.
- ▶ **“Simple Harmonic Analysis”** (Belabas-Fouvry, Taniguchi-T., ...)
Very effective for lower bound sieves.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.
- ▶ **“Simple Harmonic Analysis”** (Belabas-Fouvry, Taniguchi-T., ...)
Very effective for lower bound sieves.
Not designed to produce asymptotics.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.
- ▶ **“Simple Harmonic Analysis”** (Belabas-Fouvry, Taniguchi-T., ...)
Very effective for lower bound sieves.
Not designed to produce asymptotics.
- ▶ **Averaging with Fourier analysis** (Anderson-Bhargava-T.)

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.
- ▶ **“Simple Harmonic Analysis”** (Belabas-Fouvry, Taniguchi-T., ...)
Very effective for lower bound sieves.
Not designed to produce asymptotics.
- ▶ **Averaging with Fourier analysis** (Anderson-Bhargava-T.)
Quantitative improvements in Bhargava’s averaging method.

Five Lattice Point Counting Methods

- ▶ **Geometry of Numbers** (Gauss, Davenport-Heilbronn, ...)
General and very widely applicable.
Does not deal well with cusps.
- ▶ **Zeta Functions** (Sato-Shintani, Datskovsky-Wright, Yukie, ...)
Expect best possible results when everything is in place.
Underlying theory is difficult.
- ▶ **The Averaging Technique** (Bhargava)
Naturally separates the main body from the cusp.
Inherits some limitations from the geometry of numbers.
- ▶ **“Simple Harmonic Analysis”** (Belabas-Fouvry, Taniguchi-T., ...)
Very effective for lower bound sieves.
Not designed to produce asymptotics.
- ▶ **Averaging with Fourier analysis** (Anderson-Bhargava-T.)
Quantitative improvements in Bhargava’s averaging method.
Expect it to work in the same situations.

An explicit evaluation

Theorem (Taniguchi-T., 2011)

We have

$$\widehat{\Phi_{p^2}}(v) = \begin{cases} p^{-2} + p^{-3} - p^{-5} & v/p : \text{of type } (0), \\ p^{-3} - p^{-5} & v/p : \text{of type } (1^3), (1^2 1), \\ -p^{-5} & v/p : \text{of type } (111), (21), (3). \\ p^{-3} - p^{-5} & v : \text{of type } (1_{**}^3), \\ -p^{-5} & v : \text{of type } (1_*^3), (1_{\max}^3), \\ 0 & \text{otherwise.} \end{cases}$$

An explicit evaluation

Theorem (Taniguchi-T., 2011)

We have

$$\widehat{\Phi}_{p^2}(v) = \begin{cases} p^{-2} + p^{-3} - p^{-5} & v/p : \text{of type } (0), \\ p^{-3} - p^{-5} & v/p : \text{of type } (1^3), (1^2 1), \\ -p^{-5} & v/p : \text{of type } (111), (21), (3). \\ p^{-3} - p^{-5} & v : \text{of type } (1_{**}^3), \\ -p^{-5} & v : \text{of type } (1_*^3), (1_{\max}^3), \\ 0 & \text{otherwise.} \end{cases}$$

So:

$$\frac{1}{p^8} \sum_{v \in V(\mathbb{Z}/p^2\mathbb{Z})} |\widehat{\Phi}_{p^2}(v)| \ll p^{-7}.$$

Theorem (DHBBPBSTTTBTT)

We have

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{\frac{3}{5}+\epsilon}) + O(X^{1-\frac{1}{8-7+2}+\epsilon}).$$

Application 1: Direct products

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Application 1: Direct products

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

Application 1: Direct products

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A) X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

$$\mathrm{Disc}(KL) = * \cdot \mathrm{Disc}(K)^{|A|} \mathrm{Disc}(L)^d,$$

Application 1: Direct products

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A) X^{1/|A|}.$$

Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

where $*$ is divisible only primes **ramified in both K and L** .

Application 1: Direct products

Theorem (Wang, Masri-T.-Tsai-Wang)

Let $d \in \{3, 4, 5\}$ and let A be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A) X^{1/|A|}.$$


Idea: If K is an S_d -field and L is an A -field, then **usually** K and L are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$


where $*$ is divisible only primes **ramified in both K and L** .

(This doesn't happen too often.)

Application 2, Counting by other invariants: S_3 -cubic fields



Application 2, Counting by other invariants: S_3 -cubic fields




We have


$$\text{Disc}(K) = D(K)F(K)^2,$$

where

$$D(K) := \text{Disc}(\mathbb{Q}(\sqrt{D})), \quad F(K) = f(K(\sqrt{D})/\mathbb{Q}(\sqrt{D})).$$

S_3 -sextic fields





We have

$$\text{Disc}(K(\sqrt{D})) = D(K)^3 F(K)^4,$$

where

$$D(K) := \text{Disc}(\mathbb{Q}(\sqrt{D})), \quad F(K) = f(K(\sqrt{D})/\mathbb{Q}(\sqrt{D})).$$

Definition

For $\alpha, \beta > 0$, define an expression of the form

$$I(\alpha, \beta) := |D|^\alpha F^\beta$$

to be a **generalized discriminant**, and

Definition

For $\alpha, \beta > 0$, define an expression of the form

$$I(\alpha, \beta) := |D|^{\alpha} F^{\beta}$$

to be a **generalized discriminant**, and

$$N_I(X) := \#\{K : [K : \mathbb{Q}] = 3, I(K) < X\}$$

the associated counting function.

Sample counting results

Sample counting results

- ▶ $(\alpha = 1, \beta = 2)$:

$$I(\alpha, \beta) = |D|F^2$$

is the **usual (absolute) discriminant**, and

$$N_{|D|F^2}(X) = \frac{1}{3\zeta(3)}X + o(X)$$

is the Davenport-Heilbronn theorem.

Sample counting results

- ▶ $(\alpha = 1, \beta = 2)$ [Davenport-Heilbronn]

Sample counting results

- ▶ $(\alpha = 1, \beta = 2)$ [Davenport-Heilbronn]
- ▶ $(\alpha = 3, \beta = 4)$:

$$I(\alpha, \beta) = |D|^3 F^4$$

is the **absolute discriminant of the associated S_3 -sextic**, and

$$N_{|D|^3 F^4}(X) = C \cdot X^{1/3} + o(X^{1/3})$$

by Belabas-Fouvry and Bhargava-Wood.

Sample counting results

- ▶ $(\alpha = 1, \beta = 2)$ [Davenport-Heilbronn]
- ▶ $(\alpha = 3, \beta = 4)$:

$$I(\alpha, \beta) = |D|^3 F^4$$

is the **absolute discriminant of the associated S_3 -sextic**, and

$$N_{|D|^3 F^4}(X) = C \cdot X^{1/3} + o(X^{1/3})$$

by Belabas-Fouvry and Bhargava-Wood.

(The C_3 -cubics contribute a negligible amount.)

Sample counting results

- ▶ $(\alpha = 1, \beta = 2)$ [Davenport-Heilbronn]
- ▶ $(\alpha = 3, \beta = 4)$:

$$I(\alpha, \beta) = |D|^3 F^4$$

is the **absolute discriminant of the associated S_3 -sextic**, and

$$N_{|D|^3 F^4}(X) = C \cdot X^{1/3} + o(X^{1/3})$$

by Belabas-Fouvry and Bhargava-Wood.

(The C_3 -cubics contribute a negligible amount.)

- ▶ $(\alpha = 1, \beta = 1)$:

$$I(\alpha, \beta) = |D|F$$

is the **squarefree part of the discriminant**,

Sample counting results

- ▶ $(\alpha = 1, \beta = 2)$ [Davenport-Heilbronn]
- ▶ $(\alpha = 3, \beta = 4)$:

$$I(\alpha, \beta) = |D|^3 F^4$$

is the **absolute discriminant of the associated S_3 -sextic**, and

$$N_{|D|^3 F^4}(X) = C \cdot X^{1/3} + o(X^{1/3})$$

by Belabas-Fouvry and Bhargava-Wood.

(The C_3 -cubics contribute a negligible amount.)

- ▶ $(\alpha = 1, \beta = 1)$:

$$I(\alpha, \beta) = |D|F$$

is the **squarefree part of the discriminant**, up to 2- and 3-adic factors.

Counting by squarefree part

Let $N_C(X)$ count cubic fields where the **squarefree part of the discriminant** is $< X$.

Counting by squarefree part

Let $N_C(X)$ count cubic fields where the **squarefree part of the discriminant** is $< X$.

Theorem (Shankar-T.)

We have

$$N_C(X) = \frac{7}{5} \prod_p \left(1 + \frac{2}{p}\right) \left(1 - \frac{1}{p}\right)^2 X \log X + o(X \log X).$$

Overview of the proof

We must be able to count cubic fields with:

Overview of the proof

We must be able to count cubic fields with:

- ▶ Fixed F and bounded D ;

Overview of the proof

We must be able to count cubic fields with:

- ▶ Fixed F and bounded D ;
- ▶ Fixed D and bounded F .

Overview of the proof

We must be able to count cubic fields with:

- ▶ Fixed F and bounded D ;
- ▶ Fixed D and bounded F .

Goal: Prove that, excluding $D = -3$,

$$\#\{K \in \mathcal{F}(\Sigma) : |D(K)| < Y, F(K) < Z\} \sim C(\Sigma) \cdot Y \cdot Z,$$

Overview of the proof

We must be able to count cubic fields with:

- ▶ Fixed F and bounded D ;
- ▶ Fixed D and bounded F .

Goal: Prove that, excluding $D = -3$,

$$\#\{K \in \mathcal{F}(\Sigma) : |D(K)| < Y, F(K) < Z\} \sim C(\Sigma) \cdot Y \cdot Z,$$

where

$$C(\Sigma) = \frac{1}{2\alpha} \left(\sum_{K \in \Sigma_\infty} \frac{1}{|\text{Aut}(K)|} \right) \prod_p \left(\sum_{K \in \Sigma_p} \frac{|D(K)|_p |F(K)|_p}{|\text{Aut}(K)|} \right) \left(1 - \frac{1}{p} \right)^2.$$

Also:

$$\#\{K \in \mathcal{F}(\Sigma) : D(K) = -3, F(K) < Z\} \sim C(-3, \Sigma) \cdot Z \log Z.$$

Method 1: Uniform Davenport-Heilbronn

Define

$$N(\mathcal{F}(\Sigma)^{(f)}; Y) := \#\{K \in \mathcal{F}(\Sigma) : F(K) = f, |D(K)| < Y\},$$

Theorem (Davenport-Heilbronn, ...*, Bhargava-Taniguchi-T.)

We have

$$N(\mathcal{F}(\Sigma)^{(f)}; Y) = C_1(\Sigma, f) \cdot Y + C_2(\Sigma, f) \cdot Y^{5/6} + O(E(Y; f, \Sigma)),$$

with the 'average error bound'

$$\sum_{f \leq F} E(Y_f; f, \Sigma) \ll_{\epsilon} Y^{2/3+\epsilon} F^{4/3+\epsilon} P_{\Sigma}^{2/3},$$

uniformly in F and $Y_f \leq Y$.

*: Belabas, Belabas-Bhargava-Pomerance, Bhargava-Shankar-Tsimerman, Taniguchi-T.

Explicit Cohen-Morra

For each nonzero fundamental discriminant d , define a Dirichlet series

$$\Phi_{\Sigma,d}(s) := \frac{1}{2} + \sum_{K \in \mathcal{F}(\Sigma): \mathcal{D}(K)=d} \frac{1}{F(K)^s},$$

Theorem (Cohen-Morra, Cohen-T.)

For Σ not specifying any restriction, we have

$$\Phi_{\Sigma,d}(s) = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{2}{3^{2s}}\right) \prod_{\left(\frac{-3d}{p}\right)=1} \left(1 + \frac{2}{p^s}\right) + \sum_{L \in \mathcal{L}_3(d)} M_{3,L}(s) \prod_{\left(\frac{-3d}{p}\right)=1} \left(1 + \frac{\omega_L(p)}{p^s}\right),$$

where

$$M_{3,L}(s) := \begin{cases} 1 - 3^{-2s} & : \text{Disc}(L) = -27d \\ 1 + 2 \cdot 3^{-2s} & : \text{Disc}(L) = -3d, \end{cases}$$

Explicit Cohen-Morra

For each nonzero fundamental discriminant d , define a Dirichlet series

$$\Phi_{\Sigma,d}(s) := \frac{1}{2} + \sum_{K \in \mathcal{F}(\Sigma): \mathcal{D}(K)=d} \frac{1}{F(K)^s},$$

Theorem (Cohen-Morra, Cohen-T.)

For Σ not specifying any restriction, we have

$$\Phi_{\Sigma,d}(s) = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{2}{3^{2s}}\right) \prod_{\left(\frac{-3d}{p}\right)=1} \left(1 + \frac{2}{p^s}\right) + \sum_{L \in \mathcal{L}_3(d)} M_{3,L}(s) \prod_{\left(\frac{-3d}{p}\right)=1} \left(1 + \frac{\omega_L(p)}{p^s}\right),$$

where

$$M_{3,L}(s) := \begin{cases} 1 - 3^{-2s} & : \text{Disc}(L) = -27d \\ 1 + 2 \cdot 3^{-2s} & : \text{Disc}(L) = -3d, \end{cases}$$

- $\mathcal{L}_3(d)$ is the set of cubic fields of discriminant $-3d$ or $-27d$;

Explicit Cohen-Morra

For each nonzero fundamental discriminant d , define a Dirichlet series

$$\Phi_{\Sigma,d}(s) := \frac{1}{2} + \sum_{K \in \mathcal{F}(\Sigma): \mathcal{D}(K)=d} \frac{1}{F(K)^s},$$

Theorem (Cohen-Morra, Cohen-T.)

For Σ not specifying any restriction, we have

$$\Phi_{\Sigma,d}(s) = -\frac{1}{2} + \frac{1}{2} \left(1 + \frac{2}{3^{2s}}\right) \prod_{\left(\frac{-3d}{p}\right)=1} \left(1 + \frac{2}{p^s}\right) + \sum_{L \in \mathcal{L}_3(d)} M_{3,L}(s) \prod_{\left(\frac{-3d}{p}\right)=1} \left(1 + \frac{\omega_L(p)}{p^s}\right),$$

where

$$M_{3,L}(s) := \begin{cases} 1 - 3^{-2s} & : \text{Disc}(L) = -27d \\ 1 + 2 \cdot 3^{-2s} & : \text{Disc}(L) = -3d, \end{cases}$$

► $\mathcal{L}_3(d)$ is the set of cubic fields of discriminant $-3d$ or $-27d$;

►

$$\omega_L(p) = \begin{cases} -1 & \text{if } p \text{ is inert in } L, \\ 2 & \text{if } p \text{ is totally split in } L, \\ 0 & \text{otherwise.} \end{cases}$$

Analytic Consequence of Cohen-Morra

Define

$$N(\mathcal{F}(\Sigma)_d; Z) := \#\{K \in \mathcal{F}(\Sigma) : D(K) = d, F(K) < Z\}.$$

Then:

Theorem (Cohen-Morra)

For $\Sigma = \Sigma_{all}$ and $d \neq -3$, we have

$$N(\mathcal{F}(\Sigma)_d; Z) = \text{Res}_{s=1}(\Phi_{\Sigma,d}(s)) \cdot Z + O_{\epsilon}(|\mathcal{L}_3(d)||d|^{1/6} Z^{2/3+\epsilon}).$$

Analytic Consequence of Cohen-Morra

Define

$$N(\mathcal{F}(\Sigma)_d; Z) := \#\{K \in \mathcal{F}(\Sigma) : D(K) = d, F(K) < Z\}.$$

Then:

Theorem (Cohen-Morra)

For $\Sigma = \Sigma_{all}$ and $d \neq -3$, we have

$$N(\mathcal{F}(\Sigma)_d; Z) = \text{Res}_{s=1}(\Phi_{\Sigma,d}(s)) \cdot Z + O_{\epsilon}(|\mathcal{L}_3(d)||d|^{1/6} Z^{2/3+\epsilon}).$$

Proved variations of the above permitting **local conditions**.